

Guava / per acre



Feb–Mar sowing Multi-nutrient (K-Ca-Mg-B) 20 rows · 6 stages

	September	October, flowering starts	November, fruiting	December	January	February. Harvest
POTASH	 V-Potash Plus N 10% · K ₂ O 40% · Mg · B · chelated Fe 10 kg Fertigation 10 kg	 VL-Potash Liquid Potash 30% 1 pack Foliar, 1 L per 100 L water 1 L		 VL-Potash Liquid Potash 30% 1 pack Foliar, 1 L per 100 L water 1 L		
SECONDARY NUTRIENT	 Cala-Mag V N 10% · Ca 6% · Mg 4% · B 1% 1 pack Fertigation, or spray at 500 ml per 100 L water 1 L	 Cala-Mag V N 10% · Ca 6% · Mg 4% · B 1% 1 pack Fertigation. Repeat 15 days after flowering 1 L	 Cala-Mag V N 10% · Ca 6% · Mg 4% · B 1% 1 pack Fertigation. Repeat 15 days after fruiting 1 L			
	 MgSO₄ Magnesium Sulfate 5 kg Fertigation					
SOIL AMENDMENT	 Well-rotten farmyard manure 3-5 trolleys Soil application. Minimum 6 months old					
MICRONUTRIENTS	 VL-Boron Boron 5% 1 pack Foliar, 500 ml per 100 L water 1 L	 VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Foliar, 1 L per 100 L water 1 L	 VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Foliar, 1 L per 100 L water 1 L	 VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Foliar, 1 L per 100 L water 1 L	 VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Foliar, 1 L per 100 L water 1 L	 VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Foliar, 1 L per 100 L water 1 L
FOLIAR	 VL-NPK NPK 8-8-6 (w/v) 1 pack Spray 1 L	 VL-NPK NPK 8-8-6 (w/v) 1 pack Spray. Repeat 15 days after flowering 1 L	 VL-NPK NPK 8-8-6 (w/v) 1 pack Spray. Repeat 15 days after fruiting 1 L	 VL-NPK NPK 8-8-6 (w/v) 1 pack Spray 1 L	 VL-NPK NPK 8-8-6 (w/v) 1 pack Spray 1 L	 VL-NPK NPK 8-8-6 (w/v) 1 pack Spray 1 L

V-Potash Plus VL-Potash Liquid Cala-Mag V VL-Boron VL-Micromix VL-NPK MgSO₄ Well-rotten farmyard manure grey items: general products, not VAN brands

Indicative programme. Built on best practice and the nutrient requirement of the crop, per acre. VAN prepares farm-specific plans against your soil analysis, yield target and the other variables that decide a rate: ask for one. Vital Urea is applied by drilling at sowing, side dressing or broadcasting, never by fertigation; the fertigation rows in the nitrogen band are general urea, a different material. Every VAN batch is tested before it ships; the batch number on the bag brings its certificate.

Rows as published on van.com.pk/crops/guava.html, 27 Sep 2026. Rates are per acre unless marked. Where a row says "Drill at sowing; broadcast if no drill", the drill is first by VAN's order of methods.