

Turmeric / per acre

May-Jun sowing

S-Mg-K programme

24 rows · 5 stages

	Land Preparation	Germination	Early Growth	Grand Growth	Maturity
PHOSPHORUS	Green Phosphate P₂O₅ 32% · N 6% 1 bag Drill at sowing; broadcast if no drill 50 kg		V. Ammonium Phosphate (V-Phosphate) NPK 10-44-0 1 bag Fertigation 10 kg	V. Ammonium Phosphate (V-Phosphate) NPK 10-44-0 1 bag Fertigation 10 kg	
NITROGEN	Urea N 46% ½ bag Broadcasting 50 kg	Vital Urea N 32% · S 13% 1 bag Side dressing 25 kg	Urea N 46% ½ bag Fertigation 50 kg	Urea N 46% 1 bag Fertigation 50 kg	
POTASH	Fusion Potash K₂O 50% · N 2% · S 1.5% · B 1% · Mg 1% ½ bag Drill at sowing; broadcast if no drill 25 kg	Fusion Potash K₂O 50% · N 2% · S 1.5% · B 1% · Mg 1% ½ bag Broadcasting 25 kg	Vital Potash NPK 11.0.44 1 bag Fertigation 20 kg	Vital Potash NPK 11.0.44 1 bag Fertigation 20 kg	
SECONDARY NUTRIENT	Green Sulfur Sulfur 70% 5 bag Drill at sowing; broadcast if no drill 1 kg	V-Mag Essential Mg 8.5% · B 1% 1 bag Fertigation 5 kg	V-Mag Essential Mg 8.5% · B 1% 1 bag Fertigation 5 kg	V-Mag Essential Mg 8.5% · B 1% 1 bag Fertigation 5 kg	
SOIL AMENDMENT	Humi Grow Soil Conditioner 1 bag Drill at sowing; broadcast if no drill 8 kg				
MICRONUTRIENTS	FeSO4 (2 kg) 1 bag Broadcasting	V-Zinc Zinc 10% w/v 4¼ pack Fertigation 3 L		VL-Boron Boron 5% 1 pack Fertigation 1 L	
FOLIAR	VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Spray 1 L	VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Spray 1 L	VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Spray 1 L	VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Spray 1 L	VL-Micromix Zn 5% · Fe 2% · Mn 2% · Cu 1% (w/v) 1 pack Spray 1 L

- Green Phosphate
- V. Ammonium Phosphate (V-Phosphate)
- Vital Urea
- Fusion Potash
- Vital Potash
- Green Sulfur
- V-Mag Essential
- Humi Grow
- V-Zinc
- VL-Boron
- VL-Micromix
- Urea
- FeSO4 (2 kg)
- grey items: general products, not VAN brands

Drill at sowing, first
The seed drill's fertilizer box places the granule about 5 cm beside and below the seed row, never touching the seed. The root meets it first and the whole bag is in the root zone.

Side-dress, second
A band beside the row, by hand or planter, at sowing or the first irrigation, covered with soil.

Broadcast, last
Spread on the surface before the last ploughing or before an irrigation, and mixed in. Most of the bag is away from the root.

Indicative programme. Built on best practice and the nutrient requirement of the crop, per acre. VAN prepares farm-specific plans against your soil analysis, yield target and the other variables that decide a rate: ask for one. Vital Urea is applied by drilling at sowing, side dressing or broadcasting, never by fertigation; the fertigation rows in the nitrogen band are general urea, a different material. Every VAN batch is tested before it ships; the batch number on the bag brings its certificate.

Rows as published on van.com.pk/crops/turmeric.html, 27 Sep 2026. Rates are per acre unless marked. Where a row says "Drill at sowing; broadcast if no drill", the drill is first by VAN's order of methods.