

(a) Golub Train vs. Independent Test Shift
Training Set (n = 38)
Independent Test (n = 34)
PC2 (11.0%)
PC1 (12.5%)
(b) Cross-Platform Harmonization (5,426 Genes)
Golub (Hu6800, n = 72)
Agarwal (GSE51082) (GPL96, n = 66)
PC2 (9.7%)
PC1 (11.4%)
(c) Tissue Confounding Diagnostic
BCLL (n=41)*
MDS (n=10)*
CML (n=22)*
ALL (n=29)
AML (n=37)
ALL vs. AML Modeling:
BM: 86.2% vs. 81.1%
Fisher's Exact: OR=1.46, p=0.743 (n.s.)
-> No tissue confounding in ALL/AML
*Excluded: BCLL (92.7% PB), MDS (100% BM)
Bone Marrow (BM)
Peripheral Blood (PB)
Lymph Node (LN)
Tissue Proportion (%)
(d) 6-Class Global Landscape (n=139)
AML (n=37, Model)
PRE B-ALL (n=17, Model)
T-ALL (n=12, Model)
BCLL (n=41, Excl.)
CML (n=22, Excl.)
MDS (n=10, Excl.)
MDS -> AML continuity
PC2 (13.3%)
PC1 (26.2%)