

CLENPIQ™

(sodium picosulfate, magnesium oxide,
and anhydrous citric acid) Oral Solution
10 mg/3.5 g/12 g per 160 mL bottle


TAKING CLENPIQ

STEPS TO COMPLETE THE PREP

Day-Before-Dose Regimen

Afternoon/Early Evening Before


BETWEEN 2 PM - 6 PM



Drink 40 oz (5 cups) of clear liquids.*
Finish liquids over the next 5 hours.

Evening Before

Between 8 PM - 11 PM



Drink at least 24 oz (3 cups) of clear liquids over the next 5 hours.
Finish liquids 2 hours before your colonoscopy
or as advised by your doctor.



Hydration is important and it's part of the prep. Make sure to hydrate
before you take the prep, while you're taking the prep, and after the prep.

Prep Assistant Use the prep assistant as a guide for completing your prep.

The date and time of your colonoscopy is / / : :

You can keep hydrating with clear liquids until 12:00 PM the day of your procedure.

**ON THE DAY
BEFORE THE
COLONOSCOPY,**
stop eating all solid
food and dairy, and
start hydrating by
drinking
clear liquids.



Afternoon/Early Evening before your colonoscopy

- ☒ Drink one bottle of CLENPIQ 1st bottle
Drink CLENPIQ right from the bottle

Then, keep hydrating
Drink 5 cups (8 oz each) of clear liquid.
Check off the cups below as you go!




Start: 2:00 PM
Finish: :

Evening before your colonoscopy

- ☒ Drink the other bottle of CLENPIQ 2nd bottle
Drink CLENPIQ right from the bottle

Then, keep hydrating
Drink at least 3 cups (8 oz each) of clear liquid.
Check off the cups below as you go!



Start: 8:00 PM
Finish: :

*After your first dose, if severe bloating, swelling, or stomach pain occurs, delay the second dose until the symptoms resolve.
Please see reverse page for additional information about CLENPIQ, the importance of a clear liquid diet, and hydration.

INDICATION

CLENPIQ is a prescription medicine used by adults to clean the colon before a colonoscopy. CLENPIQ cleans your colon by causing you to have diarrhea. Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

IMPORTANT SAFETY INFORMATION

- Do not take CLENPIQ if your healthcare provider has told you that you have serious kidney problems, a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestines (bowel perforation), a very dilated intestine (toxic megacolon), problems with the emptying of food and fluid from your stomach (gastric retention), or an allergy to any of the ingredients in CLENPIQ.
- CLENPIQ and other bowel preparations can cause serious side effects, including serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause abnormal heartbeats that may result in death, seizures (this can happen even if you have never had a seizure), or kidney problems. Your chance of having fluid loss and changes in blood salts with CLENPIQ is higher if you have heart problems, have kidney problems, or take water pills or non-steroidal anti-inflammatory drugs (NSAIDs).

See complete Important Safety Information on the back.