

高等数学模拟试卷（五）参考答案

1. 解：将 $x=2$ 代入分子中，分子等于 0，且式中含有 $|x-2|$

所以 $x=2$ 是 $f(x)$ 的跳跃间断点

2. 解： $f'_-(0) = \lim_{x \rightarrow 0^-} \frac{f(x) - f(0)}{x} = \lim_{x \rightarrow 0^-} \frac{x^2 + ax}{x} = a$

$$f'_+(0) = \lim_{x \rightarrow 0^+} \frac{f(x) - f(0)}{x} = \lim_{x \rightarrow 0^+} \frac{e^x - 1}{x} = 1$$

$\therefore f(x)$ 在 $x=0$ 处可导， \therefore 应有 $f'_-(0) = f'_+(0)$ ，即 $a=1$

3. 解： $\varphi'(x) = -e^{\frac{1}{x}} \sin \frac{1}{x} \cdot \frac{-1}{x^2} = \frac{1}{x^2} e^{\frac{1}{x}} \sin \frac{1}{x}$

4. 解： $f'(x) = e^{-x} - xe^{-x} = (1-x)e^{-x}$

$$f''(x) = -e^{-x} - (1-x)e^{-x} = (x-2)e^{-x}$$

$$\text{令 } \begin{cases} f'(x) < 0 \\ f''(x) < 0 \end{cases}, \text{ 即 } \begin{cases} (1-x)e^{-x} < 0 \\ (x-2)e^{-x} < 0 \end{cases} \Rightarrow \begin{cases} 1-x < 0 \\ x-2 < 0 \end{cases} \Leftrightarrow \begin{cases} x > 1 \\ x < 2 \end{cases}$$

5. 解： $\int_1^{\sqrt{2}} dx \int_0^{\sqrt{2-x^2}} f(x, y) dy = \int_0^{\frac{\pi}{4}} d\theta \int_{\frac{1}{\cos \theta}}^{\sqrt{2}} f(r \cos \theta, r \sin \theta) r dr$

6. 解：对于 B：取 $V_n = \frac{n}{n^2 \sqrt{n}} = \frac{1}{n \sqrt{n}}$ ， $\therefore \sum_{n=1}^{\infty} \frac{1}{n \sqrt{n}}$ 收敛，所以 $\sum_{n=1}^{\infty} \frac{n+1}{n^2 \sqrt{n+1}}$ 收敛

7. 解： $|A| \neq 0 \Rightarrow A$ 可逆， $AB = AC$ 两边左乘 A^{-1}

$$A^{-1}AB = A^{-1}AC \Rightarrow B = C$$

8. 解：个数大于维数必相关，选 C

9. 解： $\lim_{x \rightarrow +\infty} \left(\frac{1+x}{2+x} \right)^{2x} = \lim_{x \rightarrow +\infty} e^{\left(\frac{1+x}{2+x} - 1 \right) \cdot 2x} = e^{-2}$


\therefore 水平渐近线方程为 $y = e^{-2}$

10. 解： $f(0) = a$

$$\lim_{x \rightarrow 0} f(x) = \lim_{x \rightarrow 0} \frac{\sin x}{\sqrt{1+x}-1} = \lim_{x \rightarrow 0} \frac{x}{\frac{1}{2}x} = 2$$

$\therefore f(x)$ 在 $x=0$ 处连续, \therefore 应有 $\lim_{x \rightarrow 0} f(x) = f(0)$, 故 $a=2$

$$\begin{aligned} 11. \text{ 解: } \int_{-2}^2 (x^2 \sin x + 1) \sqrt{4-x^2} dx &= \int_{-2}^2 x^2 \sin x \sqrt{4-x^2} dx + \int_{-2}^2 \sqrt{4-x^2} dx \\ &= 0 + \int_{-2}^2 \sqrt{4-x^2} dx = 2\pi \end{aligned}$$



$$12. \text{ 解: 方程两边对 } x \text{ 求偏导数, 得: } 2z \frac{\partial z}{\partial x} - 2yz - 2xy \frac{\partial z}{\partial x} + 2x = 0$$

$$\therefore \frac{\partial z}{\partial x} = \frac{yz-x}{z-xy}$$

$$13. \text{ 解: 令 } \lim_{n \rightarrow \infty} \left| \frac{(x+1)^{n+1}}{(n+1) \cdot 2^{n+1}} \cdot \frac{n \cdot 2^n}{(x+1)^n} \right| = \frac{1}{2} |x+1| < 1$$

$$-2 < x+1 < 2, \quad -3 < x < 1$$

$$\text{当 } x=-3 \text{ 时, 级数为 } \sum_{n=1}^{\infty} \frac{(-2)^n}{n \cdot 2^n} = \sum_{n=1}^{\infty} \frac{(-1)^n}{n} \text{ 收敛}$$

$$\text{当 } x=1 \text{ 时, 级数为 } \sum_{n=1}^{\infty} \frac{2^n}{n \cdot 2^n} = \sum_{n=1}^{\infty} \frac{1}{n} \text{ 发散}$$

收敛域 $[-3, 1)$

$$14. \text{ 解: } A^2 = \begin{pmatrix} 2 & -2 \\ 0 & 2 \end{pmatrix} \begin{pmatrix} 2 & -2 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 4 & -8 \\ 0 & 4 \end{pmatrix}$$

$$\begin{aligned}\therefore B &= A^2 - 2A + E_2 = \begin{pmatrix} 4 & -8 \\ 0 & 4 \end{pmatrix} - 2\begin{pmatrix} 2 & -2 \\ 0 & 2 \end{pmatrix} + E_2 \\ &= \begin{pmatrix} 4 & -8 \\ 0 & 4 \end{pmatrix} - \begin{pmatrix} 4 & -4 \\ 0 & 4 \end{pmatrix} + E_2 = \begin{pmatrix} 1 & -4 \\ 0 & 1 \end{pmatrix}\end{aligned}$$

$$\begin{aligned}15. \text{ 解: } \lim_{x \rightarrow 0} \frac{x - x^3 - \sin x}{x^2 \arctan x} &= \lim_{x \rightarrow 0} \frac{x - x^3 - \sin x}{x^3} = \lim_{x \rightarrow 0} \frac{1 - 3x^2 - \cos x}{3x^2} \\ &= \lim_{x \rightarrow 0} \frac{-6x + \sin x}{6x} = \lim_{x \rightarrow 0} \frac{-6 + \cos x}{6} = -\frac{5}{6}\end{aligned}$$

$$16. \text{ 解: 方程两边对 } x \text{ 求导, 得: } e^y \cdot y' + 6y + 6xy' + 2x = 0 \quad ①$$

①式两边再对 x 求导, 得:

$$e^y \cdot (y')^2 + e^y \cdot y'' + 6y' + 6y' + 6xy'' + 2 = 0 \quad ②$$

当 $x = 0$ 时, 由原方程可知 $y = 0$

将 $x = 0$, $y = 0$ 代入①式, 得: $y'(0) = 0$

将 $x = 0$, $y = 0$, $y'(0) = 0$ 代入②式, 得: $y''(0) + 2 = 0$, $y''(0) = -2$

$$\begin{aligned}17. \text{ 解: } \int \frac{x \sin x}{\cos^3 x} dx &= \frac{1}{2} \int x d \frac{1}{\cos^2 x} = \frac{x}{2 \cos^2 x} - \frac{1}{2} \int \frac{1}{\cos^2 x} dx \\ &= \frac{x}{2 \cos^2 x} - \frac{1}{2} \tan x + c\end{aligned}$$

$$18. \text{ 解: 令 } \sqrt{2x+1} = t, \text{ 则 } x = \frac{t^2-1}{2}, \text{ 当 } x = -\frac{1}{2} \text{ 时 } t = 0, \text{ 当 } x = \frac{3}{2} \text{ 时 } t = 2$$

$$\begin{aligned}\int_{-\frac{1}{2}}^{\frac{3}{2}} \frac{\sqrt{2x+1}}{2x+5} dx &= \int_0^2 \frac{t}{t^2+4} d \frac{t^2-1}{2} = \int_0^2 \frac{t^2}{t^2+4} dt \\ &= \int_0^2 dt - 4 \int_0^2 \frac{1}{t^2+2^2} dt = t \Big|_0^2 - 2 \arctan \frac{t}{2} \Big|_0^2 \\ &= 2 - \frac{\pi}{2}\end{aligned}$$

$$19. \text{ 解: } \frac{\partial z}{\partial y} = f_1' \cdot 1 + f_2' \cdot e^{x-y} \cdot (-1) = f_1' - e^{x-y} f_2'$$

$$\frac{\partial^2 z}{\partial y \partial x} = f_{11}'' \cdot 1 + f_{12}'' \cdot e^{x-y} - e^{x-y} f_2' - e^{x-y} (f_{21}' \cdot 1 + f_{22}' \cdot e^{x-y})$$


$$= f_{11}'' - e^{x-y} f_2' + e^{2(x-y)} f_{22}''$$

$$20. \text{ 解: } \iint_D (x+y) dx dy = \int_0^1 dx \int_0^x (x+y) dy + \int_1^2 dx \int_0^{\frac{1}{x}} (x+y) dy$$

$$= \int_0^1 dx \left(xy + \frac{1}{2} y^2 \right) \Big|_0^x + \int_1^2 dx \left(xy + \frac{1}{2} y^2 \right) \Big|_0^{\frac{1}{x}}$$

$$= \int_0^1 \frac{3}{2} x^2 dx + \int_1^2 \left(1 + \frac{1}{2x^2} \right) dx$$

$$= \frac{1}{2} x^3 \Big|_0^1 + \left(x - \frac{1}{2x} \right) \Big|_1^2 = \frac{7}{4}$$



$$21. \text{ 解: } \int \frac{f(x)}{x} dx = e^{2x} + c \text{ 两边对 } x \text{ 求导, 得 } \frac{f(x)}{x} = 2e^{2x}, \therefore f(x) = 2xe^{2x}$$

$$\text{所求微分方程为 } y'' + y' - 2y = 2xe^{2x}$$

$$\text{对应齐次方程的特征方程 } r^2 + r - 2 = 0, \text{ 特征根 } r_1 = 1, r_2 = -2$$

$$\text{其通解 } \bar{y} = c_1 e^x + c_2 e^{-2x}$$

$$\text{设原方程一个特解 } y^* = (ax + b)e^{2x}$$

$$y^{*'} = ae^{2x} + 2(ax + b)e^{2x}, \quad y^{*''} = 4ae^{2x} + 4(ax + b)e^{2x}$$

将 y^* 代入原方程, 得:

$$4ae^{2x} + 4(ax+b)e^{2x} + ae^{2x} + 2(ax+b)e^{2x} - 2(ax+b)e^{2x} = 2xe^{2x}$$

$$\begin{cases} 4a=2 \\ 5a+4b=0 \end{cases} \Rightarrow \begin{cases} a=\frac{1}{2} \\ b=-\frac{5}{8} \end{cases}, y^* = \left(\frac{1}{2}x - \frac{5}{8}\right)e^{2x}$$

$$\text{原方程通解 } y = \bar{y} + y^* = c_1 e^x + c_2 e^{-2x} + \left(\frac{1}{2}x - \frac{5}{8}\right)e^{2x}$$

$$22. \text{ 解: } |A| = \begin{vmatrix} 3 & -2 & 1 & 4 \\ -7 & 5 & -3 & -6 \\ 2 & 1 & -1 & 3 \\ 4 & -3 & 2 & 8 \end{vmatrix} \xrightarrow[r_4-2r_1]{r_2-3r_1, r_3+r_2} \begin{vmatrix} 3 & -2 & 1 & 4 \\ 2 & -1 & 0 & 6 \\ 5 & -1 & 0 & 7 \\ -2 & 1 & 0 & 0 \end{vmatrix} \xrightarrow{\text{按 } c_3 \text{ 展开}} \begin{vmatrix} 2 & -1 & 6 \\ 5 & -1 & 7 \\ -2 & 1 & 0 \end{vmatrix}$$

$$\xrightarrow{c_1+2c_2} \begin{vmatrix} 0 & -1 & 6 \\ 3 & -1 & 7 \\ 0 & 1 & 0 \end{vmatrix} \xrightarrow{\text{按 } r_3 \text{ 展开}} - \begin{vmatrix} 0 & 6 \\ 3 & 7 \end{vmatrix} = 18$$

$$\therefore |A^{-1}| = \frac{1}{|A|} = \frac{1}{18}$$

$$23. \text{ 证明: 令 } F(x) = x^3 + (1-x)^3 - \frac{1}{4}$$

$$F'(x) = 3x^2 - 3(1-x)^2, \text{ 令 } F'(x) = 0, \text{ 得 } x = \frac{1}{2}$$

$$\text{当 } 0 < x < \frac{1}{2} \text{ 时, } F'(x) < 0, \text{ 当 } \frac{1}{2} < x < 1 \text{ 时, } F'(x) > 0$$


$$\text{所以 } F(x) \text{ 在 } x = \frac{1}{2} \text{ 处取得最小值, 最小值 } F\left(\frac{1}{2}\right) = 0$$

$$\text{故当 } 0 \leq x \leq 1 \text{ 时, } F(x) \geq 0, \text{ 即 } x^3 + (1-x)^3 \geq \frac{1}{4}$$

$$24. \text{ 解: (1) } S = \frac{1}{2} \times 1 \times 1 + \int_0^1 (1-x^2) dx = \frac{1}{2} - \left(x - \frac{1}{3}x^3\right) \Big|_0^1 = \frac{7}{6}$$

$$(2) V_x = \pi \int_{-1}^1 1^2 dx - \pi \int_{-1}^0 (-x)^2 dx - \pi \int_0^1 (x^2)^2 dx$$

$$= \pi x \Big|_{-1}^1 - \frac{\pi}{3} x^3 \Big|_{-1}^0 - \frac{\pi}{5} x^5 \Big|_0^1 = \frac{22}{15} \pi$$



$$\begin{aligned}
 25. \text{ 解: } \bar{A} &= \begin{pmatrix} 1 & 1 & -1 & -1 & 1 & 0 \\ 2 & 2 & 1 & 0 & 1 & 1 \\ 3 & 3 & 0 & -1 & 2 & 1 \\ 1 & 1 & 2 & 1 & 0 & 1 \end{pmatrix} \xrightarrow{\substack{r_2-2r_1 \\ r_3-3r_1 \\ r_4-r_1}} \begin{pmatrix} 1 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 3 & 2 & -1 & 1 \\ 0 & 0 & 3 & 2 & -1 & 1 \\ 0 & 0 & 3 & 2 & -1 & 1 \end{pmatrix} \\
 &\xrightarrow{\substack{r_3-r_2 \\ r_4-r_2}} \begin{pmatrix} 1 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 3 & 2 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \xrightarrow{\frac{1}{3}r_2} \begin{pmatrix} 1 & 1 & -1 & -1 & 1 & 0 \\ 0 & 0 & 1 & \frac{2}{3} & -\frac{1}{3} & \frac{1}{3} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 &\xrightarrow{r_1+r_2} \begin{pmatrix} 1 & 1 & 0 & -\frac{1}{3} & \frac{2}{3} & \frac{1}{3} \\ 0 & 0 & 1 & \frac{2}{3} & -\frac{1}{3} & \frac{1}{3} \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}
 \end{aligned}$$

x_1, x_3 为约束变量, x_2, x_4, x_5 为自由变量

$$\begin{cases} x_1 = \frac{1}{3} - x_2 + \frac{1}{3}x_4 - \frac{2}{3}x_5 \\ x_2 = \frac{1}{3} - \frac{2}{3}x_4 + \frac{1}{3}x_5 \end{cases} \text{ 令 } \begin{cases} x_2 = 0 \\ x_4 = 0 \\ x_5 = 0 \end{cases}, \text{ 得原方程组的一个特解 } \eta^* = \begin{pmatrix} \frac{1}{3} \\ \frac{1}{3} \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$\text{导出组一般解} \begin{cases} x_1 = -x_2 + \frac{1}{3}x_4 - \frac{2}{3}x_5 \\ x_2 = -\frac{2}{3}x_4 + \frac{1}{3}x_5 \end{cases}$$

$$\text{令} \begin{cases} x_2 = 1 \\ x_4 = 0 \\ x_5 = 0 \end{cases}, \text{得} \xi_1 = \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \text{令} \begin{cases} x_2 = 0 \\ x_4 = 1 \\ x_5 = 0 \end{cases}, \text{得} \xi_2 = \begin{pmatrix} \frac{1}{3} \\ 0 \\ -\frac{2}{3} \\ 1 \\ 0 \end{pmatrix}$$

$$\text{令} \begin{cases} x_2 = 0 \\ x_4 = 0 \\ x_5 = 1 \end{cases}, \text{得} \xi_3 = \begin{pmatrix} -\frac{2}{3} \\ 0 \\ \frac{1}{3} \\ 0 \\ 1 \end{pmatrix}$$

$$\text{原方程组通解} \begin{pmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{pmatrix} = \begin{pmatrix} \frac{1}{3} \\ 0 \\ \frac{1}{3} \\ 0 \\ 0 \end{pmatrix} + k_1 \begin{pmatrix} -1 \\ 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + k_2 \begin{pmatrix} \frac{1}{3} \\ 0 \\ -\frac{2}{3} \\ 1 \\ 0 \end{pmatrix} + k_3 \begin{pmatrix} -\frac{2}{3} \\ 0 \\ \frac{1}{3} \\ 0 \\ 1 \end{pmatrix} \quad (\text{其中 } k_1, k_2, k_3 \text{ 为任意}$$

常数)