

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/344541693>

A REVIEW ON RECENT METHODS TO CONTROL EARLY BLIGHT OF TOMATO (*Solanum lycopersicum* L.)

Article in PLANT CELL BIOTECHNOLOGY AND MOLECULAR BIOLOGY · September 2020

CITATIONS

9

READS

4,878

2 authors:



Denita Shamurailatpam

Lovely Professional University

2 PUBLICATIONS 12 CITATIONS

[SEE PROFILE](#)



Adesh Kumar

Lovely Professional University

76 PUBLICATIONS 462 CITATIONS

[SEE PROFILE](#)

A REVIEW ON RECENT METHODS TO CONTROL EARLY BLIGHT OF TOMATO (*Solanum lycopersicum* L.)

DENITA SHAMURAILATPAM AND ADESH KUMAR*

Department of Plant Pathology, School of Agriculture, Lovely Professional University,
Phagwara - 144411, Punjab, India [DS, AK].

[*For Correspondence: E-mail: adesh.19078@lpu.co.in, phytopath06@gmail.com]

Article Information

Editor(s):

(1) Dr. Milan Stankovic, University of Kragujevc, Serbia.

Reviewers:

(1) Muhammad Shehzad Khan, The University of Agriculture, Pakistan.

(2) K. Prasannath, Eastern University, Sri Lanka.

Received: 18 June 2020

Accepted: 24 August 2020

Published: 12 September 2020

Review Article

ABSTRACT

The present review paper is focused on bringing forth the different methods used in controlling early blight disease of tomato (*Solanum lycopersicum* L.). It is a well-known fact that tomato is a main fruit consumed as a vegetable globally which provides important minerals, vitamins, fibres and antioxidants. However, tomato is susceptible to an array of diseases, of which early blight is the most destructive instigated by *Alternaria solani*, affects fruits and foliage resulting in a loss of 80% yields. Therefore, it becomes imperative to discuss the various strategies applied in controlling early blight of tomato. A number of researchers have used diverse methods to regulate early blight disease of tomato ranging from use of fungicides, plant extracts to several biological organisms. Nevertheless, it is equally important to develop and apply the most efficient management systems and good agricultural practices to rescue the plant of interest.

Keywords: *Alternaria solani*; biocontrol; fungicides; plant extracts; *Solanum lycopersicum*.

INTRODUCTION

Tomato is the utmost important and popular vegetable crop grown round the world [1,2]. Tomato belongs to the genus *Lycopersicon* and Family Solanaceae [3]. China is the leading country in production of tomato (31%), followed by India and the United States with the second and third highest producer in the world [4]. The consumption of tomato stands second after potato being rich in vitamins (K, C and A), minerals (Fe, Ca and P), amino acids, sugars, dietary fibres and antioxidant and contains 95.3% of water [5,6]. The

production of tomato suffered huge loss in production due to the attack by various bacterial, viral and fungal diseases [7]. The main diseases are fusarium wilt, early blight, damping off, late blight, tomato mosaic virus, verticillium wilt, and bacterial wilt [8,9,10]. Out of all, the most damaging disease caused by pathogenic fungus *A. solani* is early blight, which results in major loss both in quantity and quality of fruit yield [11,12]. The disease symptoms appear on stems, leaves and fruits of tomato [13]. Small and dark lesions appear on the leaves of tomato plant [13]. Necrotic and concentric rings on leaves give a target like

appearance on the surface of leaf. Lesions appeared on leaf surface are bound by yellow rings [4]. Older leaves are firstly infected by *A. solani* followed by stems, petioles, twigs and fruits of tomato plant. Control of this disease is very difficult due to wide host range and active and prolonged life cycle of *A. solani*.

DIFFERENT STRATEGIES FOR MANAGING EARLY BLIGHT

Different methods have been used for the control and better management of this disease. Methods used include cultural practices, through resistant varieties, biological control, use of chemicals and disease free planting material [14]. The use of fungicide is considered as effective method for managing early blight. The usage of resistant and tolerant varieties is also an important method to control this disease. Resistant varieties increase effect of fungicides, extend fungicide spray intervals and maintain disease. Different fungicides for combating early blight disease are available in market. Azoxystrobin, ziram, pyraclostrobin, mancozeb, copper products and potassium bicarbonate are name of some fungicides used for the same.

Use of Fungicides

The early blight of tomato is considered as the most devastating disease resulting in pre and post-harvest stages resulting in a drop in yields of 35 to 78% [4,15]. The use of fungicides is considered as the most effective approach for controlling early blight, as stated by several workers [16,17,18,14,19,20,15]. The fungicide mancozeb has been widely used since the last two decades to control early blight, reducing the severity of the disease and enhances tomato yield [4]. Farooq et al. [15] stated that hexaconazole and pyraclostrobin are the best fungicides to reduce the occurrence of the disease by inhibiting maximum growth of mycelium. While, according to Bais et al. [4], sanchar followed by mancozeb and carbendazim were superior for controlling the disease. Rani et al. [18] reported that tebuconazole and difenoconazole were the most effective fungicides in controlling early blight of tomato.

However, due to concerns about cost, toxicity risks and dangers of its residues, the use of

fungicides cannot be seen as a long-term solution. In addition, the production of pathogenic fungi resistance to synthetic pesticides is a major problem, which can have a noteworthy impact on the efficacy of chemical fungicides [18,12,9]. Inappropriate use of chemicals lead to serious human health hazards and possible harmful effects on non-target organisms and the environment [21]. Therefore, proper concentration of fungicides at proper intervals should be mandatory [16]. The use of fungicides can be altered to be in a safer range by combining with other relatively safer alternative methods [14]. An overview of the different works published in different journals in this area is presented in Table 1.

Use of Biocontrol Agents

The worldwide trend towards safe environment methods for controlling plant diseases in sustainable agriculture practice needs reduced usage of synthetic chemicals [4]. Due to this, there is an increasing interest to acquire alternative antimicrobial agents (biocontrol agents) for controlling plant diseases [22,15]. Biocontrol agent is the effective and ecological alternative approach for managing early blight. Several researchers have shown the use of biological agents as a best alternative method for the control of early blight (Table 2).

Biocontrol bacterial strains viz. *Pseudomonas fluorescens* [4], *P. aeruginosa* [39], and *P. putida* [38] have resulted in low disease severity index. A very well-known and effective biocontrol fungal strain, *Trichoderma viride* has also been used for control of early blight. *T. viride* with *Pseudomonas* strains is another effective biocontrol strategy used for disease management. Owing to strong mycoparasitism of fungal pathogens and rapid growth potential, *Trichoderma* species are considered as effective control agents [37,39]. *Trichoderma* antagonism is owed to the development of non-volatile and volatile metabolites, space, food, nutrient rivalry, the ability to secrete enzymes that degrade cell wall and antibiosis [14]. As a result, there is a growing interest in acquiring alternative antimicrobials (bio-control agents) for use in plant disease control systems. Bio control agents are found to be effective and eco-friendly. Furthermore, they are easily biodegradable, non-

phytotoxic, systemic and safe to environment [7]. Various biocontrol agents including *Pseudomonas* spp., *Bacillus* spp. and *Serratia* spp. may induce plant protection responses which are directly

related to the pathogenesis related proteins (PR) such as phenylalanine ammonia-lyase, superoxide dismutase, peroxidase, and β -1,3 glucanase defence enzymes [40].

Table 1. Different fungicides used to control early blight

S. no.	Fungicides	Method of treatment	Results	References
1	Carbendazim Mancozeb Copper oxychloride Iprodione Chlorothalonil	Poisoned food technique	-Mancozeb was most effective fungicide inhibiting sporulation and growth of mycelium of <i>A. solani</i> .	Pandey; [23]
2	Fungicides	Poisoned food technique	-Mancozeb (0.2%) and hexaconazole (0.1%) showed mycelial growth inhibition by 87.21% and 88.88% respectively. -In combination hexaconazole and zineb showed maximum inhibition of mycelial growth.	Roopa <i>et al.</i> [24]
3	Onestar Ergon Amistar	RCBD	-Onestar 23% SC and amistar 23% SC showed maximum reduction in disease (69-71%).	Saxena <i>et al.</i> [25]
4	Duter Mancozeb Carbendazim Thiophanate methyl Zineb Iprodione Captafol Copper oxychloride	RCBD	-Mancozeb (0.2%) recorded minimum percent disease incident (22.18) and 32.2% maximum disease control, which was superior to other fungicides. -Carbendazim (0.1%) showed 28.22% and copper oxychloride (0.25%) showed 24.85% disease control.	Walke <i>et al.</i> [26]
5	Mancozeb	RCBD	-As compared to control, disease intensity was reduced by mancozeb with foliar spray (15.43%, 17.90% and 20.47%).	Zghair <i>et al.</i> [27]
6	Ridomil gold Agrolaxyl Mancozeb	RCBD	-Mancozeb was most effective in controlling the disease by 47.75%.	Desta and Yesuf; [28]
7	Carbendazim Copper sulphate Copper oxychloride Captan Mancozeb Thiram Zineb	Poison food technique	-At concentration 1500 ppm, mancozeb reduced mycelial growth by 86.4%, which was followed by carbendazim (33.7%).	Sadana and Didwania; [29]
8	Mancozeb Chlorothalonil	Randomized complete block design (RCBD)	- <i>In vivo</i> results validated decline in disease control with the reduced-sensitivity isolates related to the sensitive ones.	Abu-El Samen <i>et al.</i> [30]
9	Propineb Thiophanate methyl Chlorothalonil Mancozeb Copper oxychloride	Poison food technique	-Mancozeb (89.83%) showed maximum mycelial growth inhibition followed by propineb (87.40%), thiophanate-methyl (87.10%) and copper oxychloride (79.21%). -Least inhibition shown by chlorothalonil (70.40%).	Ghazanfar <i>et al.</i> [16]
10	Mancozeb Chlorothalonil Propineb Hexaconazole Azoxystrobin Difenoconazole Tebuconazole Propiconazole Carbendazim	Poisoned food technique	-Mancozeb, hexaconazole and combi fungicide carbendazim + mancozeb recorded maximum inhibition of 88.42%, 90.58% 88.07% mycelial growth respectively.	Mahantesh <i>et al.</i> [17]
11	Tebuconazole Hexaconazole Difenoconazole Carbendazim Mancozeb	Poisoned food technique	-The most effective fungicide was tebuconazole followed by difenoconazole.	Rani <i>et al.</i> (2017)

S. no.	Fungicides	Method of treatment	Results	References
12	Trigger Solex Dew Amistar Top Corel Reflex	Poisoned food technique	-Reflex and corel showed better inhibitory effect on <i>A. solani</i> . -Solex showed no inhibitory effect.	Sarfranz et al. [18]
13	Carbendazim Mancozeb Difenoconazole Propiconazole Mancozeb Propineb Copper-oxy chloride	Poisoned food technique	-Carbendazim and mancozeb showed lowest percent disease intensity, which was followed by difenoconazole and propiconazole.	Sharma et al. [19]
14	Mancozeb RokoCarbendazim Companion Blitox 50 Sanchar	Poison food technique	-The significant maximum growth inhibition was recorded in sanchar followed by mancozeb and carbendazim.	Bais et al. [4]
15	Badge X2 Prestop Serenade Sil-Matrix Opti Regalia Oxidate Sustane	RCBD	-Badge X2 and prestop reduced disease significantly both in field and green house trails. -Sil-Matrix, regalia, oxidate and sustane were ineffective in controlling disease.	Egel et al. [20]
16	Pyraclostrobin Hexaconazole Carbendazim Mancozeb Ridomil	Poisoned food technique	-At 500 ppm, the maximum inhibition of mycelial growth was shown by pyraclostrobin (97.7%), followed by hexaconazole (91.1%), carbendazim (90%) and ridomil showed inhibition i.e, 42.2%.	Farooq et al. [15]
17	Indofil Bavistin Sulcox Tall Mancozeb	Poisoned food technique	-The most effective fungicide was Tall 25EC which showed maximum mycelial inhibition.	Roy et al. [10]

Table 2. Bio control agents used against early blight

S. no.	Biocontrol agents	Method of treatment	Result	References
1	<i>Bacillus</i> species	Dual culture technique	-Out of 45 <i>Bacillus</i> isolated, 27 showed antagonistic effects against pathogen <i>in vitro</i> and four among them showed highest effect.	Abdalla et al. [31]
2	<i>T. harzianum</i> <i>T. virens</i>	Dual culture technique	- <i>T. virens</i> displayed better inhibitory effect on radial growth and sporulation of mycelium.	Pandey, [23]
3	<i>T. harzianum</i> , <i>T. viride</i> , <i>T. virens</i>	Dual culture technique	- <i>T. harzianum</i> inhibits mycelial growth by 77.50% followed by <i>T. viride</i> 75.14%.	Roopa et al. [24]
4	<i>T. harzianum</i> <i>T. viride</i>	Dual culture technique	-Three sprays of <i>T. viride</i> (0.5%) were most effective against mycelial growth (31.47%), when compared to control (80.90%).	Yadav; [32]
5	<i>T. harzianum</i> <i>P. fluorescens</i>	Dual culture technique	- <i>T. harzianum</i> and <i>P. fluorescens</i> were effective in reducing the disease intensity.	Zghair et al. [27]
6	<i>Pseudomonas fluorescens</i> <i>Trichoderma viride</i> <i>Bacillus subtilis</i> <i>T. harzianum</i>	Dual culture technique	- <i>Bacillus subtilis</i> was most effective against mycelial growth (52.77%) of the fungus <i>A. solani</i> . - <i>P. fluorescens</i> isolates showed 47.22% and 45.55% mycelial growth inhibition. - <i>T. harzianum</i> and isolates of <i>T. viride</i> showed less than 40% growth inhibition.	Koley et al. [33]
7	<i>Pseudomonas aeruginosa</i>	RCBD	- <i>In vitro</i> studies suggested that, this bacterial strain produces a biosurfactant that inhibits the growth of pathogen at concentration of 3.0 g/l by 73%.	Lahkar et al. [34]

S. no.	Biocontrol agents	Method of treatment	Result	References
			-Concentration of crude biosurfactant in field conditions i.e, 1.50 g/l was adequate for full inhibition of <i>A. solani</i> .	
8	Rhizobacteria	Dual culture technique	-Almost all isolates shown antifungal activity against <i>A. solani</i> . -Best result observed with treatment 22 which inhibited 53.28% colony followed by treatment 25 (36.25%). -Minimum% inhibition was found in treatment 24 (19.94%).	Maurya et al. [35]
9	<i>Trichoderma spp.</i>	Dual culture technique	- <i>Trichoderma</i> isolates reduced mycelial growth of <i>Alternaria solani</i> .	Selim; [36]
10	<i>P. fluorescens</i> <i>H. armigera</i>	RCBD	-Both biocontrol treatments were found better than untreated control in reducing the occurrence of early blight in tomatoes.	Kaur et al. (2016)
11	<i>T. harzianum</i> <i>T. hamatum</i> <i>T. viride</i> <i>Penicillium glabrum</i> <i>Penicillium citrinum</i>	Dual culture technique	- <i>A. solani</i> completely overgrown (100%, class I) by the antagonists <i>T. harzianum</i> and <i>T. hamatum</i> while <i>T. viride</i> colonized half of the growth of fungus (50% overgrowth, Class III). - <i>Penicillium citrinum</i> and <i>P. glabrum</i> could not overgrow the fungus.	Devi et al. [7]
12	<i>T. viride</i> <i>P. fluorescens</i>	RCBD	-The early blight incidence was reduced by 93.4% and 88.8% over control when infected lower leaves were removed, staking of tomato plants, seed priming with <i>T. viride</i> followed by foliar spraying with <i>T. viride</i> and <i>Pseudomonas fluorescens</i> . -The <i>T. viride</i> and <i>P. fluorescens</i> foliar spray with priming of seeds also reduced the disease by 78.0% and 73.0% respectively.	Dhal et al. [37]
13	<i>T. harzianum</i> <i>T. viride</i> <i>P. fluorescens</i>	RCBD	-Minimum disease intensity recorded in treatment <i>Trichoderma viride</i> (31.44%) as compared to control (49.16%).	Jaiswal et al. (2017)
14	<i>P. aeruginosa</i> <i>P. putida</i> <i>P. fluorescens</i> <i>P. cepacia</i>	Dual culture technique	- <i>P. aeruginosa</i> showed maximum effect on early blight. - <i>P. putida</i> , <i>P. cepacia</i> and <i>P. fluorescens</i> showed no significantly different effects.	Joseph et al. [38]
15	<i>T. harzianum</i> <i>T. viride</i> <i>P. fluorescens</i> <i>B. subtilis</i>	Dual culture technique	- <i>T. harzianum</i> (UAHS 1) most effective in inhibiting the mycelial growth (80.36%) followed by <i>T. harzianum</i> (UAHS 2) (78.33%).	Mahantesh et al. [17]
16	<i>T. viride</i> <i>P. fluorescens</i> <i>T. harzianum</i> <i>B. subtilis</i>	Dual culture technique	-Maximum mycelial growth inhibition was shown by <i>T. harzianum</i> (65.93%). - <i>B. subtilis</i> showed growth inhibition of pathogen by 58.22% followed by <i>T. viride</i> (41.67%) and <i>P. fluorescens</i> (38.89%).	Rani et al. [18]
17	<i>P. aeruginosa</i> <i>S. aureus</i> <i>B. subtilis</i>	Dual culture technique	- <i>S. aureus</i> showed highest inhibitory effect. - <i>P. aeruginosa</i> , <i>B. subtilis</i> and <i>S. aureus</i> also inhibit the hyphal growth of <i>A. solani</i> .	Chanthini et al. [39]
18	<i>T. harzianum</i> <i>T. viride</i> <i>T. hamatum</i>	Dual culture technique	-Under <i>in vitro</i> conditions, <i>T. hamatum</i> showed highest inhibition, followed by <i>T. harzianum</i> and <i>T. viride</i> .	Sarfraz et al. [14]
19	<i>Serratia proteamaculans</i>	Dual culture technique	- <i>S. proteamaculans</i> significantly suppress the disease severity compared with control. -Bacterial therapy improved plant defense enzyme function and the protein associated with pathogenesis.	Youssef et al. [40]
20	<i>B. subtilis</i>	RCBD	-Early blight disease was reduced significantly by 67–83% by foliar application of <i>B. subtilis</i> either alone or in combination with the plant nutrients.	Awan and Shoaib; [5]
21	<i>T. harzianum</i> <i>T. viride</i> <i>P. fluorescens</i>	Dual culture technique	- <i>P. fluorescens</i> showed maximum inhibition (56.03%) followed by <i>T. harzianum</i> (51.58%) and <i>T. viride</i> (47.78%).	Bais et al. [4]

S. no.	Biocontrol agents	Method of treatment	Result	References
22	<i>Streptomyces puniceus</i> strain L75	Dual culture technique	-Co-inoculation of <i>A. solani</i> with L75 EA extracts reduced induced diseased areas by approximately 48, 90 and 98% respectively relative to the control after 5 days.	Hao et al. [41]
23	<i>Macrolepiota</i> sp.	Dual culture assay	-The cultural fluid of <i>Macrolepiota</i> sp. CS185 against early blight under greenhouse environment inhibit disease up to 46.6%.	Hernández-Ochoa et al. [42]
24	<i>T. viride</i> <i>T. harzianum</i>	Dual culture method	- <i>T. viride</i> was most effective against <i>A. solani</i> .	Roy et al. [10]

Use of Plant Extracts

Several researchers have recently exploited higher plant products as novel chemotherapists in plant defence, which are important sources of new agrochemicals for plant disease control [7]. The natural products are considered to be the best as alternative to synthetic chemicals due to less negative environmental impact [43]. There are number of plants species with natural constituents which are harmful to several plant pathogens [44]. Plant extracts have antimicrobial activity for controlling early blight and other plant diseases [9]. Raza et al. [43] reported that the extract of *Eucalyptus camaldulensis*, *Azadirachta indica*, *Parthenium hysterophorus* and *Datura stramonium* reduced the early blight in the laboratory conditions. Similarly, *Lantana camara* shows effect on the decline of early blight of tomato [45]. Several authors have also documented plants extract effect against early blight disease [46,44,7,3,14,12,9]. *A. sativum* also showed promising results against *A. solani* causing EB [43,1,7]. Seed and leaf extracts of neem exhibit anti-fungal property and are used as

insecticides for controlling agricultural insect pests [43]. Secondary toxic metabolites, such as coumarins, flavonols, tannins, quinones, phenolics and saponins are produced by plants [14], which are significant compounds acknowledged for their activities against a variety of microbes [1]. Because of their essential characteristics such as high molecular diversity, high bioactivity, variability and short durability of climate, essential plant oils from plants show great potential against plant disease [46]. Hong et al. [47] had noteworthy results in essential oils from cinnamon oil and origanum oil vapours against *A. solani* at 6 hours post inoculation. The use of plant extract is environmentally safe, easily biodegradable, cheaper, long-lasting and ideal for organic farming, as well as efficient biocontrol methods for plant disease management. Phytochemicals have been researched for this purpose and used to establish alternatives to chemical pesticides to control plant diseases with a lower risk to the environment and health [7,48]. The various plant extracts used for the control of early blight are presented in Table 3.

Table 3. Plant extracts used against early blight

S no.	Plant extracts	Method of treatment	Result	References
1	<i>Azadirachta indica</i> <i>Ocimum sanctum</i>	Poison food technique	-Tulsi extract reduced disease severity by 40% which is better than neem extract.	Dheeba et al. [49]
2	<i>S. liebmanni</i> <i>C. sertularioides</i> <i>U. lactuca</i> <i>P. gymnospora</i>	Poison food technique	-All sea weed extracts reduced necrotic lesions induced by <i>A. solani</i> . - <i>U. lactuca</i> and <i>P. gymnospora</i> showed better results.	Hernández-Herrera et al. [42]
3	<i>Azadirachta indica</i>	RCBD	-Under field trails, neem extract was observed effective against early blight (55.12%).	Nahak and Sahu, (2014)
4	<i>Azadirachta indica</i> <i>Zingiber officinale</i> Rosc <i>Ocimum sanctum</i> <i>Phyllanthus niruri</i> <i>Artemisia vulgaris</i> <i>Lantana camera</i>	Food poisoned technique	- <i>Zingiber officinale</i> showed better inhibitory effect on sporulation and radial growth of mycelium of <i>A. solani</i> .	Pandey; [23]

S no.	Plant extracts	Method of treatment	Result	References
5	Botanicals	Poisoned food technique	-The most effective plant extract against mycelial growth of <i>A. solani</i> was found to be <i>Jatropha</i> (62.78%).	Roopa et al. [24]
6	Plant extracts	Food poison technique	- <i>Datura stramonium</i> , <i>Azadirachta indica</i> and <i>Withania somnifera</i> showed noteworthy antifungal activities. - <i>W. somnifera</i> showed maximum inhibition of <i>A. solani</i> (62.56%), which was followed by <i>D. stramonium</i> (34.65%) and <i>A. indica</i> (25.27%).	Sahu and Patel, [50]
7	<i>C. gigantea</i> <i>A. indica</i> <i>D. alba</i> <i>A. cepa</i> <i>O. sanctum</i> <i>A. sativum</i>	Poisoned food technique	-Garlic bulb extract (10%) was observed most effective against mycelial growth (35.16%) in controlling disease intensity followed by neem leaf extract (35.68%), datura leaf extract (36.25%) and onion bulb extract (36.71%).	Yadav; [32]
8	<i>Glycyrrhiza glabra</i> <i>Syzygium aromaticum</i> , <i>Hemidesmus indicus</i> <i>Cinnamomum zeylanicum</i> <i>Inula racemosa</i> , <i>Rubia cordifolia</i> , <i>Ferula foetida</i> , <i>Saussurea lappa</i>	Poison food technique/RBD	-At dosage of 3 ml/lit, <i>C. zeylanicum</i> extract exhibited whole inhibition of <i>A. solani</i> <i>in vitro</i> conditions. - At dosage of 2 ml/lit <i>C. zeylanicum</i> was effective against <i>A. solani</i> infesting tomato in shade house trails.	Yeole et al. [51]
9	Plant extracts	Poison food technique	-Out of fifteen plant extracts, eight showed better inhibitory effect on mycelial growth of <i>A. solani</i> . - <i>Tephrosia purpurea</i> (72%) showed best results followed by <i>Capsicum annum</i> (70%) and <i>Gliricidia sepium</i> (70%).	Zafar et al. [52]
10	<i>D. stramonium</i> <i>A. indica</i> <i>L. camara</i>	Poisoned food technique	- <i>D. stramonium</i> (57.03%) was most effective against mycelial growth followed by <i>A. indica</i> (51.35%) and <i>L. camara</i> (48.02%).	Koley et al. [33]
11	<i>Coleus forskohlii</i>		- <i>Coleus forskohlii</i> roots showed inhibitory effect on spore germination of <i>A. solani</i> . - <i>C. forskohlii</i> exhibit dose dependent effect on spore germination.	Nidiry et al. [53]
	<i>Solanaceous phylloplane</i>	Poison food technique	- <i>S. phylloplane</i> significantly controlled early blight in a dose-dependent manner.	Pane and Zaccardelli; [54]
12	Plant extracts	Poison food technique	- <i>E. obliqua</i> was the most effective plant extract inhibiting mycelial growth by 88%. - <i>P. daemia</i> , <i>C. viscosa</i> and <i>P. amarus</i> showed minimum inhibition.	Sadana and Didwania; [29]
13	<i>Eucalyptus citriodora</i> <i>Cymbopogon citratus</i> <i>Rosmarinus officinalis</i> <i>Cymbopogon martinii</i> <i>Citrus sinensis</i> <i>Syzygium aromaticum</i> <i>Piper hispidinervum</i> <i>Melaleuca alternifolia</i> <i>Cinnamomum zeylanicum</i> <i>Mentha piperita</i>	Poison food technique/RCBD	- <i>C. citratus</i> , <i>S. aromaticum</i> and <i>C. zeylanicum</i> oils at concentration of 750 µL/L, <i>M. alternifolia</i> and <i>E. citriodora</i> at 2000 µL/L and <i>Mentha piperita</i> in 5000 µL/L inhibited <i>in vitro</i> conidia germination and mycelial growth completely. -Under greenhouse conditions, <i>C. citratus</i> , <i>C. martinii</i> , <i>C. zeylanicum</i> and <i>S. aromaticum</i> at 750 µL/L inhibited early blight in leaf of tomato. - <i>E. citriodora</i> oil at 5000 µL/L showed similar effects inhibiting early blight of tomato.	Abreu et al. [46]
14	Cinnamon extract	Poison food technique	-Cinnamon extract (42.5%) significantly reduced mycelial growth of <i>A. solani</i> .	Attia et al. [44]
15	<i>Tithonia diversifolia</i> , <i>Azadirachta indica</i> , <i>Euphorbia heterophylla</i> , <i>Carica papaya</i> , <i>Chromolaena odorata</i>	Poison food technique	- <i>C. papaya</i> showed maximum inhibition of mycelial growth. - <i>C. odorata</i> , <i>E. heterophylla</i> and <i>A. indica</i> showed no significant difference in disease severity -High yield was also significantly obtained with <i>C. papaya</i> .	Joseph et al. [38]
16	<i>Carica papaya</i> <i>Ricinus communis</i>	Poison food Technique/	-At 50% concentration, <i>P. juliflora</i> showed 88.9% inhibition of early blight disease.	Kalpashree and Raveesha; [55]

S no.	Plant extracts	Method of treatment	Result	References
	<i>Prosopis juliflora</i> <i>Polyalthia longifolia</i>	Dry Mycelial weight method	-The inhibition of disease shown by <i>Carica papaya</i> was 77.84% which was followed by <i>Ricinus communis</i> (70.5%) and <i>Polyalthia longifolia</i> (63%) by Poisoned Food Technique. - <i>P. juliflora</i> showed 90.7% inhibition, <i>C. papaya</i> (89.47%), <i>Ricinus communis</i> (86.1%) and <i>P. longifolia</i> (79.2%) by Dry mycelial weight method.	
17	<i>Lantana camara</i> <i>Eucalyptus globulus</i> <i>Azadirachta indica</i> <i>Allium cepa</i> <i>Zingiber officinale</i> <i>Allium sativum</i> <i>Parthenium hysterophorus</i>	Poison food technique	- <i>Allium sativum</i> plant extracts recorded lowest colony diameter of <i>Alternaria solani</i> with 87.3% inhibition of growth of fungus over control.	Kumar and Barnwal[45]
18	<i>Allium sativum</i> <i>Parthenium hysterophorus</i> <i>E. camaldulensis</i> <i>Datura stramonium</i> <i>Azadirachta indica</i>	Poison food technique	-Maximum mycelial inhibition was shown by <i>Azadirachta indica</i> plant extract (69.65%). - <i>Allium sativum</i> plant extract showed (66.15%) followed by <i>Parthenium hysterophorus</i> (59.94%), <i>Datura stramonium</i> (49.46%) and <i>Eucalyptus camaldulensis</i> (49.31%).	Raza et al. [43]
19	<i>Eucalyptus globus</i> <i>Datura alba</i> <i>Zingiber officinale</i> <i>Curcuma longa</i> <i>Allium sativum</i> <i>Melia azedarach</i>	Poison food technique	-At 20% concentration, <i>A. sativum</i> extract was most effective against <i>A. solani</i> . - <i>Datura alba</i> and <i>Eucalyptus globus</i> were ineffective, remaining plants extract significantly reduced <i>A. solani</i> growth on PDA.	Ahmad et al. [1]
20	<i>Vitex trifolia</i> <i>Allium sativum</i> <i>Azadirachta indica</i> <i>Acorus calamus</i> <i>Lantana camara</i>	Poison food technique	- <i>Allium sativum</i> showed 82.2%, 94.4% at 1, 2% and 100% at 3 and 4% concentrations. -The other plant extracts showed less than 35% inhibition on growth of the fungus.	Devi et al. [7]
21	Parthenium Lantana Water hyacinth, Pongamia Marigold,Neem, Tulsi, Onion, Garlic	Poison food technique	-Pongamia leaf extract inhibit the mycelial growth of <i>A. solani</i> (54.76%), followed by neem leaf extract (48.27%), lantana leaf extract (47.42%), and garlic bulbs (46.42%). -Least inhibition recorded in water hyacinth (36.75%).	Mahantesh et al. [17]
22	<i>Moringa oleifera</i> chloroform extract and aqueous extract	Poison food Technique	-Relatively high inhibitory effects on the germination of conidia from the lowest to the highest concentration were observed by two plant extracts.	Mvumi et al. [56]
	Clove Sesame Eucalyptus Cinnamon Castor Lemon Mustard Ginger	Poison food technique	-The castor, lemon and ginger at 3% concentration showed maximum radial growth inhibition of <i>A. solani</i> by 27, 29 and 29.6% respectively. -The inhibition shown by lemon oil was 27%, cinnamon oil, 21.6%, mustard oil 21.6%, jojoba 21%, sesame 21% and garlic 20%. -The disease was reduced by 34.9% by clove oil, 34.3% by eucalyptus and 34.2% by garlic, respectively.	Rahmatzai et al. [48]
23	<i>Gelidium serrulatum</i> <i>Ulva lactuca</i> <i>Sargassum filipendula</i>	RCBD	-Sea weed extracts reduced early blight disease efficiently. -Among weeds <i>G. serrulatum</i> showed best results.	Ramkissoon et al. [57]
24	<i>D. stramonium</i> <i>Cannabis sativa</i> <i>Ocimum sanctum</i> <i>A. indica</i>	Poison food technique	- <i>Lantana camara</i> (65.07%) showed maximum mycelial inhibition followed by <i>D. stramonium</i> (63.33%) and <i>A. indica</i> (61.33%). -Minimum mycelial growth inhibition was shown by <i>C. sativa</i> (32.59%).	Rani et al. [18]
25	Garlic bulb extract Datura leaf extract	Poisoned food technique	-The garlic bulb extract and Datura leaf extract at 5% concentration reduced disease intensity by 23.17% and 24.86% respectively.	Bhagat and Zacharia; [3]

S no.	Plant extracts	Method of treatment	Result	References
26	<i>Nigella sativa</i> <i>Trachyspermum ammi</i> <i>Azadirachta indica</i> <i>Eucalyptus globulus</i> <i>Lantana camara</i>	Poison food Technique	-Mycelial growth inhibition follows the order <i>Azadirachta indica</i> (46.66%)> <i>Eucalyptus globulus</i> (40.00%)> <i>Lantana camara</i> (32.78%) under <i>in-vitro</i> conditions. -Minimum disease intensity was recorded with <i>Azadirachta indica</i> (26.99%) and <i>Eucalyptus globulus</i> (27.25%) under <i>in-vivo</i> conditions.	Bhanage et al. [6]
27	<i>A. indica</i> <i>Ocimum sanctum</i> <i>D. stramonium</i> <i>Zingiber officinale</i> <i>Allium sativum</i> <i>Eucalyptus sp.</i>	Poison food Technique	-Significant inhibition of mycelial growth was shown by <i>Allium sativum</i> followed by <i>Azadirachta indica</i> .	Bais et al. [4]
28	<i>Azadirachta indica</i> <i>Ocimum sanctum</i> <i>Allium sativum</i>	RCBD	-Plant extracts were effective when applied to plants grown in greenhouse and field. -Highest reduction in disease incidence was shown by <i>A. sativum</i> (77.42%) and <i>A. indica</i> (62.32%).	Chohan et al. [12]
29	Ashwagandha Nithya Kalyani Tulsi Ginger Garlic Coleus Bougainvillea Eruku Sea weeds	Poisoned food technique	-Garlic extract showed highest inhibition over other treatments followed by bougainvilleas (77.4%), eruku (57.4%) and coleus (55.2%) percent inhibition over control. -Ashwagandha (24.2%) and sea weeds (0%) showed least inhibition.	Gowsik et al. [58]
30	<i>Azadirachta indica</i> <i>Prosopis juliflora</i> <i>Allium cepa</i> <i>Eucalyptus obliqua</i> <i>Pongamia pinnata</i> <i>Murraya koenigii</i> <i>Allium sativum</i> <i>Lawsonia inermis</i> <i>Curcuma longa</i> <i>Zingiber officinale</i>	Poisoned food technique	-Turmeric showed maximum inhibition i.e, 89.44% over control, which was followed by garlic (85.77%) and eucalyptus (73.03%). - <i>C. longa</i> , <i>A. cepa</i> and <i>E. obliqua</i> extracts reduced conidial germination to 91.11, 84.44 and 80.06 per cent respectively over control.	Rex et al. [9]
31	<i>Allium sativum</i> <i>Azadirachta indica</i> <i>Zingiber officinale</i> <i>Ocimum sanctum</i> <i>Adhatoda vasica</i> <i>Datura metel</i>	Poison food technique	-The <i>Allium sativum</i> plant extract was most effective against <i>A. solani</i> , followed by <i>Azadirachta indica</i> .	Roy et al. [10]

CONCLUSION

Tomato is one of the most important vegetable crops economically produced and consumed worldwide. However, tomato is susceptible to a range of biotic agents including a fungus, *A. solani* that causes early blight of tomato. This widespread disease can incur a loss of yields upto 80% of the crop. Hence, it is vital to use various strategies and control measures to control the disease in tomato. A myriad of methods and strategies have been developed and used to control early blight disease in tomato ranging from use of chemicals such as fungicides, application of plant extracts to use of several biological organisms. A number of

researchers have successfully evaluated and shown the efficacy of different fungicides with different concentrations against the disease. However, improper use of chemicals, such as fungicides pollutes the environment and becomes toxic to humans. In recent years, research is focused more on using biological plant extracts and organisms for a toxic free environment. It is important to develop good agricultural practices and efficient management systems to restraint the disease and thus, contribute to economy.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ahmad F, Raziq F, Ullah N, Khan H, Din N. In vitro and in vivo bio-assay of phytobiocidal effect of plant extracts on *Alternaria solani* causing agent of early blight disease in tomato. Archives of Phytopathology and Plant Protection. 2017; 50(11-12):568-83.
2. Chanthini, KMP, Senthil-Nathan S, Soranam R, Thanigaivel A, Karthi S, Sreenath Kumar C, Kingsley SJ and Kanagaraj Murali-Baskaran R. Bacterial compounds, as biocontrol agent against early blight (*Alternaria solani*) and tobacco cut worm (*Spodoptera litura* Fab.) of tomato (*Lycopersicon esculentum* Mill.). Archives of Phytopathology and Plant Protection. 2018;51(13-14):729-753.
3. Bhagat S, Zacharia S. Effect of plant extracts on early blight of tomato (*Lycopersicon esculentum* L.). Journal of Pharmacognosy and Phytochemistry. 2018; 7(5):1804-6.
4. Bais RK, Ratan V, Kumar S, Tiwari A. Comparative analysis of various strategies for management of early blight of tomato incited by *Alternaria solani* (Ellis and Martin) Jones and Grout.
5. Awan ZA, Shoaib A. Combating early blight infection by employing *Bacillus subtilis* in combination with plant fertilizers. Current Plant Biology. 2019; 20:100125.
6. Bhanage S, Patil NB, Singh S. Effect of selected botanicals against early blight (*Alternaria solani*) disease of tomato (*Solanum lycopersicum*) Plant Archives. 2019;19(2):2839-42.
7. Devi NO, Singh NI, Devi RT, Chanu WT. In vitro evaluation of *Alternaria solani* (Ellis and Mart.) Jones and Grout causing fruit rot of tomato by plant extracts and bio-control agents. International Journal of Current Microbiology and Applied Sciences. 2017;6(11):652-61.
8. Verma A, Kumar S, Shina HA, Jaiswal S. Evaluate the efficacy of bio-control agents and botanicals against early blight of potato caused by *Alternaria solani*. Pharma Innov J. 2018;7(3):28-30.
9. Rex B, Prabhu S, Kumar JS. Antifungal efficacies of plant extracts against *Alternaria solani* (Ellis and Martin) Jones and Grout under in vitro condition. Annals of Phytomedicine. 2019;8(1):1-5.
10. Roy CK, Akter N, Sarkar MK, Pk MU, Begum N, Zenat EA, Jahan MA. Control of Early Blight of Tomato Caused by and Screening of Tomato Varieties against the Pathogen. The Open Microbiology Journal. 2019;13(1).
11. Tomazoni EZ, Pansera MR, Pauletti GF, Moura S, Ribeiro RT, Schwambach J. In vitro antifungal activity of four chemotypes of *Lippia alba* (Verbenaceae) essential oils against *Alternaria solani* (Pleosporaceae) isolates. Anais da Academia Brasileira de Ciências. 2016;88(2):999-1010.
12. S, Perveen R, Anees M, Azeem M, Abid M. Estimation of secondary metabolites of indigenous medicinal plant extracts and their in vitro and in vivo efficacy against tomato early blight disease in Pakistan. Journal of Plant Diseases and Protection. 2019;126(6):553-63.
13. Shoaib A, Awan ZA, Khan KA. Intervention of antagonistic bacteria as a potential inducer of disease resistance in tomato to mitigate early blight. Scientia Horticulturae. 2019;252:20-8.
14. Sarfraz M, Khan SA, Moosa A, Farzand A, Ishaq U, Naeem I, Khan WA. Promising antifungal potential of selective botanical extracts, fungicides and trichoderma isolates against *Alternaria solani*. Cercetari Agronomice in Moldova. 2018;51(1):65-74.
15. Farooq S, Jat RR, Gupta A, Singh R, Majeed M, Nabi SU, Bashir N, Shah OS. Evaluation of different fungicides against *Alternaria solani* (Ellis & Martin) Sorauer cause of early blight of tomato under laboratory conditions.
16. Ghazanfar MU, Raza W, Ahmed KS, Qamar J, Haider N, Rasheed MH. Evaluation of different fungicides against *Alternaria solani* (Ellis & Martin) Sorauer cause of early blight of tomato under laboratory conditions. Evaluation. 2016; 1(5).
17. Mahantesh SB, Karegowda C, Kavitha SV, Kavita TH, Punith ND. In vitro evaluation

- of fungicides, bio agents and natural plant extracts against early blight caused by *A. solani*. IJCS. 2017;5(5):1346-50.
18. Rani S, Singh R, Gupta S. Development of integrated disease management module for early blight of tomato in Jammu. Journal of Pharmacognosy and Phytochemistry. 2017; 6(2):268-73.
 19. Sharma RK, Patel DR, Chaudhari DR, Kumar V, Patel MM. Effect of Some Fungicides against Early Blight of Tomato (*Lycopersicon esculentum* Mill.) Caused by *Alternaria solani* (Ell. & Mart.) Jones and Grout and Their Impact on Yield. Int. J. Curr. Microbiol. App. Sci. 2018;7(07): 1395-401.
 20. Egel DS, Hoagland L, Davis J, Marchino C, Bloomquist M. Efficacy of organic disease control products on common foliar diseases of tomato in field and greenhouse trials. Crop Protection. 2019;122:90-7.
 21. Rajeshkumar S. Antifungal impact of nanoparticles against different plant pathogenic fungi. In Nanomaterials in Plants, Algae and Microorganisms. Academic Press. 2019;197-217.
 22. Joseph A, Igbinosa OB, Alori ET, Ademiluyi BO, Aluko AP. Effectiveness of *Pseudomonas* species in the management of tomato early blight pathogen *Alternaria solani*. African Journal of Microbiology Research. 2017;11(23):972-6.
 23. Pandey RK. Studies on management of early blight disease caused by *Alternaria solani* on tomato crops through fungicides, crude plant extracts and bio control agents. Journal of Natural Resource and Development. 2014;9(2):1-2.
 24. Roopa RS, Yadahalli KB, Kavyashree MC. Evaluation of natural plant extracts, antagonists and fungicides against early blight caused by *A. solani* *in vitro*. The Bioscan. 2014;9(3):1309-12.
 25. Saxena A, Sarma BK, Singh HB. Effect of azoxystrobin based fungicides in management of chilli and tomato diseases. Proceedings of the National Academy of Sciences, India Section B: Biological Sciences. 2016;86(2):283-9.
 26. Walke AG, Potdukhe SR, Meena S, Meena AK, Kakralya GL, Choudhary CS. Efficacy of fungicides *in vitro* and *in vivo* of early blight of tomato (*Lycopersicon esculentum*) caused by *Alternaria solani*. The Journal of Plant Science Research. 2014;30(2):123.
 27. Zghair QN, Lal AA, Mane MM, Sobita S. Effect of bioagents and fungicide against early blight disease of tomato (*Lycopersicon esculentum* L.). International Journal of Plant Protection. 2014;7(2):330-3.
 28. Desta M, Woldia PO, Yesuf M. A true endophyte of the endangered tree, *Cynometra travancorica* in the efficacy and economics of fungicides and their application schedule for early blight (*Alternaria solani*) management and yield of tomato at south Tigray, Ethiopia. Journal of Plant Pathology & Microbiology; 2015.
 29. Sadana D, Didwania N. Bioefficacy of fungicides and plant extracts against *Alternaria solani* causing early blight of tomato. In International Conference on Plant, Marine and Environmental Sciences 2015;2:38-42.
 30. Abu-El Samen F, Goussous SJ, Al-Shudifat A, Makhadmeh I. Reduced sensitivity of tomato early blight pathogen (*Alternaria solani*) isolates to protectant fungicides, and implication on disease control. Archives of Phytopathology and Plant Protection. 2016; 49(5-6):120-36.
 31. Abdalla SA, Algarni SA, Ibrahim EA, Naim AM. *In vitro* screening of *Bacillus* isolates for biological control of early blight disease of tomato in shambat soil. World J. Agric. Res. 2014;2(2):47-50.
 32. Yadav R. Eco-friendly management of early blight of potato caused by *Alternaria solani*. Res. Environ. Life Sci. 2014;7(2): 81-84.
 33. Koley S, Mahapatra SS, Kole PC. *In vitro* efficacy of bio-control agents and botanicals on the growth inhibition of *Alternaria solani* causing early leaf blight of tomato. Int. J. Bio-res. Env. Agril. Sci.; 2015.
 34. Lahkar J, Borah SN, Deka S, Ahmed G. Biosurfactant of *Pseudomonas aeruginosa* JS29 against *Alternaria solani*: the causal organism of early blight of tomato. BioControl. 2015;60(3):401-11.

35. Maurya U, Lal EP, Yadav OP, Prakash A, Singh AK. Plant growth promoting rhizobacteria and their activity against early blight of tomato. *Indian Journal of Life Sciences*. 2015;5(1):57.
36. Selim ME. Effectiveness of Trichoderma biotic applications in regulating the related defense genes affecting tomato early blight disease. *J Plant Pathol Microb*. 2015; 6(311):2.
37. Dhal A, Beura SK, Dash SK, Tripathy L, Swain SK, Sethi D. Eco-friendly and integrated approaches for management of early blight disease in tomato. *Int J Curr Microbiol App Sci*. 2017;6(10):3052-7.
38. Joseph A, Igbinosa OB, Alori ET, Ademiluyi BO, Aluko AP. Effectiveness of *Pseudomonas* species in the management of tomato early blight pathogen *Alternaria solani*. *African Journal of Microbiology Research*. 2017;11(23):972-6.
39. Chanthini KM, Senthil-Nathan S, Soranam R, Thanigaivel A, Karthi S, Sreenath Kumar C, Kingsley SJ, Kanagaraj Murali-Baskaran R. Bacterial compounds, as biocontrol agent against early blight (*Alternaria solani*) and tobacco cut worm (*Spodoptera litura* Fab.) of tomato (*Lycopersicon esculentum* Mill.). *Archives of Phytopathology and Plant Protection*. 2018;51(13-14):729-53.
40. Youssef SA, Tartoura KA, Greash AG. Serratia proteamaculans mediated alteration of tomato defense system and growth parameters in response to early blight pathogen *Alternaria solani* infection. *Physiological and Molecular Plant Pathology*. 2018;103:16-22.
41. Hao L, Zheng X, Wang Y, Li S, Shang C, Xu Y. Inhibition of tomato early blight disease by culture extracts of a *Streptomyces puniceus* isolate from mangrove soil. *Phytopathology*. 2019; 109(7):1149-56.
42. Hernández-Herrera RM, Virgen-Calleros G, Ruiz-López M, Zañudo-Hernández J, Délano-Frier JP, Sánchez-Hernández C. Extracts from green and brown seaweeds protect tomato (*Solanum lycopersicum*) against the necrotrophic fungus *Alternaria solani*. *Journal of Applied Phycology*. 2014;26(3):1607-14.
43. Raza W, Ghazanfar MU, Iftikhar Y, Ahmed KS, Haider N, Rasheed MH. Management of early blight of tomato through the use of plant extracts. *Management*. 2016;1(5).
44. Attia MS, Sharaf AE, Zayed AS. Protective action of some bio-pesticides against early blight disease caused by *Alternaria solani* in tomato plant. *JISSET International Journal of Innovative Science, Engineering and Tech*. 2017;4:67-94.
45. Kumar TR and Barnwal M. Evaluation of new fungicides for management of early blight of tomato. *The Bioscan*, 2016;11(4): 2751-2756.
46. Abreu CL, Feres CI, Furtado EL, Ming LC, Marques MO, Acációd RS, Riffel A, Goulart HF, Santana AE, Bernardes FS, Câmara FL. Efficient control of conidium germination, mycelial growth and early blight in tomato in vitro with essential oils under farm conditions. *African Journal of Agricultural Research*. 2016;11(43):4401-12.
47. Hong JK, Jo YS, Ryoo DH, Jung JH, Kwon HJ, Lee YH, Chang SW, Park CJ. *Alternaria* spots in tomato leaves differently delayed by four plant essential oil vapours. *Research in Plant Disease*. 2018;24(4):293.
48. Rahmatzai N, Zaitoun AA, Madkour MH, Ahmady A, Hazim Z, Mousa MA. *In vitro* and in vivo antifungal activity of botanical oils against *Alternaria solani* causing early blight of tomato. *International Journal of Biosciences*. 2017;10(1):91-9.
49. Dheeba B, Niranjana R, Sampathkumar P, Kannan K, Kannan M. Efficacy of neem (*Azadirachta indica*) and tulsi (*Ocimum sanctum*) leaf extracts against early blight of tomato. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*. 2015;85(1):327-36.
50. Sahu DK, Khare CP, Patel R. Eco friendly management of early blight of tomato using botanical plant extracts. *Journal of Industrial Pollution Control*. 2014;30(2): 215-8.
51. Yeole GJ, Teli NP, Kotkar HM, Mendki PS. *Cinnamomum zeylanicum* extracts and their formulations control early blight of

- tomato. Journal of Biopesticides. 2014; 7(2):110.
52. Zafar H, Shaukat SS, Sheikh AH. Detection of antifungal activity of various plant extracts against *Alternaria solani*, the cause of early blight of tomato. International Journal of Biology and Biotechnology (Pakistan); 2014.
 53. Nidiry ES, Ganeshan G, Lokesha AN. Antifungal activity of the extractives of *Coleus forskohlii* roots and Forskolin. Pharmaceutical Chemistry Journal. 2015; 49(9):624-6.
 54. Pane C, Zaccardelli M. Evaluation of *Bacillus* strains isolated from *solanaceous phylloplane* for biocontrol of *Alternaria* early blight of tomato. Biological control. 2015;84:11-8.
 55. Kalpashree MM, Raveesha KA. Efficacy of aqueous plant extracts against *Alternaria solani*, the causative agent of early blight of tomato. International J. Herb. Med. 2016; 4(6):184-8.
 56. Mvumi C, Ngadze E, Marais D, du Toit ES, Mvumi BM. Moringa (*Moringa oleifera*) leaf extracts inhibit spore germination of *Alternaria solani*, causal agent of early blight disease of tomato (*Solanum lycopersicum*). South African Journal of Plant and Soil. 2017;34(3):161-5.
 57. Ramkissoon A, Ramsubhag A, Jayaraman J. Phytoelicitor activity of three Caribbean seaweed species on suppression of pathogenic infections in tomato plants. Journal of Applied Phycology. 2017; 29(6):3235-44.
 58. Gowsik Raja S, Kannan R, Rajinimala N, Sabarinathan KG. *In vitro* evaluation of plant extracts against early blight of tomato caused by *Alternaria solani*. Journal of Pharmacognosy and Phytochemistry. 2019; 8(3):4767-9.