


Метод извлечения математических формул



Surya/pix2tex

LATEX

```
<math display="block">\begin{split} & \& \rho''' x_1 \\ & = p_{12} \setminus, dp_{13} \setminus - p_{13} \setminus, dp_{12} \setminus \equiv \\ & (p_3 \setminus, p_{1'} \setminus, p_{2'} \setminus) \setminus \& \rho''' x_2 \\ & \equiv p_{12} \setminus, dp_{23} \setminus - p_{23} \setminus, dp_{12} \setminus \\ & \equiv (p_3 \setminus, p_{3'} \setminus, p_{2'} \setminus) \setminus \& \rho''' x_3 \\ & \equiv p_{12} \setminus, dp_{23} \setminus - p_{23} \setminus, dp_{12} \setminus \\ & \equiv (p_3 \setminus, p_{1'} \setminus, p_{2'} \setminus) \setminus \& \rho''' x_4 \\ & \equiv p_{12} \setminus, dp_{12} \setminus + p_{14} \setminus, dp_{13} \setminus + \\ & p_{14} \setminus, dp_{23} \setminus \equiv -p_{12} \setminus, dp_{34} \setminus - \\ & p_{23} \setminus, dp_{14} \setminus \equiv (p_3 \setminus, p_{1'} \setminus, p_{2'} \setminus) \setminus \end{split}</math>
```

KaTeX

$$\begin{aligned} \rho''' x_1 &\equiv p_{12} dp_{13} - p_{13} dp_{12} \equiv (p_3 p'_1 p'_2) p_1 \\ \rho''' x_2 &\equiv p_{12} dp_{23} - p_{23} dp_{12} \equiv (p_3 p'_3 p'_2) p_2 \\ \rho''' x_3 &\equiv p_{13} dp_{23} - p_{23} dp_{13} \equiv (p_3 p'_1 p'_2) p_3 \\ \rho''' x_4 &\equiv p_{34} dp_{12} + p_{14} dp_{13} - p_{12} dp_{34} - p_{23} dp_{14} \equiv (p_3 p'_1 p'_2) p_{47} \end{aligned}$$

Ключевые этапы модуля:

1. formula_pipeline
2. crop по bbox
3. Surya -> LaTeX
4. fallback: pix2tex -> LaTeX
5. FormulaBlockResult
6. PageContent
7. KaTeX в UI