2/26/25, 5:37 PM hw5csds343

CSDS 343 HW5


1

a

Construct a FSA equivalent to the REG EX


$$1[((01)^* + 1 + 01)^* + 1]^*0$$

The above REG EX is equivalent to this regex: $1(01+1)^*0$ Below is the FSA equivalent to the REG EX



b

Construct a REG EX equivalent to the FSA



Regex Equivalent: $[1 + 0(11)^*(0 + 10)]^*01(11)^*$

2

Prove the reverse of a regular language is also a regular language, (A,A^R)

There are 2 ways to prove this true (please grade using the 2nd method):

If A is a regular language then there exists a FSA F that can decide A. A^R would then be decided by the transpose of F, that is F^T can decide A^R . This is done by having the q_0 of F be the accept state of F^T and the various accept states of F are the initial states of F^T . However we cannot have multiple initial states of F^T , so to circumvent this we create a q_0 that ϵ transitions to each of the accept states of F.

Alternatively, if A is a regular language then there exists a regular expression for A. A^R would then be decided by the reverse of the regular expression. We can prove this by showing the reverses of each regular expression rule:

- 1. $\epsilon \to^R \epsilon$ holds true
- 2. $a \rightarrow^R a$, where $a \in \Sigma$, holds true
- 3. $(x \cup y) \to^R (x^R \cup y^R)$, where x and y are regular expressions, holds true
- 4. $(xy) \rightarrow^R (y^R x^R)$, where x and y are regular expressions, holds true
- 5. $x^* \rightarrow^R x^*$, where x is a regular expression, holds true

Thus since we have created a reverse function \to^R we can show that A^R is a regular language by reversing the regular expression for A and using the reverse regex for strings in A^R .


3

localhost:5613 2/4

2/26/25, 5:37 PM hw5csds343

Prove $B = \{w \in \Sigma_3^* | \text{the bottom row of w is the sum of the top 2 rows} \}$ is a regular language. Hint prove B^R is regular.

We can show this language is regular by proving its reverse is regular. A language can be proven regular if it can be decided by a FSA.



How this FSA works is we alternate between q0 (no carry) and q1 (carry over) if there is a 1 to carry over from the previous string. Then we only accept if there are no more columns to read. Note that any columns not mentioned at a particular state lead to a reject. This FSA will recognize any string in B^R .

Thus sinse ${\cal B}^R$ is a regular language we know that ${\cal B}$ is a regular language by question 2.

4

Prove the following languages are not regular languages:

a

Palindromes over the alphabet $\Sigma = \{a, b, c\}$

localhost:5613 3/4

2/26/25, 5:37 PM hw5csds343

We can show this by proof by contradiction of the pumping lemma.

Suppose palindromes are regular then $\exists p$ s.t. all strings s in L with |s|>p can be divided into s=xyz such that $|xy|\leq p$ and |y|>0

Given the string a^pba^p is a valid palindrome, we can divide it into string $xy=a^p$ because $|xy|=|a^p|< p$ and |y|=0.

The pumping lemma suggests xy^kz is a palindrome for all $k\in\mathbb{N}$

But what about xy^2z ? In order for $xy^2z=a^{p+|y|}ba^p$ to be a palindrome p+|y|=p so |y|=0 which is a contradiction because |y|>0.

Therefore palindromes are not regular.

b

The language $L = \{abbaaabbbb...a^{k-2}b^{k-1}a^k|k \geq 3\}$

We can show this by proof by contradiction of the pumping lemma.

Suppose this language is regular. The the reverse of this language (L^R) is also regular. If L^R is regular then $\exists p$ s.t. all strings s in L with |s|>p can be divided into s=xyz such that $|xy|\leq p$ and |y|>0

Given the string $a^pb^{p-1}a^{p-2}\ldots\in L^R$, we can divide it into string $xy=a^p$ because $|xy|=|a^p|< p$ and |y|=0.

The pumping lemma suggests $xy^kz\in L^R$ for all $k\in\mathbb{N}$

But what about xy^2z ? In order for $xy^0z=a^{p-|y|}b^{p-1}a^{p-2}$ to be in L p-|y|=p so |y|=0 which is a contradiction because |y|>0.

Therefore L is not regular because its reverse is not regular

localhost:5613 4/4