

# ¶ Stat 405 Final

Meera Borle, Isis Burgos, Naomi Consiglio, Carson Foster, Aidan Gerber

December 10th, 2022

## Introduction

Our main dataset is a mortality dataset from the National Bureau of Economic Research. There are new versions of it each year and we are using the 2019 version. Each row is one death based on information filed in death certificates in the 50 states plus Washington DC. Columns are factors regarding the status of persons involved in deaths that can be gleaned from public death certificates. The dataset is focused on the different causes of death. Notable information the dataset contains is education, sex, age classification, day of month, place of death, weekday, manner of death, cause of death, and different risk factors that the deceased had.

## Secondary Dataset

For our secondary dataset, we are using the Behavioral Risk Factor Surveillance System Survey. This survey includes different free text survey questions from across the United States and territories with responses broken out by subgroup. There is also information on sample size, percent affirmative response, and confidence interval bounds. We will combine the secondary dataset by matching up subgroups between the death dataset and the risk factor dataset and trying to use aggregate statistics to analyze how risk factors can be matched with causes of death.


# Questions


We are interested in what different factors are correlated with higher death rates based on cause of death, age groups, and weekdays. Depending on the different causes, different policies can be recommended. Moreover, we can break down the different causes of death by demographics such as race, age, and sex to determine where resources should specifically be directed.

## Exploration

### Deaths by Weekday


First, we plotted weekday of death versus death counts. There were the most deaths on Tuesday. However, days have an average of 7839.79 deaths and 2019 had an extra Tuesday so adjusting for that, the most deaths were on Fridays.





## Deaths by Month



The most deaths occur in the coldest and darkest months of the year which are February, January, December, and March. Summer months have lower deaths by around 10-11%.



## Deaths by Age

Next, we plotted age versus death counts. Deaths were most prevalent among older age groups such as those between 70 and 84, although deaths start increasing more quickly at age 60. There is also a spike in those less than 1 day old. However, those greater than 1 day old do not frequently die.

Deaths by Age



## Deaths by Manner

Here, we plotted the manner of death versus age and counted how many people of a certain aged died based on a certain manner of death. A few key finding of this analysis shows that the majority of people die from natural causes, especially those aged 60+ and less than 1 day old, and accidental causes, spanning across all age groups. What this plot may help to inform us about is the behavior and activities that people in a general age group may commonly engage in that may have lead to their manner of passing. By being observant of the manners of death based on age group, preventative methods can be used to decrease the number of accidental related deaths if we are able to determine commonly engaged activities for age groups. Using this plot will help us answer the cause of death among the different age groups, and further promote research in what actual activities people are participating in that lead to their manner of death.


Under 1 hour  
1-23 hours  
1 day  
2 days  
3 days  
4 days  
5 days  
6 days  
7-13 days  
14-20 days  
21-27 days  
1 month  
2 months  
3 months  
4 months  
5 months  
6 months  
7 months  
8 months  
9 months  
10 months  
11 months  
1 year  
2 years  
3 years  
4 years  
5 years  
6 years  
7 years  
8 years  
9 years  
10 years  
11 years  
12 years  
13 years  
14 years  
15 years  
16 years  
17 years  
18 years  
19 years  
20 years  
21 years  
22 years  
23 years  
24 years  
25 years  
26 years  
27 years  
28 years  
29 years  
30 years  
31 years  
32 years  
33 years  
34 years  
35 years  
36 years  
37 years  
38 years  
39 years  
40 years  
41 years  
42 years  
43 years  
44 years  
45 years  
46 years  
47 years  
48 years  
49 years  
50 years  
51 years  
52 years  
53 years  
54 years  
55 years  
56 years  
57 years  
58 years  
59 years  
60 years  
61 years  
62 years  
63 years  
64 years  
65 years  
66 years  
67 years  
68 years  
69 years  
70 years  
71 years  
72 years  
73 years  
74 years  
75 years  
76 years  
77 years  
78 years  
79 years  
80 years  
81 years  
82 years  
83 years  
84 years  
85 years  
86 years  
87 years  
88 years  
89 years  
90 years  
91 years  
92 years  
93 years  
94 years  
95 years  
96 years  
97 years  
98 years  
99 years  
100 years  
Age not stated

Age




## Deaths by Age and Gender

In this plot, we plotted the number of deaths versus age ranges while demonstrating how many men compared to women passed away in each age category. In each of the bars, the red fill represents the amount of women who passed away in that particular age range while the blue accounts for the amount of men. The percentage seen in each bar represents the proportion of men in a given age range that passed away. This analysis shows that the majority of people under the age of 80 who pass away tend to be men, as nearly every bar from ages 0-80 shows the proportion of male deaths to be above 50%. This proportion of male deaths goes down after 80 years of age, and is likely because women who are of an older age tend to live to a complete life expectancy. What this plot may help to inform us about is the differences in male and females lives and life expectancies. Further research into differences in lifestyle choices for men versus women as a whole may help better explain why women tend to live longer than men. Furthermore, this plot accompanied with a plot on cause of death by gender, may assist in determining what kind of, potentially more risky, behaviors men may partake in during their lifetimes that lead to an earlier death than women.




## **Cause of Death by Education**

For most causes of death, level of education does not have an impact on what proportion of people have that cause of death. The largest difference belongs to “Certain conditions originating in the perinatal period” with high occurrences in those with 8th grade or less education and those with unknown education and nearly no occurrences in all others. Another large proportion difference is in “Congenital malformations” where 8th grade or less has a much higher mortality proportion than other education levels. For causes of death that are not highly tied to conditions at birth, “Syphilis” and “Assault (homicide)” have the highest differing proportions. “Syphilis” has a quite small sample size but unknown education has the highest mortality proportion. For “Assault (homicide)”, 9 - 12th grade, no diploma has the highest mortality proportion.



## Free Text Analysis: Selected Health Conditions by Age Group

Using regular expressions, we polled the BRFSS data set for questions related to heart conditions, cancer, and depression, while grouping by the age of respondents. Furthermore, we restricted entries to those where participants responded positively, indicating that they did have those conditions. Interestingly, all age groups except 65+ had a high incidence of depression, hovering around the 20% mark. This dips significantly to 15% for the 65+ age group, perhaps because mental health was more stigmatized during their lives and psychological diagnoses were less readily available. In addition, the orange and olive green lines (for heart attack and coronary heart disease, respectively) have a significant degree of overlap, which makes sense, given the conditions. The green line (corresponding to non-skin cancer), is slightly higher than the blue line (corresponding to skin cancer) for all age groups. Furthermore, both cancer have positive slopes, indicating that as your age increases, you become more likely to be diagnosed with cancer.



#### Question

- Ever told you had a heart attack (myocardial infarction)?
- Ever told you had angina or coronary heart disease?
- Ever told you had any other types of cancer?
- Ever told you had skin cancer?
- Ever told you that you have a form of depression?