


d'Alembert's Method and the One-Dimensional Heat Equation

روش دالامبر و معادله انتقال حرارت یک بعدی



In this chapter

5.1 d'Alembert's Method for the One-Dimensional Wave Equation	2
5.2 The Heat Equation	3
5.3 Solving the One-Dimensional Heat Equation	4
5.4 The One-Dimensional Heat Equation on an Infinite Rod	6
5.5 The Error Function	8

5.1 d'Alembert's Method for the One-Dimensional Wave Equation

$$\begin{cases} \frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2} \\ u(0, t) = u(L, t) = 0 \\ u(x, 0) = f(x) \\ u_t(x, 0) = g(x) \end{cases}$$

We rewrite and solve the equation using the following change of variables.

$$u(x, t) \Rightarrow u(\varphi, \psi), \quad \begin{cases} \varphi = x + ct \\ \psi = x - ct \end{cases}$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial \varphi} \frac{\partial \varphi}{\partial x} + \frac{\partial u}{\partial \psi} \frac{\partial \psi}{\partial x} = \frac{\partial u}{\partial \varphi} + \frac{\partial u}{\partial \psi}$$

$$\frac{\partial u}{\partial t} = \frac{\partial u}{\partial \varphi} \frac{\partial \varphi}{\partial t} + \frac{\partial u}{\partial \psi} \frac{\partial \psi}{\partial t} = c \frac{\partial u}{\partial \varphi} - c \frac{\partial u}{\partial \psi}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial u}{\partial \varphi} + \frac{\partial u}{\partial \psi} \right) = \left(\frac{\partial^2 u}{\partial \varphi^2} + \frac{\partial^2 u}{\partial \varphi \partial \psi} \right) \cdot 1 + \left(\frac{\partial^2 u}{\partial \psi \partial \varphi} + \frac{\partial^2 u}{\partial \psi^2} \right) \cdot 1 = \frac{\partial^2 u}{\partial \varphi^2} + 2 \frac{\partial^2 u}{\partial \varphi \partial \psi} + \frac{\partial^2 u}{\partial \psi^2}$$

$$\text{(using } \frac{\partial^2 u}{\partial \varphi \partial \psi} = \frac{\partial^2 u}{\partial \psi \partial \varphi} \text{.)}$$

Exercise 5.1

In the same way, prove the identity

$$\frac{\partial^2 u}{\partial t^2} = c^2 \left(\frac{\partial^2 u}{\partial \varphi^2} - 2 \frac{\partial^2 u}{\partial \varphi \partial \psi} + \frac{\partial^2 u}{\partial \psi^2} \right)$$

Substituting the two relations above into the one-dimensional wave equation:

$$c^2 \left(\frac{\partial^2 u}{\partial \varphi^2} - 2 \frac{\partial^2 u}{\partial \varphi \partial \psi} + \frac{\partial^2 u}{\partial \psi^2} \right) = c^2 \left(\frac{\partial^2 u}{\partial \varphi^2} + 2 \frac{\partial^2 u}{\partial \varphi \partial \psi} + \frac{\partial^2 u}{\partial \psi^2} \right) \Rightarrow \frac{\partial^2 u}{\partial \varphi \partial \psi} = 0$$

The general solution:

$$\begin{aligned} \frac{\partial u}{\partial \varphi} = \phi_1(\varphi) &\Rightarrow u(\varphi, \psi) = \int \phi_1(\varphi) d\varphi + \Psi(\psi) = \Phi(\varphi) + \Psi(\psi) \Rightarrow u(x, t) \\ &= \Phi(x + ct) + \Psi(x - ct) \end{aligned}$$

Applying the initial conditions:

$$u(x, 0) = \Phi(x) + \Psi(x) = f(x)$$

$$u_t(x, t) = \frac{\partial \Phi}{\partial \varphi} \frac{\partial \varphi}{\partial t} + \frac{\partial \Psi}{\partial \psi} \frac{\partial \psi}{\partial t} = c\Phi'(\varphi) - c\Psi'(\psi) = c\Phi'(x + ct) - c\Psi'(x - ct)$$

$$u_t(x, 0) = c\Phi'(x) - c\Psi'(x) = c[\Phi'(x) - \Psi'(x)] = g(x)$$

$$\begin{cases} \Phi(x) + \Psi(x) = f(x) \\ \Phi'(x) - \Psi'(x) = \frac{1}{c}g(x) \end{cases} \Rightarrow \begin{cases} \Phi'(x) + \Psi'(x) = f'(x) \\ \Phi'(x) - \Psi'(x) = \frac{1}{c}g(x) \end{cases} \Rightarrow \begin{cases} \Phi'(x) = \frac{1}{2}f'(x) + \frac{1}{2c}g(x) \\ \Psi'(x) = \frac{1}{2}f'(x) - \frac{1}{2c}g(x) \end{cases}$$

$$\begin{cases} \Phi(x) = \frac{1}{2}f(x) + \frac{1}{2c} \int_0^x g(s) ds + k_1 \\ \Psi(x) = \frac{1}{2}f(x) - \frac{1}{2c} \int_0^x g(s) ds + k_2 \end{cases}$$

$$u(x, t) = \frac{1}{2} \left(f(x + ct) + \frac{1}{c} \int_0^{x+ct} g(s) ds \right) + \frac{1}{2} \left(f(x - ct) - \frac{1}{c} \int_0^{x-ct} g(s) ds \right) + k_1 + k_2$$

★ **Key result 5.2** | d'Alembert's formula

$$u(x, t) = \frac{1}{2} [f(x + ct) + f(x - ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds + K, \quad u(0, 0) = 0 \Rightarrow K = 0$$

$$u(x, t) = \frac{1}{2} [f(x + ct) + f(x - ct)] + \frac{1}{2c} \int_{x-ct}^{x+ct} g(s) ds$$

✎ **Exercise 5.3**

Prove that if the boundary conditions are to hold, the functions $f(x)$ and $g(x)$ in the formula above must both be periodic with period $2L$ and odd.

5.2 The Heat Equation

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}, \quad \frac{\partial u}{\partial t} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right), \quad \frac{\partial u}{\partial t} = c^2 \left(\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right)$$

■ The One-Dimensional Heat Equation




Figure 5.1 A one-dimensional rod $0 \leq x \leq L$ with temperature distribution $u(x, t)$

For a unique solution we need boundary conditions and initial conditions:

- Boundary conditions: $u(0, t) = u(L, t) = 0$
- Initial conditions: $u(x, 0) = f(x)$

5.3 Solving the One-Dimensional Heat Equation

$$\begin{cases} \frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \\ u(0, t) = u(L, t) = 0 \\ u(x, 0) = f(x) \end{cases}$$

Here we solve the equation by **separation of variables**. Assume $u(x, t) = X(x) \times T(t)$.

$$\frac{\partial^2 u}{\partial x^2} = X''(x)T(t), \quad \frac{\partial u}{\partial t} = X(x)T'(t) \quad \Rightarrow \quad X(x)T'(t) = c^2 X''(x)T(t) \quad \Rightarrow \quad \frac{T'}{c^2 T} = \frac{X''}{X} = k$$

k must necessarily be negative (for $k = 0$ and $k > 0$ one gets $X(x) = 0$ exactly as in the wave problem). With $k = -\lambda^2$:

$$\frac{T'}{c^2 T} = \frac{X''}{X} = -\lambda^2 \quad \Rightarrow \quad \begin{cases} X'' + \lambda^2 X = 0 \\ T' + c^2 \lambda^2 T = 0 \end{cases}$$

$$X(x) = A \cos \lambda x + B \sin \lambda x, \quad X(0) = 0 \Rightarrow A = 0, \quad X(L) = 0 \Rightarrow B \sin \lambda L = 0 \Rightarrow \lambda = \frac{n\pi}{L} = \lambda_n$$

$$X_n(x) = B_n \sin \lambda_n x, \quad T' + c^2 \lambda_n^2 T = 0 \Rightarrow T_n(t) = C_n e^{-(c\lambda_n)^2 t}$$

$$u_n(x, t) = X_n(x) \times T_n(t) = B_n \sin\left(\frac{n\pi}{L}x\right) \times C_n e^{-(\frac{cn\pi}{L})^2 t}$$

$$u(x, t) = \sum_{n=1}^{\infty} A_n e^{-(c\lambda_n)^2 t} \sin(\lambda_n x), \quad \lambda_n = \frac{n\pi}{L}$$

The initial condition $u(x, 0) = f(x)$:

$$u(x, 0) = \sum_{n=1}^{\infty} A_n \sin \lambda_n x = f(x)$$

$f(x)$ has been set equal to a periodic odd function, so to compute the unknown Fourier coefficients we must regard $f(x)$ as a periodic odd function (with period $2L$):

$$A_n = \frac{2}{2L} \int_{-L}^L f(x) \sin\left(\frac{2\pi}{2L}nx\right) dx = \frac{1}{L} \int_{-L}^L f(x) \sin\left(\frac{n\pi}{L}x\right) dx = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi}{L}x\right) dx$$

★ **Key result 5.4** | Solution of the one-dimensional heat equation (separation of variables)

$$u(x, t) = \sum_{n=1}^{\infty} A_n e^{-(c\lambda_n)^2 t} \sin(\lambda_n x), \quad \lambda_n = \frac{n\pi}{L}, \quad A_n = \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi}{L}x\right) dx$$

Example 5.5 | Example 1

Find the solution of the following one-dimensional heat equation.

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad u(0, t) = u(10, t) = 0, \quad u(x, 0) = f(x) = \begin{cases} x & 0 \leq x < 5 \\ 10 - x & 5 \leq x \leq 10 \end{cases}$$

$$\begin{aligned} A_n &= \frac{2}{L} \int_0^L f(x) \sin\left(\frac{n\pi}{L}x\right) dx = 0.2 \int_0^5 x \sin\left(\frac{n\pi}{10}x\right) dx + 0.2 \int_5^{10} (10 - x) \sin\left(\frac{n\pi}{10}x\right) dx \\ &= 0.2 \left[(x) \left(\frac{-10}{n\pi} \cos \frac{n\pi}{10}x \right) - (1) \left(\frac{-100}{n^2\pi^2} \sin \frac{n\pi}{10}x \right) \right]_0^5 \\ &\quad + 0.2 \left[(10 - x) \left(\frac{-10}{n\pi} \cos \frac{n\pi}{10}x \right) - (-1) \left(\frac{-100}{n^2\pi^2} \sin \frac{n\pi}{10}x \right) \right]_5^{10} = \frac{40}{n^2\pi^2} \sin \frac{n\pi}{2} \\ u(x, t) &= \sum_{n=1}^{\infty} \left(\frac{40}{n^2\pi^2} \sin \frac{n\pi}{2} \right) e^{-(\frac{cn\pi}{10})^2 t} \sin\left(\frac{n\pi}{10}x\right) \end{aligned}$$

(with $c = 1$.)

Example 5.6 | Example 2 (nonzero boundary values)

Find the solution of the following one-dimensional heat equation.

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}, \quad u(0, t) = T_0, \quad u(L, t) = T_1, \quad u(x, 0) = f(x)$$

$$u(x, t) = w(x, t) + v(x) \quad \Rightarrow \quad \begin{cases} w(0, t) + v(0) = T_0 \\ w(L, t) + v(L) = T_1 \end{cases}$$

$$\frac{\partial w}{\partial t} = c^2 \left(\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 v}{\partial x^2} \right)$$

$$\begin{cases} \frac{\partial^2 v}{\partial x^2} = 0 \\ v(0) = T_0 \\ v(L) = T_1 \end{cases} \Rightarrow v(x) = \left(\frac{T_1 - T_0}{L} \right) x + T_0, \quad \begin{cases} \frac{\partial w}{\partial t} = c^2 \frac{\partial^2 w}{\partial x^2} \\ w(0, t) = w(L, t) = 0 \end{cases} \Rightarrow w(x, t)$$

$$= \sum_{n=1}^{\infty} A_n e^{-(c\lambda_n)^2 t} \sin(\lambda_n x)$$

$$u(x, t) = \underbrace{\sum_{n=1}^{\infty} A_n e^{-(c\lambda_n)^2 t} \sin(\lambda_n x)}_{w(x, t)} + \underbrace{\left(\frac{T_1 - T_0}{L} \right) x + T_0}_{v(x)}, \quad \lambda_n = \frac{n\pi}{L}$$

The initial condition:

$$\begin{aligned} \sum_{n=1}^{\infty} A_n \sin \frac{n\pi}{L} x &= f(x) - \left(\frac{T_1 - T_0}{L} \right) x - T_0 \triangleq f_1(x) \Rightarrow A_n \\ &= \frac{2}{L} \int_0^L \left[f(x) - \left(\frac{T_1 - T_0}{L} \right) x - T_0 \right] \sin \frac{n\pi}{L} x dx \end{aligned}$$

Example 5.7 | Example 3 (insulated ends)

Find the solution of the following one-dimensional heat equation.

$$\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}, \quad u_x(0, t) = u_x(L, t) = 0, \quad u(x, 0) = f(x)$$

We solve this problem by **separation of variables**: $u(x, t) = X(x) \times T(t)$.

$$\frac{T'}{c^2 T} = \frac{X''}{X} = -\lambda^2 \Rightarrow \begin{cases} X'' + \lambda^2 X = 0 \Rightarrow X(x) = A \cos \lambda x + B \sin \lambda x \\ T' + c^2 \lambda^2 T = 0 \end{cases}$$

$$\begin{aligned} X'(x) &= -A\lambda \sin \lambda x + B\lambda \cos \lambda x, & X'(0) &= 0 \Rightarrow B = 0, & X'(L) &= 0 \Rightarrow -A\lambda \sin \lambda L \\ &= 0 \Rightarrow \lambda L = n\pi \Rightarrow \lambda_n &= \frac{n\pi}{L} \end{aligned}$$

$$X_n(x) = A_n \cos \lambda_n x, \quad T_n(t) = C_n e^{-(c\lambda_n)^2 t}$$

$$u(x, t) = \sum_{n=1}^{\infty} B_n e^{-(c\lambda_n)^2 t} \cos(\lambda_n x), \quad \lambda_n = \frac{n\pi}{L}$$

The initial condition $u(x, 0) = \sum_{n=1}^{\infty} B_n \cos\left(\frac{n\pi}{L}x\right) = f(x)$ gives

$$B_n = \frac{2}{L} \int_0^L f(x) \cos\left(\frac{n\pi}{L}x\right) dx$$

5.4 The One-Dimensional Heat Equation on an Infinite Rod

$$\begin{cases} \frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \\ u(\infty, t) = u(-\infty, t) = 0 \\ u(x, 0) = f(x), \quad -\infty < x < \infty \end{cases}$$

Assuming $u(x, t) = X(x) \times T(t)$:

$$\begin{aligned} X(x)T'(t) &= c^2 X''(x)T(t) \Rightarrow \frac{T'}{c^2 T} = \frac{X''}{X} = -\lambda^2 \\ \Rightarrow \begin{cases} X(x) = A_1(\lambda) \cos \lambda x + B_1(\lambda) \sin \lambda x = X_\lambda(x) \\ T(t) = C_1(\lambda) e^{-c^2 \lambda^2 t} = T_\lambda(t) \end{cases} \end{aligned}$$

$$\begin{aligned} u_\lambda(x, t) &= [A(\lambda) \cos \lambda x + B(\lambda) \sin \lambda x] e^{-c^2 \lambda^2 t} \Rightarrow u(x, t) \\ &= \frac{1}{\pi} \int_0^\infty [A(\lambda) \cos \lambda x + B(\lambda) \sin \lambda x] e^{-c^2 \lambda^2 t} d\lambda \end{aligned}$$

Checking the equation:

$$\begin{aligned} \frac{\partial^2 u}{\partial x^2} &= \frac{1}{\pi} \int_0^\infty -\lambda^2 [A \cos \lambda x + B \sin \lambda x] e^{-c^2 \lambda^2 t} d\lambda, & \frac{\partial u}{\partial t} \\ &= \frac{1}{\pi} \int_0^\infty -c^2 \lambda^2 [A \cos \lambda x + B \sin \lambda x] e^{-c^2 \lambda^2 t} d\lambda \Rightarrow \frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2} \checkmark \end{aligned}$$

Applying the initial condition: $u(x, 0) = \frac{1}{\pi} \int_0^\infty [A(\lambda) \cos \lambda x + B(\lambda) \sin \lambda x] d\lambda = f(x)$, i.e. the Fourier

integral (Chapter 3):

$$\begin{aligned}
 A(\lambda) &= \mathfrak{F}_C\{f(x)\} = \int_{-\infty}^{\infty} f(x) \cos \lambda x \, dx, & B(\lambda) &= \mathfrak{F}_S\{f(x)\} = \int_{-\infty}^{\infty} f(x) \sin \lambda x \, dx \\
 u(x, t) &= \frac{1}{\pi} \int_0^{\infty} \left[\left(\int_{-\infty}^{\infty} f(v) \cos \lambda v \, dv \right) \cos \lambda x + \left(\int_{-\infty}^{\infty} f(v) \sin \lambda v \, dv \right) \sin \lambda x \right] e^{-c^2 \lambda^2 t} \, d\lambda \\
 &= \frac{1}{\pi} \int_0^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) [\cos \lambda v \cos \lambda x + \sin \lambda v \sin \lambda x] \, dv \right\} e^{-c^2 \lambda^2 t} \, d\lambda \\
 &= \frac{1}{\pi} \int_0^{\infty} \left\{ \int_{-\infty}^{\infty} f(v) \cos \lambda (x - v) \, dv \right\} e^{-c^2 \lambda^2 t} \, d\lambda
 \end{aligned}$$

★ **Key result 5.8** | First form of the solution

$$u(x, t) = \frac{1}{\pi} \int_{-\infty}^{\infty} f(v) \left\{ \int_0^{\infty} e^{-c^2 \lambda^2 t} \cos \lambda (x - v) \, d\lambda \right\} \, dv$$

Mathematical Lemmas

☺ **Theorem 5.9** | Lemma (a)

$$I = \int_0^{\infty} e^{-x^2} \, dx = \frac{\sqrt{\pi}}{2}$$

✓ **Proof.**

$$\begin{aligned}
 I^2 &= \int_0^{\infty} e^{-x^2} \, dx \times \int_0^{\infty} e^{-y^2} \, dy = \int_0^{\infty} \int_0^{\infty} e^{-(x^2+y^2)} \, dx \, dy \stackrel{\rho^2=x^2+y^2}{=} \int_0^{\pi/2} \int_0^{\infty} \rho e^{-\rho^2} \, d\rho \, d\theta \\
 &= \int_0^{\pi/2} \left[\frac{-1}{2} e^{-\rho^2} \right]_0^{\infty} \, d\theta = \frac{\pi}{4}
 \end{aligned}$$

☺ **Theorem 5.10** | Lemma (b)

$$\int_0^{\infty} e^{-s^2} \cos(2bs) \, ds = \frac{\sqrt{\pi}}{2} e^{-b^2}$$

✓ **Proof.**

$$I(b) = \int_0^{\infty} e^{-s^2} \cos(2bs) \, ds \Rightarrow \frac{dI(b)}{db} = - \int_0^{\infty} 2se^{-s^2} \sin(2bs) \, ds$$

Integrating by parts ($du = -2se^{-s^2} \, ds$, $v = \sin 2bs$):

$$\begin{aligned}
 \frac{dI(b)}{db} &= e^{-s^2} \sin 2bs \Big|_0^{\infty} - 2b \int_0^{\infty} e^{-s^2} \cos 2bs \, ds = -2b I(b) \Rightarrow I(b) = k e^{-b^2}, \quad I(0) = k \\
 &= \int_0^{\infty} e^{-s^2} \, ds = \frac{\sqrt{\pi}}{2}
 \end{aligned}$$

With the substitution $c\lambda\sqrt{t} = s$:

$$\int_0^\infty e^{-c^2\lambda^2 t} \cos \lambda(x-v) d\lambda = \frac{1}{c\sqrt{t}} \int_0^\infty e^{-s^2} \cos \left[2 \frac{(x-v)}{2c\sqrt{t}} s \right] ds = \frac{1}{c\sqrt{t}} \frac{\sqrt{\pi}}{2} e^{-\left(\frac{x-v}{2c\sqrt{t}}\right)^2}$$

★ **Key result 5.11** | Second form of the solution

$$u(x, t) = \frac{1}{2c\sqrt{\pi t}} \int_{-\infty}^\infty f(v) e^{-\frac{(x-v)^2}{4c^2 t}} dv$$

With the substitution $z = \frac{v-x}{2c\sqrt{t}}$ (i.e. $v = x + 2c\sqrt{t}z$ and $dv = 2c\sqrt{t}dz$):

★ **Key result 5.12** | Third form of the solution

$$u(x, t) = \frac{1}{\sqrt{\pi}} \int_{-\infty}^\infty f(x + 2c\sqrt{t}z) e^{-z^2} dz$$

5.5 The Error Function

In mathematics the error function is most often defined as

$$\operatorname{erf}(x) \triangleq \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

Properties of the error function:

- odd symmetry: $\operatorname{erf}(-x) = -\operatorname{erf}(x)$
- asymptotic property: $\operatorname{erf}(\infty) = 1$
- its Maclaurin expansion is given below (start from the expansion of e^x and prove this identity):

$$\operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \left[x - \frac{x^3}{1! \times 3} + \frac{x^5}{2! \times 5} - \frac{x^7}{3! \times 7} + \dots \right] = \frac{2}{\sqrt{\pi}} \sum_{n=1}^\infty (-1)^{n+1} \frac{x^{2n-1}}{(n-1)!(2n-1)}$$




Figure 5.2 The graph of the error function $\operatorname{erf}(x)$ for $-2 \leq x \leq 2$

Example 5.13 | Example 1

Find the solution of the equation above for the following initial condition.

$$f(x) = \begin{cases} U_0 & |x| < 1 \\ 0 & |x| \geq 1 \end{cases}$$

We use the second form of the solution:

$$\begin{aligned} u(x, t) &= \frac{U_0}{2c\sqrt{\pi t}} \int_{-1}^1 e^{-\frac{(x-v)^2}{4c^2 t}} dv \stackrel{z = \frac{v-x}{2c\sqrt{t}}}{=} \frac{U_0}{\sqrt{\pi}} \int_{-\frac{1+x}{2c\sqrt{t}}}^{\frac{1-x}{2c\sqrt{t}}} e^{-z^2} dz \\ &= \frac{U_0}{\sqrt{\pi}} \left\{ \int_{-\frac{1+x}{2c\sqrt{t}}}^0 e^{-z^2} dz + \int_0^{\frac{1-x}{2c\sqrt{t}}} e^{-z^2} dz \right\} = \frac{U_0}{\sqrt{\pi}} \frac{\sqrt{\pi}}{2} \left\{ \frac{2}{\sqrt{\pi}} \int_0^{\frac{1+x}{2c\sqrt{t}}} e^{-z^2} dz + \frac{2}{\sqrt{\pi}} \int_0^{\frac{1-x}{2c\sqrt{t}}} e^{-z^2} dz \right\} \\ u(x, t) &= \frac{U_0}{2} \left[\operatorname{erf}\left(\frac{1+x}{2c\sqrt{t}}\right) + \operatorname{erf}\left(\frac{1-x}{2c\sqrt{t}}\right) \right] \end{aligned}$$




Figure 5.3 The temperature distribution $u(x, t)$ at $t = 0, \frac{1}{8}, \frac{1}{2}, 1, 2$ and 8 for the initial pulse $U_0 = 100$ and $c = 1$

 Example 5.14 | Example 2

Find the solution of the equation above for the following initial condition.

$$f(x) = \begin{cases} 1 & x > 0 \\ 0 & x \leq 0 \end{cases}$$

We use the second form of the solution:

$$\begin{aligned} u(x, t) &= \frac{1}{2c\sqrt{\pi t}} \int_0^\infty e^{-\frac{(x-v)^2}{4c^2t}} dv \quad z = \frac{v-x}{2c\sqrt{t}} \quad \frac{1}{\sqrt{\pi}} \int_{-\frac{x}{2c\sqrt{t}}}^\infty e^{-z^2} dz \\ &= \frac{1}{\sqrt{\pi}} \left\{ \int_{-\frac{x}{2c\sqrt{t}}}^0 e^{-z^2} dz + \int_0^\infty e^{-z^2} dz \right\} \\ u(x, t) &= \frac{1}{2} \left[\operatorname{erf}\left(\frac{x}{2c\sqrt{t}}\right) + 1 \right] \end{aligned}$$