


Integration of Complex Functions

انتگرال توابع مختلط



In this chapter

12.1	The Definite Integral and the Path of Integration	2
12.2	The Definite Integral of Complex Functions	4
12.3	Simple Closed Curves and Simply Connected Domains	6
12.4	Integration by Antiderivative	6
12.5	Integrals of Complex Functions Along Closed Paths	7
12.6	Cauchy's Integral Formula	10
12.7	Application: Evaluating Real Integrals	12
12.8	Cauchy's Theorem with Several Non-Analytic Points	16
12.9	Derivatives of an Analytic Function	17

12.1 The Definite Integral and the Path of Integration

The integral of complex functions, like that of real functions, appears in two forms:

- the indefinite integral, which amounts to finding an antiderivative;
- the definite integral, known as the integral along a path.

In the setting of complex functions the definite integral is also called the path integral or *line integral*, and is written

$$\int_C f(z) dz$$

C is called the path of integration.




Figure 12.1 A path of integration C from the point A to the point B in the z -plane

■ The Path of Integration

A path of integration in the complex plane consists of a starting point, an intermediate route and an end point. In many cases the best way to describe a path is to use an auxiliary variable which, as it increases, traverses the path from start to end in the desired direction.

$$z(t) = x(t) + iy(t), \quad a \leq t \leq b, \quad t, a, b \in \mathbb{R}$$

$$z(a) = A, \quad z(b) = B, \quad z, A, B \in \mathbb{C}$$




Figure 12.2 The path C , directed from $A = z(a)$ to $B = z(b)$

 Example 12.1

$$z(t) = t + i2t, \quad 1 \leq t \leq 2$$

This path is the line segment from the point $1 + 2i$ to the point $2 + 4i$.




Figure 12.3 The path $z(t) = t + i2t$, $1 \leq t \leq 2$

 Example 12.2

$$z(t) = \cos t + i \sin t, \quad 0 \leq t \leq 2\pi$$

$$z(t) = e^{it}, \quad 0 \leq t \leq 2\pi$$

This path is the unit circle, traversed counterclockwise.




Figure 12.4 The path $z(t) = e^{it}$, $0 \leq t \leq 2\pi$ (the unit circle)

■ The Derivative of the Path

$$\dot{z}(t) = \frac{d}{dt}z(t) = \lim_{\Delta t \rightarrow 0} \frac{z(t + \Delta t) - z(t)}{\Delta t} = \frac{d}{dt}x(t) + i \frac{d}{dt}y(t) = \dot{x}(t) + i\dot{y}(t)$$




Figure 12.5 The derivative of the path $\dot{z}(t)$ as the limit of $\frac{z(t+\Delta t)-z(t)}{\Delta t}$

12.2 The Definite Integral of Complex Functions

$$\int_C f(z) dz = \int_C (u + iv)(dx + i dy) = \int_C (u dx - v dy) + i \int_C (v dx + u dy)$$

$$\int_C f(z) dz = \int_t f(z(t)) \dot{z}(t) dt$$

Definition 12.3 Steps for computing a complex definite integral along a path

1. Describe the path parametrically: $z(t) = x(t) + iy(t)$, $a \leq t \leq b$.
2. Compute the derivative of $z(t)$ with respect to t : $\dot{z}(t) = \dot{x}(t) + i\dot{y}(t)$.
3. Substitute $z(t)$ into $f(z)$.
4. Integrate $f(z(t))\dot{z}(t)$ with respect to t for $a \leq t \leq b$.

Example 12.4

Compute the integral of the function $f(z) = \bar{z}$ along the path shown.

$$z(t) = t + it^2, \quad -1 \leq t \leq 1, \quad \dot{z}(t) = 1 + i2t$$

$$f(z) = \bar{z} = x - iy, \quad f(z(t)) = \bar{z}(t) = t - it^2$$

$$\begin{aligned} \int_{-1+i}^{1+i} \bar{z} dz &= \int_{t=-1}^1 (t - it^2)(1 + i2t) dt \\ &= \int_{t=-1}^1 [(t + 2t^3) + i(2t^2 - t^2)] dt = \int_{t=-1}^1 [(t + 2t^3) + it^2] dt \\ &= \left(\frac{1}{2}t^2 + \frac{2}{4}t^4 \right) + i \times \frac{1}{3}t^3 \Big|_{-1}^1 = 0 + i \frac{2}{3} \end{aligned}$$




Figure 12.6 The path $z(t) = t + it^2$, $-1 \leq t \leq 1$, from $-1 + i$ to $1 + i$

Example 12.5

Compute the integral of the function $f(z) = \bar{z}$ along the path shown.

$$z(t) = t + i, \quad -1 \leq t \leq 1, \quad \dot{z}(t) = 1$$

$$f(z) = \bar{z} = x - iy, \quad f(z(t)) = \bar{z}(t) = t - i$$

$$\int_{-1+i}^{1+i} \bar{z} dz = \int_{t=-1}^1 (t - i)(1) dt = \left(\frac{t^2}{2} - it \right) \Big|_{-1}^1 = 0 - i2$$

So for this complex function the value of the integral depends on the chosen path.




Figure 12.7 The path $z(t) = t + i$, $-1 \leq t \leq 1$, from $-1 + i$ to $1 + i$

Some Properties of the Definite Integral

$$\int_C [k_1 f_1(z) + k_2 f_2(z)] dz = k_1 \int_C f_1(z) dz + k_2 \int_C f_2(z) dz \quad (\text{linearity})$$

$$\int_{z_0}^Z f(z) dz = - \int_Z^{z_0} f(z) dz \quad (\text{reversing the direction of integration})$$

$$\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz \quad (\text{splitting the path of integration})$$




Figure 12.8 A path C from z_0 to Z split into two pieces C_1 and C_2

12.3 Simple Closed Curves and Simply Connected Domains

Definition 12.6 Simple closed curve

A simple closed curve is a closed curve that does not intersect itself.




Figure 12.9 Simple closed curves and closed curves that are not simple

Definition 12.7 Simply connected domain

A domain D is simply connected if, whenever a simple closed curve lies entirely in it, all the points inside the curve belong to this domain.




Figure 12.10 Simply connected and not simply connected domains

12.4 Integration by Antiderivative

Theorem 12.8

Let the complex function $f(z)$ be analytic in the simply connected domain D . If $F(z)$ is an antiderivative of $f(z)$, i.e. $F'(z) = f(z)$, then

$$\int_{z_0}^{z_1} f(z) dz = F(z_1) - F(z_0)$$

Example 12.9

$$\int_0^{1+i} z^2 dz = \frac{1}{3} z^3 \Big|_0^{1+i} = \frac{1}{3} (1+i)^3 = \frac{2}{3} (-1+i)$$

Example 12.10

$$\int_{-\pi i}^{\pi i} \cos z \, dz = \sin z \Big|_{-\pi i}^{\pi i} = 2 \sin(\pi i) = 2i \sinh \pi = 23.097 i$$

Example 12.11

$$\int_{8+\pi i}^{8-3\pi i} e^{z/2} \, dz = 2e^{z/2} \Big|_{8+\pi i}^{8-3\pi i} = 2(e^{4-3\pi i/2} - e^{4+\pi i/2}) = 0$$

Example 12.12

$$\int_{-i}^i \frac{dz}{z} = \ln i - \ln(-i) = \frac{i\pi}{2} - \left(-\frac{i\pi}{2}\right) = i\pi$$

12.5 Integrals of Complex Functions Along Closed Paths

The integral along a simple closed path C , written

$$\oint_C f(z) \, dz$$

is of great importance.

☺ Theorem 12.13 | Cauchy's integral theorem

If the function $f(z)$ is analytic in the simply connected domain D , then for every simple closed path C in this domain,

$$\oint_C f(z) \, dz = 0$$




Figure 12.11 A simple closed path C in the simply connected domain D

✓ **Proof.**

As we saw earlier, since the complex function is analytic, we have

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

$$\oint_C f(z) dz = \oint_C (u + iv)(dx + i dy) = \oint_C (u dx - v dy) + i \oint_C (v dx + u dy)$$

By Green's theorem (which you learned in Mathematics II),

$$\oint_C (L dx + M dy) = \iint_R \left(\frac{\partial M}{\partial x} - \frac{\partial L}{\partial y} \right) dx dy$$

Hence

$$\oint_C (u dx - v dy) + i \oint_C (v dx + u dy) = \iint_R \underbrace{\left(-\frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} \right)}_0 dx dy + i \iint_R \underbrace{\left(\frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right)}_0 dx dy = 0$$

where the first term vanishes because of the second Cauchy–Riemann condition and the second because of the first.

■

 Example 12.14

If the function is analytic in the whole complex plane, the integral along any closed path is zero:

$$\oint_C \cos z dz = 0, \quad \oint_C e^z dz = 0, \quad \oint_C z^n dz = 0, \quad n = 1, 2, 3, \dots$$

 Example 12.15

If the function has non-analytic points that lie neither on the path nor inside C , the integral is still zero:

$$\oint_{|z|=1} \sec z dz, \quad \oint_{|z|=1} \frac{dz}{z^2 + 4}$$

(the path is the unit circle.)

Example 12.16

If the function is not analytic, the integral along a closed path **may** be nonzero:

$$\oint_{|z|=1} \bar{z} dz, \quad z(t) = \cos t + i \sin t = e^{it}, \quad \dot{z}(t) = ie^{it}$$

$$\oint_{|z|=1} \bar{z} dz = \int_0^{2\pi} e^{-it} ie^{it} dt = 2\pi i$$

Example 12.17

If the function is not analytic on the path or inside it, the integral along a closed path **may** be zero:

$$\oint_C \frac{1}{z^2} dz = 0$$

Exercise 12.18

Prove that $\oint_C \frac{1}{z^2} dz = 0$.

Remark | Consequence 1

The connectedness of the domain is essential in Cauchy's integral theorem.

Remark | Consequence 2

If $f(z)$ is analytic in the simply connected domain D , the definite integral does not depend on the choice of path.

$$\int_{C_1} f(z) dz + \int_{C_2^*} f(z) dz = 0 \Rightarrow \int_{C_1} f(z) dz = - \int_{C_2^*} f(z) dz \Rightarrow \int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

Hence the integral from the starting point to the end point does not depend on the chosen path.




Figure 12.12 The paths C_1 , C_2 and C_2^* (C_2 with reversed direction) between z_1 and z_2

■ A Very Important Example

$$\oint_C \frac{dz}{z - z_0} = ? \quad C : z(t) = z_0 + re^{it}$$




Figure 12.13 A circle C with centre z_0 and radius r

$$z(t) = z_0 + re^{it}, \quad \dot{z}(t) = rie^{it}$$

$$\oint_C \frac{dz}{z - z_0} = \int_0^{2\pi} \frac{rie^{it}}{re^{it}} dt = \int_0^{2\pi} i dt = 2\pi i$$

★ Key result 12.19

$$\oint_C \frac{dz}{z - z_0} = 2\pi i$$

12.6 Cauchy's Integral Formula

☺ Theorem 12.20 | Cauchy's integral formula

If the function $f(z)$ is analytic in the simply connected domain D and z_0 is a point inside C , then

$$\frac{1}{2\pi i} \oint_C \frac{f(z)}{z - z_0} dz = f(z_0)$$

✓ **Proof.**

As we saw earlier, since the function $\frac{f(z)}{z - z_0}$ is analytic inside the connected domain Γ , we have

$$\oint_{\Gamma} \frac{f(z)}{z - z_0} dz = 0 = \int_{ACA'} \frac{f(z)}{z - z_0} dz + \int_{A'B'} \frac{f(z)}{z - z_0} dz + \int_{B'C'B} \frac{f(z)}{z - z_0} dz + \int_{BA} \frac{f(z)}{z - z_0} dz$$

$$\lim_{\substack{A \rightarrow A' \\ B \rightarrow B'}} \int_{BA} \frac{f(z)}{z - z_0} dz = - \int_{A'B'} \frac{f(z)}{z - z_0} dz$$

Consequently (the two integrals along the connecting segments cancel)

$$\lim_{\substack{A \rightarrow A' \\ B \rightarrow B'}} \int_{ACA'} \frac{f(z)}{z - z_0} dz + \int_{B'C'B} \frac{f(z)}{z - z_0} dz = 0 \Rightarrow \oint_C \frac{f(z)}{z - z_0} dz = \oint_{C'} \frac{f(z)}{z - z_0} dz$$

(here C' is traversed counterclockwise). Continuing:

$$\begin{aligned} \oint_C \frac{f(z)}{z - z_0} dz &= \oint_{C'} \frac{f(z)}{z - z_0} dz = \oint_{C'} \frac{f(z) - f(z_0) + f(z_0)}{z - z_0} dz \\ &= \oint_{C'} \frac{f(z) - f(z_0)}{z - z_0} dz + \oint_{C'} \frac{f(z_0)}{z - z_0} dz \\ &= \oint_{C'} g(z) dz + f(z_0) \oint_{C'} \frac{1}{z - z_0} dz = 0 + 2\pi i f(z_0) = 2\pi i f(z_0) \end{aligned}$$

where

$$g(z) \triangleq \begin{cases} \frac{f(z) - f(z_0)}{z - z_0} & z \neq z_0 \\ f'(z_0) & z = z_0 \end{cases}$$

is analytic inside C' . Therefore

$$\frac{1}{2\pi i} \oint_C \frac{f(z)}{z - z_0} dz = f(z_0)$$

■




Figure 12.14 The path Γ in the proof of Cauchy's integral formula: the curve C , the small circle C' centred at z_0 , and the two connecting segments AA' and BB'

12.7 Application: Evaluating Real Integrals

Example 12.21

Evaluate the integral $\int_{-\infty}^{\infty} \frac{dx}{1+x^2}$ using Cauchy's integral formula.

Consider the function $g(z) = \frac{1}{1+z^2}$. Using the path shown in the figure we find the integral along the closed path.

$$\oint_C \frac{dz}{1+z^2} = \oint_C \left[\frac{1}{z+i} \right] dz = 2\pi i \frac{1}{i+i} = \pi, \quad f(z) = \frac{1}{z+i}, \quad z_0 = i$$

On the other hand

$$\oint_C \frac{dz}{1+z^2} = \int_{-R}^R \frac{dx}{1+x^2} + \int_0^\pi \frac{iRe^{it} dt}{1+R^2e^{i2t}}$$

In the expression above, the first term as $R \rightarrow \infty$ is the required value.

$$\oint_C \frac{dz}{1+z^2} = \pi = \int_{-\infty}^{\infty} \frac{dx}{1+x^2} + I, \quad I = \lim_{R \rightarrow \infty} \int_0^\pi \frac{iRe^{it} dt}{1+R^2e^{i2t}} = 0$$

Therefore

$$\int_{-\infty}^{\infty} \frac{dx}{1+x^2} = \pi$$




Figure 12.15 The semicircular path in the upper half-plane; the poles are at $z = \pm i$

Example 12.22

Evaluate the integral $\int_{-\infty}^{\infty} \frac{\cos \omega x}{1+x^2} dx$ using Cauchy's integral formula ($\omega > 0$).

Consider the function $g(z) = \frac{e^{i\omega z}}{1+z^2}$. Using the path shown in the figure (the same semicircle as before) we find the integral along the closed path.

$$\oint_C \frac{e^{i\omega z}}{1+z^2} dz = \oint_C \frac{\left[\frac{e^{i\omega z}}{z+i} \right]}{z-i} dz = 2\pi i \frac{e^{i\omega i}}{i+i} = \pi e^{-\omega}, \quad f(z) = \frac{e^{i\omega z}}{z+i}, \quad z_0 = i$$

On the other hand

$$\oint_C \frac{e^{i\omega z} dz}{1+z^2} = \int_{-R}^R \frac{e^{i\omega x} dx}{1+x^2} + \int_0^\pi \frac{e^{i\omega Re^{it}} iRe^{it} dt}{1+R^2 e^{i2t}}$$

But:

$$I_1 = \lim_{R \rightarrow \infty} \int_0^\pi \frac{e^{i\omega Re^{it}} iRe^{it} dt}{1+R^2 e^{i2t}} \rightarrow 0$$

Proof:

$$e^{i\omega Re^{it}} = e^{i\omega R(\cos t + i \sin t)} = e^{i\omega R \cos t} e^{-\omega R \sin t}$$

$$\lim_{R \rightarrow \infty} |e^{i\omega Re^{it}}| = \lim_{R \rightarrow \infty} |e^{i\omega R \cos t}| \times |e^{-\omega R \sin t}| = 1 \times 0 = 0, \quad t \in [0, \pi]$$

On the other hand, in the integral I_1 the remaining factor of the integrand tends to zero as R tends to infinity. So the limit of I_1 as $R \rightarrow \infty$ is zero.

Therefore

$$\pi e^{-\omega} = \oint_C \frac{e^{i\omega z} dz}{1+z^2} = \int_{-R}^R \frac{e^{i\omega x} dx}{1+x^2} = \int_{-R}^R \frac{(\cos x + i \sin x) dx}{1+x^2} = \int_{-R}^R \frac{\cos x dx}{1+x^2} + i \int_{-R}^R \frac{\sin x dx}{1+x^2}$$

The second integral is zero. Why?

$$\int_{-\infty}^{\infty} \frac{\cos \omega x}{1+x^2} dx = \pi e^{-\omega}$$

Example 12.23

Evaluate the integral $I = \int_{-\infty}^{\infty} \frac{dx}{(x^2 + 1)(x^2 + 4)}$ using Cauchy's integral formula.

$$\frac{1}{(x^2 + 1)(x^2 + 4)} = \frac{A}{x^2 + 1} + \frac{B}{x^2 + 4} = \frac{\frac{1}{3}}{x^2 + 1} + \frac{-\frac{1}{3}}{x^2 + 4}$$

$$\int_{-\infty}^{\infty} \frac{dx}{(x^2 + 1)(x^2 + 4)} = \frac{1}{3} \int_{-\infty}^{\infty} \frac{dx}{x^2 + 1} - \frac{1}{3} \int_{-\infty}^{\infty} \frac{dx}{x^2 + 4}$$

First integral: $I_1 = \int_{-\infty}^{\infty} \frac{dx}{x^2 + 1}$ and $g_1(z) = \frac{1}{z^2 + 1}$:

$$\oint_C g_1(z) dz = \oint_C \frac{1}{z^2 + 1} dz = \oint_C \left[\frac{1}{z + i} \right] dz = 2\pi i \frac{1}{i + i} = \pi$$

$$\oint_C \frac{1}{z^2 + 1} dz = \pi = \int_{-R}^R \frac{dx}{1 + x^2} + \int_0^\pi \frac{iRe^{it} dt}{1 + R^2 e^{i2t}}$$

$$\lim_{R \rightarrow \infty} \oint_C \frac{1}{z^2 + 1} dz = \int_{-R}^R \frac{dx}{1 + x^2} + 0 = \pi \Rightarrow I_1 = \pi$$

Second integral: $I_2 = \int_{-\infty}^{\infty} \frac{dx}{x^2 + 4}$ and $g_2(z) = \frac{1}{z^2 + 4}$:

$$\oint_C g_2(z) dz = \oint_C \frac{1}{z^2 + 4} dz = \oint_C \left[\frac{1}{z + 2i} \right] dz = 2\pi i \frac{1}{2i + 2i} = \frac{\pi}{2}$$

$$\oint_C \frac{1}{z^2 + 4} dz = \frac{\pi}{2} = \int_{-R}^R \frac{dx}{x^2 + 4} + \int_0^\pi \frac{iRe^{it} dt}{R^2 e^{i2t} + 4}$$

$$\lim_{R \rightarrow \infty} \oint_C \frac{1}{z^2 + 4} dz = \int_{-R}^R \frac{dx}{x^2 + 4} + 0 = \frac{\pi}{2} \Rightarrow I_2 = \frac{\pi}{2}$$

As a result

$$I = \frac{1}{3} I_1 - \frac{1}{3} I_2 = \frac{1}{3} \pi - \frac{1}{3} \times \frac{\pi}{2} = \frac{\pi}{6}$$




Figure 12.16 The semicircular path; the poles are at $z = \pm i$ and $z = \pm 2i$

Example 12.24

Evaluate the integral $I = \int_0^{\infty} \frac{dx}{x^4 + 1}$ using Cauchy's integral formula.

$$g(z) = \frac{1}{z^4 + 1}, \quad z^4 + 1 = 0 \Rightarrow z_1 = e^{i\pi/4}, \quad z_2 = e^{i3\pi/4}, \quad z_3 = e^{i5\pi/4}, \quad z_4 = e^{i7\pi/4}$$

Only the two poles z_1 and z_2 lie inside the path C :

$$\begin{aligned} \oint_C g(z) dz &= \oint_C \frac{1}{z^4 + 1} dz = \oint_{C_1} \frac{1}{z^4 + 1} dz + \oint_{C_2} \frac{1}{z^4 + 1} dz \\ &= \oint_{C_1} \frac{1}{(z - z_1)(z - z_2)(z - z_3)(z - z_4)} dz + \oint_{C_2} \frac{1}{(z - z_1)(z - z_2)(z - z_3)(z - z_4)} dz \\ &= 2\pi i \frac{1}{(z - z_2)(z - z_3)(z - z_4)} \Big|_{z=z_1} + 2\pi i \frac{1}{(z - z_1)(z - z_3)(z - z_4)} \Big|_{z=z_2} \\ &= 2\pi i \frac{1}{\underbrace{(e^{i\pi/4} - e^{i3\pi/4})}_{\sqrt{2}} \underbrace{(e^{i\pi/4} - e^{i5\pi/4})}_{\sqrt{2}+i\sqrt{2}} \underbrace{(e^{i\pi/4} - e^{i7\pi/4})}_{i\sqrt{2}}} \\ &\quad + 2\pi i \frac{1}{\underbrace{(e^{i3\pi/4} - e^{i\pi/4})}_{-\sqrt{2}} \underbrace{(e^{i3\pi/4} - e^{i5\pi/4})}_{i\sqrt{2}} \underbrace{(e^{i3\pi/4} - e^{i7\pi/4})}_{-\sqrt{2}+i\sqrt{2}}} = \frac{\pi}{\sqrt{2}} \end{aligned}$$

On the other hand

$$\begin{aligned} \oint_C \frac{1}{z^4 + 1} dz &= \frac{\pi}{\sqrt{2}} = \int_{-R}^R \frac{dx}{x^4 + 1} + \int_0^{\pi} \frac{iRe^{it} dt}{R^4 e^{i4t} + 1} \\ \lim_{R \rightarrow \infty} \oint_C \frac{1}{z^4 + 1} dz &= \int_{-\infty}^{\infty} \frac{dx}{x^4 + 1} + \lim_{R \rightarrow \infty} \int_0^{\pi} \frac{iRe^{it} dt}{R^4 e^{i4t} + 1} = \frac{\pi}{\sqrt{2}} \\ \int_{-\infty}^{\infty} \frac{dx}{x^4 + 1} &= \frac{\pi}{\sqrt{2}} \Rightarrow \int_0^{\infty} \frac{dx}{x^4 + 1} = \frac{\pi}{2\sqrt{2}} \end{aligned}$$




Figure 12.17 The semicircular path and the circles C_1 and C_2 around the poles z_1 and z_2 ; the poles z_3 and z_4 lie in the lower half-plane

12.8 Cauchy's Theorem with Several Non-Analytic Points

☹️ Theorem 12.25

If the function $f(z)$ is analytic at all points of the domain D except the points $z_1, z_2, z_3, \dots, z_n$ inside C , then

$$\oint_C f(z) dz = \oint_{C_1} f(z) dz + \oint_{C_2} f(z) dz + \dots + \oint_{C_n} f(z) dz$$

where C_k is a circle centred at z_k that lies inside C and does not contain any of the other points $z_1, \dots, z_n$.




Figure 12.18 A path C and the non-analytic points $z_1, \dots, z_n$ inside it

✓ **Proof.**

Build the path Γ from C (counterclockwise), the small circles $C_1, \dots, C_n$ (clockwise) and the segments connecting them to C ; the integral along each connecting segment is traversed twice in opposite directions and cancels. The function is analytic inside Γ , so

$$\oint_{\Gamma} f(z) dz = 0 \Rightarrow \oint_C f(z) dz + \oint_{C_1^{\circlearrowright}} f(z) dz + \dots + \oint_{C_n^{\circlearrowright}} f(z) dz = 0$$

$$\oint_C f(z) dz = -\oint_{C_1^{\circlearrowright}} f(z) dz - \dots - \oint_{C_n^{\circlearrowright}} f(z) dz = \oint_{C_1^{\circlearrowleft}} f(z) dz + \dots + \oint_{C_n^{\circlearrowleft}} f(z) dz$$

where $C_k^{\circlearrowright}$ is the circle C_k traversed clockwise and $C_k^{\circlearrowleft}$ the same circle traversed counterclockwise. ■




Figure 12.19 The path Γ for the proof: C , the circles $C_1, \dots, C_n$ and the connecting segments

12.9 Derivatives of an Analytic Function

☺ **Theorem 12.26** | Cauchy's integral formula for the derivative

If the function $f(z)$ is analytic in the simply connected domain D and z_0 is a point inside C , then

$$\frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^2} dz = f'(z_0)$$

✓ **Proof.**

$$\begin{aligned}
f'(z_0) &= \lim_{\Delta t \rightarrow 0} \frac{f(z_0 + \Delta t) - f(z_0)}{\Delta t}, \quad f(z_0) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{z - z_0} dz, \quad f(z_0 + \Delta t) \\
&= \frac{1}{2\pi i} \oint_C \frac{f(z)}{z - z_0 - \Delta t} dz \\
\frac{f(z_0 + \Delta t) - f(z_0)}{\Delta t} &= \frac{1}{2\pi i \Delta t} \left\{ \oint_C \frac{f(z)}{z - z_0 - \Delta t} dz - \oint_C \frac{f(z)}{z - z_0} dz \right\} \\
&= \frac{1}{2\pi i \Delta t} \oint_C f(z) \left[\frac{1}{z - z_0 - \Delta t} - \frac{1}{z - z_0} \right] dz \\
&= \frac{1}{2\pi i \Delta t} \oint_C f(z) \left[\frac{\Delta t}{(z - z_0 - \Delta t)(z - z_0)} \right] dz \\
&= \frac{1}{2\pi i} \oint_C f(z) \left[\frac{1}{(z - z_0 - \Delta t)(z - z_0)} \right] dz \\
f'(z_0) &= \lim_{\Delta t \rightarrow 0} \frac{f(z_0 + \Delta t) - f(z_0)}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{1}{2\pi i} \oint_C f(z) \left[\frac{1}{(z - z_0 - \Delta t)(z - z_0)} \right] dz \\
&= \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^2} dz
\end{aligned}$$

■

☹ **Theorem 12.27** | Generalization of Cauchy's integral formula

If the function $f(z)$ is analytic in the simply connected domain D and z_0 is a point inside C , then

$$\frac{n!}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^{n+1}} dz = f^{(n)}(z_0)$$

✍ **Exercise 12.28**

Prove it by mathematical induction.

📖 **Example 12.29**

Evaluate the following integral. (C is the unit circle.)

$$\begin{aligned}
I &= \oint_C \frac{z^2}{(2z - 1)^2} dz \\
I &= \oint_C \frac{z^2}{4\left(z - \frac{1}{2}\right)^2} dz = \frac{1}{4} \oint_C \frac{z^2}{\left(z - \frac{1}{2}\right)^2} dz = \frac{1}{4} \times 2\pi i \times \frac{d}{dz}(z^2) \Big|_{z=1/2} = \frac{1}{4} \times 2\pi i \times (2z) \Big|_{z=1/2} \\
&= \frac{\pi}{2} i
\end{aligned}$$

($z = \frac{1}{2}$ lies inside the unit circle.)

Example 12.30

Evaluate the following integral. (C is the unit circle.)

$$I = \oint_C \frac{e^{-z} \sin z}{z^2} dz$$

$$f(z) = e^{-z} \sin z, \quad z_0 = 0, \quad f'(z) = -e^{-z} \sin z + e^{-z} \cos z, \quad f'(z_0) = f'(0) = 1$$

$$I = 2\pi i f'(z_0) = 2\pi i$$

Exercise 12.31 | Problem

If $f(z)$ is analytic in the whole z -plane and bounded in the whole plane, i.e. $|f(z)| < M$, prove that $f(z) = k$, $k \in \mathbb{C}$.

Choose the path C to be a circle of radius R centred at z_0 , and let R tend to infinity.

$$f'(z_0) = \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^2} dz$$

$$\begin{aligned} |f'(z_0)| &= \left| \frac{1}{2\pi i} \oint_C \frac{f(z)}{(z - z_0)^2} dz \right| \leq \frac{1}{2\pi} \oint_C \frac{|f(z)|}{|z - z_0|^2} |dz| \leq \frac{1}{2\pi} \frac{M}{R^2} \oint_C |dz| = \frac{1}{2\pi} \frac{M}{R^2} \cdot 2\pi R \\ &= \frac{M}{R} \rightarrow 0 \quad (R \rightarrow \infty) \end{aligned}$$

$$\Rightarrow |f'(z_0)| = 0 \Rightarrow f'(z) = 0 \Rightarrow f(z) = k$$




Figure 12.20 A circle C with centre z_0 and radius R

 Exercise 12.32

Evaluate the following integrals:

1. $\oint_C \frac{z^2}{(2z-1)^4} dz$, C : the unit circle
2. $\int_{1+i}^{1+3i} e^z \sin z \, dz$
3. $\int_{1+i}^{2+2i} \operatorname{Re}\{z^2\} \, dz$ along a straight-line path
4. $\oint_C \frac{e^{\cos z} \sin z}{1+z^6} dz$ along the path below




Figure 12.21 The path of exercise 4: a circle with centre $3 + 3i$ and radius $r = 1$