


# Evaluating Definite Integrals by Residues

محاسبه انتگرال‌های معین به روش مانده‌ها



## In this chapter

|             |                                     |   |
|-------------|-------------------------------------|---|
| <b>14.1</b> | Integrals of the First Class .....  | 3 |
| <b>14.2</b> | Integrals of the Second Class ..... | 4 |

 **Theorem 14.1** | The residue theorem

If the function  $f(z)$  is analytic everywhere inside the path  $C$  except at the points  $z_1, z_2, \dots, z_n$ , and the residues of the function at these points are  $A_1^1, A_1^2, \dots, A_1^n$  respectively, then

$$\oint_C f(z) dz = 2\pi i \sum_{k=1}^n A_1^k$$

Many different kinds of integrals can be solved by the method of residues. Two classes of them are mentioned here:

 **Definition 14.2** | Integrals of the first class

In these integrals the denominator of the integrand is a polynomial, and the numerator is a polynomial, an exponential, or a sine or cosine function (which can also be expressed exponentially).

 **Definition 14.3** | Integrals of the second class

In these integrals the integrand is a function of sine or cosine.

### 14.1 Integrals of the First Class

#### Example 14.4

Evaluate the integral  $\int_0^\infty \frac{\cos \omega x}{(1+x^2)^2} dx$  ( $\omega > 0$ ).

Consider the function  $f(z) = \frac{e^{i\omega z}}{(1+z^2)^2}$ . It is not analytic at  $z = i$  and  $z = -i$ . The path  $C$  is the upper semicircle of radius  $R$ .


$$\begin{aligned} \oint_C \frac{e^{i\omega z}}{(1+z^2)^2} dz &= 2\pi i \times \text{Res} \left\{ \frac{e^{i\omega z}}{(1+z^2)^2} \right\} \Big|_{z=i} \\ &= 2\pi i \times \left[ \frac{d}{dz} \left\{ (z-i)^2 \frac{e^{i\omega z}}{(z-i)^2(z+i)^2} \right\} \right]_{z=i} \\ &= 2\pi i \times \left[ \frac{i\omega e^{i\omega z}(z+i) - 2e^{i\omega z}}{(z+i)^3} \right]_{z=i} \\ &= 2\pi i \times e^{-\omega} \left( \frac{i\omega(2i) - 2}{(2i)^3} \right) \\ &= 2\pi i \times e^{-\omega} \frac{\omega + 1}{4i} = \frac{\pi}{2} (\omega + 1) e^{-\omega} \end{aligned}$$

On the other hand

$$\oint_C \frac{e^{i\omega z}}{(1+z^2)^2} dz = \int_{-R}^R \frac{e^{i\omega x}}{(1+x^2)^2} dx + \int_0^\pi \frac{e^{i\omega R e^{it}} R i e^{it}}{(1+R^2 e^{i2t})^2} dt$$

and as  $R \rightarrow \infty$  the integral along the semicircle vanishes, while the imaginary part of the integral along the real axis (the odd function  $\frac{\sin \omega x}{(1+x^2)^2}$ ) is zero:

$$\begin{aligned} \oint_C \frac{e^{i\omega z}}{(1+z^2)^2} dz &\xrightarrow{R \rightarrow \infty} 2 \int_0^\infty \frac{\cos \omega x}{(1+x^2)^2} dx \\ \int_0^\infty \frac{\cos \omega x}{(1+x^2)^2} dx &= \frac{\pi}{4} (\omega + 1) e^{-\omega} \end{aligned}$$



**Figure 14.1** The upper semicircular path; the double pole  $z = i$  lies inside the path and  $z = -i$  outside it

## 14.2 Integrals of the Second Class

$$\int_0^{2\pi} f(a \cos t, b \sin t) dt$$

To solve such integrals we put  $z = e^{it}$ . Then


$$dz = ie^{it} dt \Rightarrow dt = \frac{dz}{ie^{it}} = \frac{dz}{iz}$$

$$\sin t = \frac{e^{it} - e^{-it}}{2i} = \frac{z - z^{-1}}{2i}, \quad \cos t = \frac{e^{it} + e^{-it}}{2} = \frac{z + z^{-1}}{2}$$

### ★ Key result 14.5

$$\int_0^{2\pi} f(a \cos t, b \sin t) dt = \oint_C f\left(a \frac{z + z^{-1}}{2}, b \frac{z - z^{-1}}{2i}\right) \frac{dz}{iz}$$

Here  $C$  is the unit circle, and the integral is easily solved with the method of residues.



**Figure 14.2** The unit circle  $z = e^{it}$ ,  $0 \leq t \leq 2\pi$

### Example 14.6

Evaluate the integral  $\int_0^{2\pi} \frac{1}{2 + \cos x} dx$ .

$$z = e^{ix}, \quad dz = ie^{ix} dx, \quad dx = \frac{dz}{iz}, \quad \cos x = \frac{z + z^{-1}}{2} = \frac{z^2 + 1}{2z}$$


$$2 + \cos x = 2 + \frac{z^2 + 1}{2z} = \frac{z^2 + 4z + 1}{2z}$$

$$\begin{aligned} \int_0^{2\pi} \frac{1}{2 + \cos x} dx &= \oint_C \frac{2z}{z^2 + 4z + 1} \frac{dz}{iz} = \frac{2}{i} \oint_C \frac{1}{z^2 + 4z + 1} dz \\ &= \frac{2}{i} \times 2\pi i \times \sum_k \operatorname{Res} \left\{ \frac{1}{z^2 + 4z + 1} \right\} \Big|_{z=z_k} \end{aligned}$$

$$z^2 + 4z + 1 = 0 \Rightarrow z_1 = -2 + \sqrt{3} \approx -0.27, \quad z_2 = -2 - \sqrt{3} \approx -3.73$$

Only  $z_1$  lies inside the unit circle.

$$\begin{aligned} \int_0^{2\pi} \frac{1}{2 + \cos x} dx &= 4\pi \times \operatorname{Res} \left\{ \frac{1}{z^2 + 4z + 1} \right\} \Big|_{z=z_1} \\ &= 4\pi \left[ (z + 2 - \sqrt{3}) \frac{1}{(z + 2 - \sqrt{3})(z + 2 + \sqrt{3})} \right]_{z_1 = -2 + \sqrt{3}} = \frac{2\pi}{\sqrt{3}} \end{aligned}$$



**Figure 14.3** The unit circle and the poles  $z_1 \approx -0.27$  (inside) and  $z_2 \approx -3.73$  (outside)

### Exercise 14.7 | Applying the residue theorem to integrals

Using the residue theorem, evaluate the following definite integrals:

$$I_1 = \int_0^{2\pi} \frac{dt}{1 - 2p \cos t + p^2}, \quad I_2 = \int_{-\infty}^{\infty} \frac{\sin 2x}{x^2 + x + 1} dx$$

$$I_3 = \int_{-\infty}^{\infty} \frac{x}{(x^2 - 2x + 2)^2} dx, \quad I_4 = \int_{-\infty}^{\infty} \frac{\cos^4 x}{(x^2 + 1)(x^2 + 4)} dx$$