

# Benefits of Bayesian Modelling Approaches For Conservation Science Under Climate Change

A.K. Ettinger<sup>1,a</sup>, H. Eyster<sup>2</sup>, D. Loughnan<sup>3</sup>, X. Wang<sup>3</sup>, E.M. Wolkovich<sup>3</sup>, M. Auger-Methe<sup>3</sup>, R. Zenil-Ferguson<sup>4</sup>, V. Leos Barajas<sup>5</sup>, Leithen M’Gonigle<sup>6</sup>, Hanna Jackson<sup>6</sup>, and Others from Symposium working group?<sup>7</sup>

<sup>1</sup>The Nature Conservancy, Seattle, Washington, USA

<sup>2</sup>TNC

<sup>3</sup>UBC

<sup>4</sup>UKY

<sup>5</sup>University of Toronto

<sup>6</sup>SFU

<sup>7</sup>Multiple other institutions?

<sup>a</sup>Corresponding author; email: [ailene.ettinger@tnc.org](mailto:ailene.ettinger@tnc.org); mailing address: 74 Wall Street. Seattle, WA 98121, USA

October 17, 2024

**Author contributions:** All authors conceived of this manuscript, which began through conversations at a Bayesian Generative Modelling Symposium at University of British Columbia in 2024; AKE led the writing of the manuscript, all authors contributed portions of text and revised the manuscript.

**Data Accessibility**

**Running title**

**Key words**

**Paper type** *Perspective* in Conservation Letters (3,000 words). Back-up/next journal to submit to is Conservation Science and Practice

**Focal Audience** Conservation biologists and other scientists who contribute to or would like to influence conservation and environmental policy (e.g., IPCC, IUCN)

**Abstract**

# Introduction

Conservation science in the 21st century seeks to address the dual crises of climate change and rapid biodiversity loss. These are problems that require urgent action globally, as loss of earth’s biodiversity and its benefits are accelerating (Brondizio et al., 2019; Ripple et al., 2017; Tittensor et al., 2014). Climate change has added complexity to conservation biology’s traditional goal and approaches for conserving biodiversity. Climate change both affects and is affected by biodiversity conservation (e.g., nature-based climate solutions or ‘natural climate solutions’). Thus, integrating approaches to bolster adaptation and resilience to climate change, as well as greenhouse gas mitigation, are critical components of planning and implementing conservation in the era of climate change.

The need for robust and usable conservation science under climate change is necessary at scales ranging from local to global. For example, preserve managers seek guidance about how to best steward natural resources for climate resilience (Nadeau et al., 2015), international climate policy relies on scientific data and publications for systematic observation of climate systems and impacts to people. IPCC (2007). Developing the evidence base for urgent climate and biodiversity questions often requires synthesizing multiple data sources and incomplete datasets, given the complex ecological systems in which many conservation science problems are grounded. Thus, a critical part of building the evidence base is ensuring reproducibility and transparency, including clear communication of uncertainty (Ellis et al., 2024; IPCC, 2007).

Bayesian data analysis provides a framework and approaches that support these needs of conservation science in the 21st century. Bayesian approaches facilitate synthesis of multiple sources of data to update probabilities of focal outcomes of interest after examining new data (e.g., priors, see Box 1). Bayesian methods are well-suited to decision making, as they moving beyond strict null-hypothesis testing to provide a quantitative measure of the probability of a hypothesis being true given the available data. Some fields within conservation biology and natural resource management have adopted Bayesian methods (e.g., wildlife mark and recapture models or occupancy models (Kery and Schaub, 2011), fisheries(Doll and Jacquemin, 2018)), but historically they have not been widely used in conservation biology.

## Aim

We aim to highlight features of Bayesian approaches that are well-suited to conservation science to practitioners who may be unfamiliar with Bayesian methods, and hope to accelerate more widespread adoption of Bayesian data analytical approaches in conservation. We believe that, with more widespread adoption, these approaches could enhance progress of conservation science. We first review some benefits of using Bayesian methods to address conservation science questions, focusing on three key features: the iterative workflow process, flexible modelling frameworks, and integration of uncertainty. We then show that Bayesian approaches have been steadily increasing in ecology and natural resource management, highlighting that the time is right for more widespread use and integration into conservation science, policy, and practice. Finally, we share resources that we hope will make Bayesian methods more approachable to those who have not used them before, including example code and analyses through three case studies relevant to current conservation problems, and a list of resources to get started with these methods.

# Benefits of Bayesian Modelling for Conservation Science

## Iterative workflows that align with adaptive management practices in conservation

Conservation scientists often need to develop new alternative approaches that differ from current ‘business-as-usual’ approaches. The need to test new approaches, coupled with the fact that ecosystems are dynamic and often yield unexpected behaviors (Levin et al., 2012; Gross, 2013), have led to practices of adaptive management in conservation (Holling and Walters, 1978) (Fig. 1). Cycles of adaptive management include stages of assessing threats and conservation goals, planning strategies, implementation, and monitoring, implementing those plans, analyzing results of initial implementation, adapting implementation based on these analyses, and sharing learning from the process. This cycle of iteration continues until the end of the project, ideally when conservation goals are met, and the intention is for this evidence based approach to lead to improved conservation outcomes (<https://www.conservationbydesign.org/>).

Bayesian modelling workflows are similarly iterative and could be integrated into the cyclical process of adaptive conservation science and management. In our experience, conservation processes often postpone data analysis until late in the cycle; this is visually represented in the Conservation by Design diagram shown in (Fig. 1). We believe that integrating the modelling workflow throughout the cycle would lead to valuable insights earlier in process, result in more robust conservation science and, perhaps, then better outcomes after implementation. Model development could occur in the assessing stage, and tested with simulated data in the planning stage. Data collection occurs as part of monitoring during implementing stage and then models are run with empirical data as part of the analyzing stage. Model checking and revising occurs during adapting stages and then sharing learning about modelling as well as conservation implementation occurs in parallel.

## Flexible frameworks for accomodating complex data

Conservation problems are complicated and addressing them, especially in the era of climate change, requires integrating social, economic, biological, and physical information to provide the evidence base upon which decisions can be made. Focal populations are often small, and ecological and social data are often non-normal, ‘unbalanced’ and nested within hierarchical or non-hierarchical groupings that are non-independent (e.g., species or other groupings related to evolutionary history/genetics, spatial clustering such as plots or sites, or temporal clustering such as day or year); these qualities challenges many traditional, commonly used statistical approaches (e.g., analysis of variance). Further, conservation evidence comes in many forms, including from quantitative studies, community knowledge, expert knowledge, traditional ecological knowledge, and others. Conservation decision-making requires integrating these multiple sources of information to provide an evidence base upon which decisions can be grounded (Stern and Humphries, 2022).

Bayesian modelling approaches are well-suited to address these constraints, given their flexibility and power to provide robust estimates under a wide range of conditions (e.g., Case Study 1). Bayesian models can be specified to include a wide range of data distributions via ready-to-use packages in programs like R (e.g.), and expanded to infinite model specifications when the user codes them by hand. Multiple sources and types of data can be amalgamated into Bayesian Belief Networks (Marcot et al., 2001; Newton et al., 2007), and extant information can be used to inform ‘prior distributions’ used in Bayesian modelling (O’Leary, 2008; McNeish, 2016). (For more on what ‘prior distributions’ or ‘priors’ are, see Box 1.) Bayesian methods are also well-suited to accommodate the small population sizes that are often a focus of conservation because they are at high risk of extinction (Stinchcombe et al., 2002). This is because Bayesian methods do not rely on asymptotic behavior (as frequentist statistics due (McNeish, 2016)), so are better able to accommodate small sample sizes.

The flexibility of Bayesian methods is also beneficial and well-suited to conservation planning because they move beyond ‘null hypothesis’ frameworks and a focus on p-values to facilitate comparison of support for a variety of hypotheses or interventions (van Zyl, 2018). Null hypothesis significance testing and conventions of rejecting the null hypothesis when  $p < 0.05$  have long been dominant in conservation biology, as in ecology and related biological fields (e.g., ecotoxicology Erickson and Rattner, 2020). P-value-focused conventions are becoming less prevalent, but many biologists are unsure of alternative ways to analyse and interpret their data (Halsey, 2019). Bayesian approaches and workflows offer an alternative framework, facilitating, for example, assessments of which interventions will most likely bring about conservation gains (Prato, 2005), whether the current versus alternative management practices produce similar results (Gallistel, 2009), and whether a population has passed a particular threshold, such as declining ‘rapidly’ versus ‘moderately’, as defined by a particular probability (Brooks et al., 2008). The flexibility and power of Bayesian modelling require training to implement, as well as thoughtful specification and careful interpretation. We believe this is not unique to Bayesian modelling; rather it is applicable to all statistical tools and approaches.

## **Integrated approaches for quantifying and propagating uncertainty**

Bayesian analyses are particularly useful for decision-making because they are adept at integrating not only a range of information types, but also the uncertainty associated with these information types (e.g., Stern and Humphries, 2022). The integration of multiple datasets required by many conservation problems in turn necessitates quantifying and sometimes propagating uncertainty across multiple sources and/or multiple modeling steps (see Case Study 2). Bayesian approaches enable straightforward quantification and propagation of uncertainty, including for some conservation problems that can require analyses for which frequentist statistics are unable to compute the associated uncertainties (Bolker et al., 2009; Bates, 2006), and for which the intuitive interpretation matches the technical definition, yielding interpretable results, particularly for non-statistician colleagues and decision-makers (Fornacon-Wood et al., 2021).

Moreover, Bayesian methods enable uncertainty to be propagated through multiple analyses, ensuring that end results represent the full uncertainty of the process under study. (Draper, 1995; Gilbert et al., 2023; Eyster et al., 2022; Saunders et al., 2019). For example, using Bayesian methods, one can calculate the abundance of birds in different types of management landscapes such as traditional agriculture and perennial polycultures, and then propagate the uncertainty associated with those abundances into a downstream analysis to test whether the bird communities in the alternative perennial polyculture landscape are maintained simply as an ecological sink (Eyster et al., 2022).

## **Increasing use of Bayesian approaches**

Ecology and conservation science have become more quantitative and data-driven fields in recent decades, and use of Bayesian approaches appears to be increasing (?). The rise in Bayesian approaches likely reflects an acknowledgement of their utility, given the benefits they offer, including those we mention here.

Within the general increase in use of Bayesian methods, however, there appears to be notable variation by subfield, with more widespread use in fisheries and wildlife biology, and less in forestry and plant ecology (Fig. 2). This may reflect adoption of particular model types within subfields, such as mark-and-capture studies within wildlife biology (e.g. REFs), or high-demand needs within natural resource management that can be met by Bayesian modelling, such as fisheries (e.g., REF). We see opportunities for more widespread use the fields that do not appear to have taken up Bayesian approaches as rapidly. For example, within plant ecology, incorporation of Bayesian models may allow better integration of observation and process models, similar to what has been used in wildlife biology. For analysis of long-term data this may allow for better separation of trends due to var

Though Bayesian methods are increasing, they are not yet standard, widely used approaches that are well-integrated into global systems of conservation and climate science, policy and practice (e.g., IPCC, IUCN). For example, IPCC-described methods for uncertainty propagation do not include Bayesian approaches (IPCC, 2007), though these approaches would offer straightforward implementation (e.g., Case Study 2).

## **A future with more widespread use of Bayesian modelling in Conservation**

The urgency of conservation problems and complexity of ecological systems requires flexible, powerful modelling approaches. We envision a future with more widespread adoption of Bayesian modelling approaches in conservation training and implementation. For example, conservation and ecology students, across undergraduate and graduate levels, should receive analytical training to provide strong foundations in Bayesian statistics. Many introductory statistics classes today still focus on t-tests and linear regressions, which are often not appropriate for ecological datasets. It doesn't have to be this way! Implementing Bayesian modelling is easier than ever before, with increased access to computational resources and training (see, e.g., Resources to Get Started). Further, we believe that the IPCC and other global institutions should include guidelines for Bayesian approaches, given their increasing use (Fig. 2) and as biological feedbacks get more integrated into the climate/biophysical analyses that dominated early IPCC work. These flexible and powerful approaches can help us develop more robust understanding of complex ecological systems, and facilitate fuller use of the data we have to address urgent conservation problems under climate change.

## **Case Studies**

### **Case Study 1: Robust estimates of trends in species or populations of concern**

In some cases, Bayesian approaches can lead to importantly different conclusions than common Fisherian approaches, such as null hypothesis testing (citep:wade2000bayesian). Consider, for example, sampling five populations of a species across its range—from north to south—to monitor for changes in the population size with climate change. After collecting data for ten years, a traditional null hypothesis testing approach to analyzing the data, using the common Type I error value ( $\alpha$ ) of 0.05 would find changes in only two of the populations—the furthest north population, which appears to be increasing, and the second furthest north population, also increasing. The three other populations are not significantly changing under this approach (Fig. 3 left). In contrast, a Bayesian approach (using weakly informative priors centered at zero, all code provided in supplement) would likely focus on the posteriors, where small differences in the variance do not appear so different (Fig. 3 right). Here, a clear trend emerges where trends in population correlate with position in range—with the most northern population increasing the most and the most southern population decreasing the most. This pattern across the range is the type predicted by climate change and may be missed with a classic Fisherian approach (combining null hypothesis testing with threshold values for 'significance'), leading to potential very different conservation and management decisions.

### **Case Study 2: Incorporating and propagating uncertainty**

One major approach that integrates biodiversity conservation and climate change mitigation is 'natural climate solutions' (NCS) (Ellis et al., 2024), also called nature-based climate solutions. NCS are intentional human actions (or 'NCS pathways') that protect, restore, and improve management of forests, wetlands, grasslands, oceans, and agricultural lands to mitigate climate change (Griscom et al., 2017). ADD a bit more

detail here.

We simulate flux data from a field study quantifying greenhouse gas fluxes (methane, carbon dioxide) s in peatlands with and without grazing in Ecuador (Sanchez et al) to show how Bayesian approaches can be used to propoagate uncertainty in a straightforward way.

- carbon Mitigation= flux X extent, Fig. 4
- uncertainty propogation using posterior
- see <https://github.com/AileneKane/bayes4cons/tree/main/analyses/ncs.R>

### Case Study 3: State-space model and priors example

We demonstratte use of priors via a population model, described in Auger-Methe et al 2021 and based on Jamieson and Brooks (2004) and Dennis and Taper (1994). This is a simple population model with density dependence, made up of process and observation components, as described at <https://github.com/AileneKane/bayes4cons/blob/>

In this scenario, we have re-introduced 10 adult females and some 10 adult males from an extirpated species in 2003 in a conservation area, and we are monitoring their growth for the past 20 years to see if they have reached carrying capacity and what is that carrying capacity.

This is a species that has a long life span (e.g. 20+ years) and creates long term pairs that can produce maximum 2 offspring when in good conditions, and therefore could be able to almost double in size ever year. Here the conservation area does not allow the animal to fulfill it's full growth ( $\beta_0 = 1.3$ ). From a repeated assessments of the efficiency of the line transect in early years, when all of the re-introduce animals were individually marked, we know that we can miss many individuals and have estimated the  $\sigma_o = 10$ .

As for most population of this species the birth rate, which is the main source of biological stochasticity, vary by less than 5

When we fit the model with vague priors, there are many warnings, indicating the model is problematic. However, when we use knowledge gained from previous studies of species' biology to inform priors, we are able to fit models.

## Box 1: Defining Bayesian Analysis (i.e., Inference and Workflow)

Bayesian methods allow us to combine prior knowledge of a system with available data to quantitatively measure the probability of a model or hypothesis being true. A Bayesian model treats parameters as random variables for which a known probability distribution is defined based on prior knowledge of the parameter uncertainty. This allows the development of bespoke and uniquely formulated models for any given data-generating process.

At the foundation of this statistical approach is the use of Bayes' Theorem (eqn 1). Consisting of three components, this theorem combines a prior distribution ( $A$ ) as defined using data from previous experiments, expert opinions, or literature with a likelihood function ( $f(B|A)$ ) to estimate probability distributions for each parameter ( $P(A|B)$ ).

$$P(A|B) = \frac{f(B|A)(A)}{P(B)} \quad (1)$$

Since the results of Bayesian methods are probabilistic, they allow us to make stronger inferences based on a model's predicted uncertainty. As new data is collected or inferences made, we can improve our model priors and iteratively improve our understanding of the data-generating processes that underlie ecological systems. Model estimates can be further used to perform simulations and more robustly forecast changes under future conditions.

## Box 2: Resources to Get Started

### Priors

Default priors. Chosen without critical thought or evaluation. Fear of being too subjective. Defending prior choice promotes good statistical inference

Resources on priors:

- Banner et al. 2020 <https://besjournals.onlinelibrary.wiley.com/doi/full/10.1111/2041-210X.13407>

Other resources:

- A guide to Bayesian model checking for ecologists (Conn et al., 2018)
- Gentle Introduction to Bayesian statistics (Van de Schoot et al., 2014) <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4611111/>
- Bayesian model selection for ecologists. (Hooten and Hobbs, 2015) <https://doi.org/10.1890/14-0661.1>
- Bayesian Inference for ecologists. (?)
- Why becoming Bayesian? (Clark, 2005) <https://doi.org/10.1111/j.1461-0248.2004.00702.x>
- Bayesian Workflow (?)

## References

- Bates, D. 2006. R| lmer, p-values and all that. R-help mailing list .
- Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and evolution. *Trends in Ecology & Evolution* 24:127–135. PT: J; UT: WOS:000264615200003.
- Brondizio, E., S. M. Díaz, J. Settele, H. Ngo, M. Gueze, Y. Aumeeruddy-Thomas, X. Bai, A. Geschke, Z. Molnár, A. Niamir, et al. 2019. Assessing a planet in transformation: Rationale and approach of the ipbes global assessment on biodiversity and ecosystem services .
- Brooks, S. P., S. N. Freeman, J. J. Greenwood, R. King, and C. Mazzetta. 2008. Quantifying conservation concern—bayesian statistics, birds and the red lists. *Biological Conservation* 141:1436–1441.
- Clark, J. S. 2005. Why environmental scientists are becoming bayesians. *Ecology letters* 8:2–14.
- Conn, P. B., D. S. Johnson, P. J. Williams, S. R. Melin, and M. B. Hooten. 2018. A guide to bayesian model checking for ecologists. *Ecological Monographs* 88:526–542.
- Doll, J. C., and S. J. Jacquemin. 2018. Introduction to bayesian modeling and inference for fisheries scientists. *Fisheries* 43:152–161.

- Draper, D. 1995. Assessment and propagation of model uncertainty. *Journal of the Royal Statistical Society Series B: Statistical Methodology* 57:45–70.
- Ellis, P. W., A. M. Page, S. Wood, J. Fargione, Y. J. Masuda, V. Carrasco Denney, C. Moore, T. Kroeger, B. Griscom, J. Sanderman, et al. 2024. The principles of natural climate solutions. *Nature Communications* 15:547.
- Erickson, R. A., and B. A. Rattner. 2020. Moving beyond  $p < 0.05$  in ecotoxicology: A guide for practitioners. *Environmental Toxicology and Chemistry* 39:1657–1669.
- Eyster, H. N., D. S. Srivastava, M. Kreitzman, and K. M. A. Chan. 2022. Functional traits and metacommunity theory reveal that habitat filtering and competition maintain bird diversity in a human shared landscape. *Ecography* 2022.
- Fornacon-Wood, I., H. Mistry, C. Johnson-Hart, J. P. O’Connor, G. J. Price, and C. Faivre-Finn. 2021. A bayesian approach to evaluate the impact of change in igrt protocol using real world data. *Radiotherapy and Oncology* 161.
- Gallistel, C. R. 2009. The importance of proving the null. *Psychological review* 116:439.
- Gilbert, N. A., H. N. Eyster, and E. F. Zipkin. 2023. Propagating uncertainty in ecological models to understand causation. *Frontiers in Ecology and the Environment* 21.
- Griscom, B. W., J. Adams, P. W. Ellis, R. A. Houghton, G. Lomax, D. A. Miteva, W. H. Schlesinger, D. Shoch, J. V. Siikamäki, P. Smith, et al. 2017. Natural climate solutions. *Proceedings of the National Academy of Sciences* 114:11645–11650.
- Gross, M. 2013. The social-ecological co-constitution of nature through ecological restoration: experimentally coping with inevitable ignorance and surprise. *In* S. Lockie, D. A. Sonnenfeld, and D. R. Fisher, eds., *Routledge international handbook of social and environmental change*. Routledge/Taylor & Francis Group, London New York.
- Halsey, L. G. 2019. The reign of the p-value is over: what alternative analyses could we employ to fill the power vacuum? *Biology letters* 15:20190174.
- Holling, C. S., and C. Walters. 1978. Adaptive environmental assessment and management .
- Hooten, M. B., and N. T. Hobbs. 2015. A guide to bayesian model selection for ecologists. *Ecological monographs* 85:3–28.
- IPCC. 2007. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, 2007. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Kery, M., and M. Schaub. 2011. Bayesian population analysis using WinBUGS: a hierarchical perspective. Academic Press.
- Levin, S., T. Xepapadeas, A.-S. Crépin, J. Norberg, A. de Zeeuw, C. Folke, T. Hughes, K. Arrow, S. Barrett, G. Daily, P. Ehrlich, N. Kautsky, K.-G. Mäler, S. Polasky, M. Troell, J. R. Vincent, and B. Walker. 2012. Social-ecological systems as complex adaptive systems: modeling and policy implications. *Environment and Development Economics* 18:111–132.
- Marcot, B. G., R. S. Holthausen, M. G. Raphael, M. M. Rowland, and M. J. Wisdom. 2001. Using bayesian belief networks to evaluate fish and wildlife population viability under land management alternatives from an environmental impact statement. *Forest ecology and management* 153:29–42.
- McNeish, D. 2016. On using bayesian methods to address small sample problems. *Structural Equation Modeling: A Multidisciplinary Journal* 23:750–773.



- Nadeau, C. P., A. K. Fuller, and D. L. Rosenblatt. 2015. Climate-smart management of biodiversity. *Ecosphere* 6:1–17.
- Newton, A., G. Stewart, A. Diaz, D. Golicher, and A. Pullin. 2007. Bayesian belief networks as a tool for evidence-based conservation management. *Journal for Nature Conservation* 15:144–160.
- O’Leary, R. A. 2008. Informed statistical modelling of habitat suitability for rare and threatened species. Ph.D. thesis. Queensland University of Technology.
- Prato, T. 2005. Bayesian adaptive management of ecosystems. *Ecological Modelling* 183:147–156.
- Ripple, W. J., C. Wolf, T. M. Newsome, M. Hoffmann, A. J. Wirsing, and D. J. McCauley. 2017. Extinction risk is most acute for the world’s largest and smallest vertebrates. *Proceedings of the National Academy of Sciences* 114:10678–10683.
- Saunders, S. P., M. T. Farr, A. D. Wright, C. A. Bahlai, J. W. Ribeiro, S. Rossman, A. L. Sussman, T. W. Arnold, and E. F. Zipkin. 2019. Disentangling data discrepancies with integrated population models. *Ecology* 100.
- Stern, E. R., and M. M. Humphries. 2022. Interweaving local, expert, and indigenous knowledge into quantitative wildlife analyses: A systematic review. *Biological Conservation* 266:109444.
- Stinchcombe, J., L. C. Moyle, B. R. Hudgens, P. L. Bloch, S. Chinnadurai, and W. F. Morris. 2002. The influence of the academic conservation biology literature on endangered species recovery planning. *Conservation Ecology* 6.
- Tittensor, D. P., M. Walpole, S. L. Hill, D. G. Boyce, G. L. Britten, N. D. Burgess, S. H. Butchart, P. W. Leadley, E. C. Regan, R. Alkemade, et al. 2014. A mid-term analysis of progress toward international biodiversity targets. *Science* 346:241–244.
- Van de Schoot, R., D. Kaplan, J. Denissen, J. B. Asendorpf, F. J. Neyer, and M. A. Van Aken. 2014. A gentle introduction to bayesian analysis: Applications to developmental research. *Child development* 85:842–860.
- van Zyl, C. J. 2018. Frequentist and bayesian inference: A conceptual primer. *New Ideas in Psychology* 51:44–49.

## Figures




Figure 1: **The iterative nature of Bayesian workflows aligns well with cycles of adaptive management in conservation.** Integrating analytical approaches through cyclical stages of conservation planning and implementation– rather than only in one later stage– is likely to lead to better planning and implementation outcomes.




Figure 2: Proportion papers using Bayes in XX major conservation journals since 2000




Figure 3: Trends in population size over time (left) analyzed with a traditional Fisherian approach using null hypothesis testing (using an  $\alpha$  of 0.05 to reject the null hypothesis of a slope of zero) versus a Bayesian approach, which focuses on the posterior distribution (right).




Figure 4: NCS Example: Uncertainty propagation