

# Temperature and photoperiod interactively drive spring phenology across all species in a temperate forest community

Authors:

D. F. B. Flynn<sup>1,2†</sup> & E. M. Wolkovich<sup>1,2†\*</sup>

*Author affiliations:*

<sup>1</sup>Arnold Arboretum of Harvard University, 1300 Centre Street, Boston, Massachusetts, USA;

<sup>2</sup>Organismic & Evolutionary Biology, Harvard University, 26 Oxford Street, Cambridge, Massachusetts, USA

\*Corresponding author: lizzie@oeb.harvard.edu

†Authors contributed equally

## Abstract

Accurate predictions of future spring plant phenology with continued climate change are critical for robust projections of future growing seasons, plant communities and a related suite of critical ecosystem services. Despite tremendous amounts of data, progress towards prediction has been hindered because the major cues known to drive phenology—chilling temperatures in fall and winter, photoperiod, and spring forcing temperatures—generally covary in nature. Further, research to date using controlled environments to separate these factors suggests that the cues are interactive, meaning accurate predictions of plant responses to climate change will be complex and non-linear (?). Other research, however, has suggested many species may be dominated by one of the three possible cues (?), with a tradeoff between photoperiod and forcing temperature sensitivities, meaning some species' responses would be simple to predict. Here we present results of a full-factorial experiment manipulating all three cues (spring forcing temperatures, photoperiod, and intensity of winter chilling) across 28 woody species and from two North American forests at two latitudes (42.5°N and 46°N). We found responses to photoperiod and temperature cues were largely coordinated across species; namely, species highly sensitive to temperature were also highly sensitive to photoperiod. Chilling exerted the strongest effect on phenology, with more intense chilling at 1.5°C resulting in less pronounced effects than at 4°C. Cues varied across species leading to staggered leafout within each community and supporting the idea that phenology may be a critical aspect of species' temporal niches (??). Our results suggest that predicting the spring phenology of communities will be difficult as all species we studied could have complex, non-linear responses to future warming.

1 Plant phenology—the timing of recurring life history events, such as leafout and flowering—  
2 is critical to the structure and function of ecosystems. Plant phenology determines the timing  
3 of the basal resource in most systems, and thus shapes food webs and mutualistic networks (?).  
4 Spring plant phenology drives local ecosystem properties, from the length of the growing season  
5 to energy balance between land and atmosphere, and scales up to impact global carbon cycles  
6 (?).

7 Phenology is also one of the major biological indicators of climate change, with plant phe-  
8 nology shifting across the globe 4-6 days/°C with warming (?). While this average response is  
9 strikingly consistent when considered across diverse datasets (?), it masks tremendous varia-  
10 tion. Variation is extreme when examined across species (?), but additional variation can be  
11 seen within species across time (???). Understanding this variation has been the goal of much  
12 recent work (?), with research focusing on two major linked aims: (1) identifying and quanti-  
13 fying the environmental cues that drive spring phenology (i.e., leafout and budburst), and (2)  
14 identifying what drives variation in cues between different species.

15 Decades of study on wild species spring phenology—mainly focused on temperate woody  
16 species—show that three major cues drive budburst and leafout: warm spring temperatures  
17 (forcing), length and intensity of winter temperature (chilling), and increasing daylength (pho-  
18 toperiod). Across studies increasing temperatures in the spring appear to be a dominant factor  
19 that controls spring phenology, yet many of these studies have been observational—making it  
20 nearly impossible to tease out the generally co-varying effects of longer days and reduced cold  
21 temperatures, which generally reduce chilling (?). In contrast, studies from controlled envi-  
22 ronments (e.g., growth chambers or greenhouses) have highlighted the additional importance  
23 of photoperiod and chilling (?????), with longer days and increased chilling leading to more  
24 rapid leafout (?). Many of these cues are known to interact and possibly modify one another  
25 in some species. Insufficient chilling may be offset by additional forcing, and photoperiod and  
26 chilling often appear to interact, as long photoperiod enhances cell growth, compensating for  
27 a lack of chilling during plants' winter dormancy (???). Finally, photoperiod and forcing are  
28 known to interactively determine spring phenology—especially through their complex impacts  
29 on dormancy release (?).

30 Yet, while such complexities have been identified in some species, a growing body of hypothe-

ses and experimental studies has suggested many species are dominated by one cue and may lack any response to other cues (?). If true, this would have critical implications for predicting responses to climate change, as a species dominated by a forcing cue could march forever earlier in the timing of leafout with continued warming, while species with strong photoperiod cues could stop advancing at some threshold point (?). Alternatively, if all three cues—temperature, photoperiod, and chilling—are interchangeable then a species experiencing a mild winter with insufficient chilling (as predicted with climate change) could still break bud given sufficiently long photoperiods and warm temperatures (?).

Research to date shows cues clearly vary across species, and recent efforts have focused on understanding and predicting this variation. Studies have focused on attributes of species: origin of species on shorter and longer timescales (??), the successional stage (i.e., pioneer or climax communities) to which species traditionally belong (??), and a variety of possibly related traits (??).

Most of these studies hinge on an often implicit assumption that phenology—by helping define the temporal niche of a species—is a critical axis along which plant species assemble within communities (??). Support for this hypothesis comes from work showing that phenology is often staggered within communities, and from the special case of plant invasions, where research has suggested climate change has provided open temporal niche space for new species to occupy (??).

Improved understanding and predictions of phenology with climate change would benefit from a fuller understanding of the interacting environmental cues that drive phenology within (and eventually across) communities. To this aim we studied how forcing, photoperiod and chilling cues vary across a community of 28 woody plant species from two temperate forest locations, separated by 4°N latitude. We used clipped dormant branches, which have been shown to approximate whole plant responses (?), and forced them in controlled environments that varied forcing temperatures, photoperiod and chilling. We predicted that: (1) Cues would vary across species, driving staggered leafout across the spring, and (2) within-species cues would trade off, such that some species would be dominated by one or another cue, while others would show a mix of cues. To our knowledge this is the first multi-species study to assess all three cues in one experiment through a controlled environment approach, while several studies have done

this for one species (????), all other studies of all three cues have used multiple experiments (e.g., ?) or relied on field sampling to assess one or more cues (e.g., ???).

## Results & Discussion

In total we monitored 2,136 clippings and took over 19,320 observations of phenology (budburst and leafout as assessed by the BBCH scale) in an experiment comprising 12 unique treatments that ran 82 days. Higher forcing temperatures, longer photoperiod, and more chilling all caused large advances in budburst and leafout (Fig. ??, Tables S2-S3).

Forcing temperatures and chilling caused the largest advances in budburst and leafout, and these two effects offset one another, as shown by their interactive delayed response (Fig. ??). The interactive effects of forcing and chilling has been noted repeatedly before (e.g., ??) and highlights that insufficient chilling can be overcome by additional forcing—a hypothesis suggested by recent studies that have found shifting temperature sensitivities in observational data over time (??). We found similar effects of chilling across two different base temperatures, with only minor differences: responses to the colder (1.5°C) chilling treatment were similar or more muted compared to responses to the warmer (4°C) chilling treatment (Fig. ??, Tables S2-S3). This could indicate that either plants cannot assess chilling temperatures below some threshold (??) or that both our chilling treatments met the chilling requirements for our studied species and thus produced similar results.

The two forest sites showed very similar responses, with only a very minor delay in overall timing for the more northern site, and a more pronounced effect of site through its interaction with chilling (Fig. ??, Tables S2-S3). The full effect of chilling depended on site, with a larger impact of site on the lower chilling temperature (1.5°C). This could indicate that chilling requirements vary across populations due to local adaptation, or it could be due to the field chilling experienced before we took cuttings for our experiments (see Table S4). Effects of forcing temperatures and photoperiod were not clearly impacted by site.

At the community level we found that all species were responsive to all cues (forcing temperatures, photoperiod and chilling, Fig. S2-3), and cues were generally staggered such that each species would budburst and leafout at a distinct time compared to other species (Fig. S4). This provides support for the idea that spring phenology is an important component of the temporal

niche (??) for temperate forest species.

In contrast to our expectations that within a species cues would trade off (i.e., a species could be dominated by one cue over all others), we found that species tended to show coordinated cues, especially between forcing and photoperiod (Fig. ??a,c). Thus, a species with a strong response to forcing temperature generally also had a strong response to photoperiod, and similarly a species with a comparatively weak response to forcing also had a weaker response to photoperiod. This was also seen somewhat with chilling (Fig. ??b,d), though we have fewer species with which to assess the relationship (see Methods).

Our finding that all species responded strongly to all three cues is at odds with some recently published work (??), but is coherent with many other studies (e.g., ??) and related process-based models of woody plant phenology (??). The contrasting results may be due to varying methodologies. Our study used controlled environments (growth chambers) to manipulate all three cues, while other studies have used multiple field sampling dates (i.e., sampling once each month from January to March in the northern hemisphere) to assess the effect of one cue, most often chilling (???). As chilling increases across a winter season, so do forcing temperatures and photoperiod as well, meaning it may be hard to fully assess any one cue using this method. Such methods may thus underestimate the full suite of cues that a species uses to control spring phenology. They may, however, have the advantage of providing more realistic environmental conditions by capturing realistic shifts in all three cues across the winter-spring season (?), and thus play an important role in helping predict near-term impacts of climate change. Our findings are probably impacted by collecting in January, when species had likely not had requirements for any cue fully met. Indeed, all methods using clipped end of tree branches are affected by the climate of the season when the experiment was conducted, which provides an additional confounding effect to all such experiments. An alternative reason we may have found contrasting results compared to previous studies could be differences in species sampling: our species list, however, includes a number of species classified as non- or low-responsive to photoperiod (i.e., *Alnus incana*, *Aronia melanocarpa*, ?), making this explanation seems less likely.

Responses to cues were qualitatively similar across both budburst and leafout, but quantitatively varied greatly between the two phenophases (Fig. ??-2), with responses generally greater to leafout (Fig. ??). This change was dramatic for the response to forcing and photoperiod

where the advance in days more than doubled from budburst to leafout (Fig. ??, Table S2-S3). Species cues varied depending on the phenophase considered, meaning species responses also shuffled between the two stages (Fig. ??). For example, shrubs tended to show weaker photoperiod effects to budburst (Fig. ??a) but this was not seen as consistently in the response to leafout (Fig. ??c). This also fundamentally means that the species that bursts bud first will not necessarily leaf out first.

These quantitatively diverging findings for each phenophase suggest complex dynamics in the early season within a community of woody plant species. Increasing evidence suggests the period between budburst and leafout is when plants are at greatest risk of tissue loss from frost (?), and these new insights have come at the same time research suggests spring frosts may increase with climate change (??). For early season species in particular, this period may be critical to their current and future performance. Our results suggest that the cues for each stage are not identical and supports other work suggesting cues on bud swelling and budburst may be distinct from the cues governing the development afterwards (?). Understanding budburst is particularly difficult as it is the first observable event after a series of important physiological events required for budburst (??) and our results echo calls for increased research in this topic (?), which spans both molecular, cellular and whole plant areas of study (???).

### *Conclusions*

Across the two communities we studied, our results suggest species within a community have paced budburst and leafout due to a mix of all three major environmental cues: forcing temperatures, photoperiod and chilling. In contrast to our and others' hypothesis (?), we found no evidence of any species being dominated by one or another cue; instead, species tended to show coordination across cues, especially between forcing and photoperiod cues. Thus, accurately predicting the phenology of any one of our studied species under diverse environmental conditions would require considering how all three cues will change in concert. Shifting climate has already clearly altered forcing and potentially chilling across the globe (??) with trends expected to only continue and possibly accelerate; in contrast, photoperiod has not and will not shift. These trends combined with our results mean that all 28 species we studied could potentially show complex, non-linear responses in the future, with cascading community consequences.

## Methods

### Field sampling

Woody plant cuttings were made in January 2015 for 28 species which occurred in both Harvard Forest (HF, 42.5°N, 72.2°W) and the Station de Biologie des Laurentides in St-Hippolyte, Québec (SH, 45.9°N, 74.0°W). The typical late January temperatures are -3.4 and -22°C, respectively. Weather station data from each field site was obtained for calculations of chilling units (see Table S4).

Species were chosen based on the dominant forest vegetation at each site, aiming to maximize the number of shared species between the two sites. Of the 28 species, at least 19 occurred at both sites. Comparing only shared species, the mean days to budburst and leafout across all treatments for Harvard Forest and St. Hippolyte was 25.6/36.8 and 24.8/36.1 days, respectively (Table S1). For each species, up to 15 representative healthy, mature individuals with branches accessible by pole pruners from the ground were tagged in late summer and fall 2014. In winter 2015, six individuals were located and 4-16 cuttings taken from each individual, depending on size of the individual and number of treatments to be applied. Cuttings were kept cold and transported back to the Arnold Arboretum in Boston, MA.

### Growth Chamber Study

Cuttings were placed in growth chambers at the Arnold Arboretum in Erlenmeyer flasks distilled water, with water changed every 7-10 days. The base of cuttings was re-cut at each water change under water to prevent callusing. For 11 of the 28 species, sufficient cuttings were obtained from each individual tree to apply the full set of 12 experimental treatments: 2 temperature (20°C / 10°C warm vs. 15°C / 5°C cool) x 2 photoperiod (12 vs. 8 h) x 3 chilling (no additional chilling, additional 33 d at 4°C, or 33 d at 1.5°C) treatments. For the remaining 17 species, only sufficient cuttings were obtained to apply the temperature and photoperiod treatments, without the additional chilling levels. The total number of cuttings for a given species thus ranged from 24 to 144, depending on presence at each site and application of the chilling treatment (Fig. S1). Lighting was a combination of halogen incandescent bulbs and T5HO fluorescent lamps with the lamploft adjusted to provide roughly 400  $\mu\text{moles}/\text{m}^2/\text{s}$  as measured by Apogee QSO-A5E quantum PAR light sensors in each chamber (sensor set to the height of the cuttings). Treatments were rotated across chambers every two weeks, as was flask position within chamber,



to remove any possible bias of chamber or flask position.

Phenology of the cuttings was assessed using a modified BBCH scale ?, with observations on each of the 2,136 cuttings made every 2-3 days for the course of the 82-day experiment, a total of 48 observation days. The phenological stages assessed in the present study are budburst, defined as beginning of sprouting or bud breaking or shoot emergence (Code 07 in ?) and leaf out, defined as first leaves unfolded (Code 11 in (?)). Additional stages up to flowering and stem elongation were also recorded and we provide a photographic guide to help visualize stages across species(?). In total, we made 19,318 phenological observations at the cutting level.

### Statistical analysis

For the two phenology responses measured, budburst and leafout, we fit Bayesian mixed-effect hierarchical models using site, warming, photoperiod, and chilling treatments as predictors (fixed effects) and species as a modeled groups (random effects). This approach allowed us to calculate the posterior probabilities of the effects for each of the abiotic drivers individually and interactively across all species sampled. For each model, two-way interactions for effects of site, warming, and each of the chilling treatments were included. The models were fit using the programming languages Stan (?)([www.mc-stan.org](http://www.mc-stan.org)), accessed via the *rstan* package (version 2.15.1) in R (?), version 3.3.3.

The model was fit as follows:

$$\begin{aligned}
y_i \sim N(&\alpha_{sp[i]} + \beta_{site_{sp[i]}} + \beta_{temperature_{sp[i]}} + \beta_{photoperiod_{sp[i]}} + \beta_{chilling1_{sp[i]}} + \beta_{chilling2_{sp[i]}} \\
&+ \beta_{temperature \times photoperiod_{sp[i]}} + \beta_{temperature \times site_{sp[i]}} + \beta_{photoperiod \times site_{sp[i]}} \\
&+ \beta_{temperature \times chilling1_{sp[i]}} + \beta_{temperature \times chilling2_{sp[i]}} \\
&+ \beta_{photoperiod \times chilling1_{sp[i]}} + \beta_{photoperiod \times chilling2_{sp[i]}} \\
&+ \beta_{site \times chilling1_{sp[i]}} + \beta_{site \times chilling2_{sp[i]}})
\end{aligned}$$

Each of the 14  $\beta$  coefficients was modeled at the species level, as follows:

$$\begin{aligned}
&1. \beta_{site_{sp}} \sim N(\mu_{site}, \sigma^2_{site}) \\
&\dots \\
&14. \beta_{site \times chilling2_{sp}} \sim N(\mu_{site \times chilling2}, \sigma^2_{site \times chilling2})
\end{aligned}$$

For the  $\mu$  and  $\sigma$  parameters, weakly informative priors were chosen.

We ran four chains simultaneously, with 4 000 warm-up iterations followed by 3 997 sampling iterations, resulting in 15 998 posterior samples for each parameter. We used a non-centered parameterization on all interactions terms and assessed good model performance through  $\hat{R}$  close to 1 and high  $n_{eff}$  as well as visual consideration of chain convergence and posteriors.

## Acknowledgements

We thank T. Savas for technical and field assistance as well as countless observing hours, J. Samaha, H. Eyster for help with field collections, and E. Borjigin-Wang, J. Samaha, N. Farrant, and T. Chen for recording chamber observations.

## References

- Augspurger, C. K. 2009. Spring 2007 warmth and frost: phenology, damage and refoliation in a temperate deciduous forest. *Functional Ecology* 23:1031–1039.
- Basler, D., and C. Körner. 2012. Photoperiod sensitivity of bud burst in 14 temperate forest tree species. *Agricultural and Forest Meteorology* 165:73–81.
- . 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* 34:377–388.
- Caffarra, A., and A. Donnelly. 2011. The ecological significance of phenology in four different tree species: effects of light and temperature on bud burst. *Int J Biometeorol* 55:711–21.
- Caffarra, A., A. Donnelly, I. Chuine, and M. B. Jones. 2011. Modelling the timing of *Betula pubescens* bud-burst. I. Temperature and photoperiod: A conceptual model. *Climate Research* 46:147.
- Carpenter, B., A. Gelman, M. Hoffman, D. Lee, B. Goodrich, M. Betancourt, M. A. Brubaker, J. Guo, P. Li, and R. Allen. 2016. Stan: A probabilistic programming language. *Journal of Statistical Software* (in press).
- Chuine, I. 2000. A unified model for budburst of trees. *Journal of Theoretical Biology* 207:337–347.
- Chuine, I., M. Bonhomme, J. M. Legave, I. G. de Cortazar-Atauri, G. Charrier, A. Lacointe, and T. Ameglio. 2016. Can phenological models predict tree phenology accurately in the future? the unrevealed hurdle of endodormancy break. *Global Change Biology* 22:3444–3460.
- Chuine, I., G. Cambon, and P. Comtois. 2000. Scaling phenology from the local to the regional level: advances from species-specific phenological models. *Global Change Biology* 6:943–952.
- Chuine, I., and P. Cour. 1999. Climatic determinants of budburst seasonality in four temperate-zone tree species. *New Phytologist* 143:339–349.
- Cleland, E. E., I. Chuine, A. Menzel, H. A. Mooney, and M. D. Schwartz. 2007. Shifting plant phenology in response to global change. *Trends in ecology & evolution* 22:357–365.

- Cook, B. I., E. M. Wolkovich, and C. Parmesan. 2012. Divergent responses to spring and winter warming drive community level flowering trends. *Proceedings of the National Academy of Sciences of the United States of America* 109:9000–9005.
- Dai, J., H. Wang, and Q. Ge. 2013. The decreasing spring frost risks during the flowering period for woody plants in temperate area of eastern China over past 50 years. *Journal of Geographical Sciences* 23:641–652.
- Falusi, M., and R. Calamassi. 1996. Geographic variation and bud dormancy in beech seedlings (*Fagus sylvatica* L). *Annales des Sciences Forestières* 53:967–979.
- Finn, G. A., A. E. Straszewski, and V. Peterson. 2007. A general growth stage key for describing trees and woody plants. *Annals of Applied Biology* 151:127–131.
- Foley, M. E., J. V. Anderson, and D. P. Horvath. 2009. The effects of temperature, photoperiod, and vernalization on regrowth and flowering competence in *Euphorbia esula* (Euphorbiaceae) crown buds. *Botany* 87:986–992.
- Fu, Y. S. H., H. F. Zhao, S. L. Piao, M. Peaucelle, S. S. Peng, G. Y. Zhou, P. Ciais, M. T. Huang, A. Menzel, J. P. Uelas, Y. Song, Y. Vitasse, Z. Z. Zeng, and I. A. Janssens. 2015. Declining global warming effects on the phenology of spring leaf unfolding. *Nature* 526:104–+.
- Ghelardini, L., A. Santini, S. Black-Samuelsson, T. Myking, and M. Falusi. 2010. Bud dormancy release in elm (*Ulmus spp.*) clones—a case study of photoperiod and temperature responses. *Tree physiology* 30:264–274.
- Gotelli, N. J., and G. R. Graves. 1996. The temporal niche. Pages 95–111 *in* N. J. Gotelli and G. R. Graves, eds. *Null Models in Ecology*. Smithsonian Institution, Washington, DC.
- Guy, R. D. 2014. The early bud gets to warm. *New Phytologist* 202:7–9.
- Harrington, C. A., P. J. Gould, and J. B. St Clair. 2010. Modeling the effects of winter environment on dormancy release of Douglas-fir. *Forest Ecology and Management* 259:798–808.
- Heide, O. 1993. Daylength and thermal time responses of budburst during dormancy release in some northern deciduous trees. *Physiologia Plantarum* 88:531–540.

- IPCC. 2014. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Körner, C., and D. Basler. 2010. Phenology under global warming. *Science* 327:1461–1462.
- Laube, J., T. H. Sparks, C. Baessler, and A. Menzel. 2015. Small differences in seasonal and thermal niches influence elevational limits of native and invasive balsams. *Biological Conservation* 191:682–691.
- Laube, J., T. H. Sparks, N. Estrella, J. Hoffer, D. P. Ankerst, and A. Menzel. 2014. Chilling outweighs photoperiod in preventing precocious spring development. *Global Change Biology* 20:170–182.
- Lechowicz, M. J. 1984. Why do temperate deciduous trees leaf out at different times? Adaptation and ecology of forest communities. *The American Naturalist* 124:821–842.
- Lenz, A., G. Hoch, Y. Vitasse, and C. Körner. 2013. European deciduous trees exhibit similar safety margins against damage by spring freeze events along elevational gradients. *New Phytologist* 200:1166–1175.
- Loreau, M., and C. de Mazancourt. 2008. Species synchrony and its drivers: Neutral and nonneutral community dynamics in fluctuating environments. *American Naturalist* 172:E48–E66.
- Morin, X., M. J. Lechowicz, C. Augspurger, J. O’Keefe, D. Viner, and I. Chuine. 2009. Leaf phenology in 22 north american tree species during the 21st century. *Global Change Biology* 15:961–975.
- Myking, T., and O. Heide. 1995. Dormancy release and chilling requirement of buds of latitudinal ecotypes of *Betula pendula* and *B. pubescens*. *Tree Physiology* 15:697–704.
- Polgar, C., A. Gallinat, and R. B. Primack. 2014. Drivers of leaf-out phenology and their implications for species invasions: insights from thoreau’s concord. *New phytologist* 202:106–115.
- R Development Core Team. 2017. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

- Richardson, A. D., D. Y. Hollinger, D. B. Dail, J. T. Lee, J. W. Munger, and J. O'keefe. 2009. Influence of spring phenology on seasonal and annual carbon balance in two contrasting new england forests. *Tree Physiology* 29:321–331.
- Rinne, P. L. H., A. Welling, J. Vahala, L. Ripel, R. Ruonala, J. Kangasjarvi, and C. van der Schoot. 2011. Chilling of dormant buds hyperinduces FLOWERING LOCUS T and recruits GA-Inducible 1,3-beta-Glucanases to reopen signal conduits and release dormancy in *Populus*. *Plant Cell* 23:130–146.
- Rutishauser, T., J. Luterbacher, C. Defila, D. Frank, and H. Wanner. 2008. Swiss spring plant phenology 2007: Extremes, a multi-century perspective, and changes in temperature sensitivity. *Geophysical Research Letters* 35.
- Savas, T., D. F. B. Flynn, and E. M. Wolkovich. 2017. A standardized photographic guide to woody plant spring phenology.
- Singh, R. K., T. Svystun, B. AlDahmash, A. M. Jonsson, and R. P. Bhalerao. 2017. Photoperiod- and temperature-mediated control of phenology in trees - a molecular perspective. *New Phytologist* 213:511–524.
- Skuterud, R., J. Dietrichson, et al. 1994. Budburst in detached birch shoots (*betula pendula*) of different varieties winter-stored in darkness at three different temperatures. *Silva Fennica* .
- Søgaard, G., Ø. Johnsen, J. Nilsen, and O. Junttila. 2008. Climatic control of bud burst in young seedlings of nine provenances of norway spruce. *Tree Physiology* 28:311–320.
- Sønsteby, A., and O. M. Heide. 2014. Chilling requirements of contrasting black currant (*ribes nigrum* l.) cultivars and the induction of secondary bud dormancy. *Scientia Horticulturae* 179:256–265.
- Stocker, T., D. Qin, and G. Platner. 2013. Climate change 2013: The physical science basis. Working Group I Contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. Summary for Policymakers (IPCC, 2013) .
- Vitasse, Y., and D. Basler. 2014. Is the use of cuttings a good proxy to explore phenological

responses of temperate forests in warming and photoperiod experiments? *Tree Physiology*  
34:174–183.

Vitasse, Y., A. Lenz, and C. Korner. 2014. The interaction between freezing tolerance and  
phenology in temperate deciduous trees. *Frontiers in Plant Science* 5.

Weinberger, J. H. 1950. Chilling requirements of peach varieties. *Proceedings of the American*  
*Society for Horticultural Science* 56:122–128.

Willis, C. G., B. R. Ruhfel, R. B. Primack, A. J. Miller-Rushing, J. B. Losos, and C. C. Davis.  
2010. Favorable climate change response explains non-native species' success in Thoreau's  
woods. *PLoS ONE* 5:e8878.

Wolkovich, E. M., B. I. Cook, J. M. Allen, T. M. Crimmins, J. L. Betancourt, S. E. Travers,  
S. Pau, J. Regetz, T. J. Davies, N. J. Kraft, T. R. Ault, K. Bolmgren, S. J. Mazer, G. J.  
McCabe, B. J. McGill, C. Parmesan, N. Salamin, M. D. Schwartz, and E. E. Cleland. 2012.  
Warming experiments underpredict plant phenological responses to climate change. *Nature*  
485:494–7.

Wolkovich, E. M., B. I. Cook, and T. J. Davies. 2014. Progress towards an interdisciplinary  
science of plant phenology: building predictions across space, time and species diversity. *New*  
*Phytologist* 201:1156–1162.

Wolkovich, E. M., T. J. Davies, H. Schaefer, E. E. Cleland, B. I. Cook, S. E. Travers, C. G.  
Willis, and C. Davis. 2013. Temperature-dependent shifts in phenology contribute to the  
success of exotic species with climate change. *American Journal of Botany* 100:1407– 1421.

Worrall, J., and F. Mergen. 1967. Environmental and genetic control of dormancy in *picea abies*.  
*Physiologia Plantarum* 20:733–745.

Yu, H. Y., E. Luedeling, and J. C. Xu. 2010. Winter and spring warming result in delayed spring  
phenology on the Tibetan Plateau. *Proceedings of the National Academy of Sciences of the*  
*United States of America* 107:22151–22156.

Zohner, C. M., B. M. Benito, J. D. Fridley, J. C. Svenning, and S. S. Renner. 2017. Spring

337        predictability explains different leaf-out strategies in the woody floras of north america, europe  
338        and east asia. *Ecology Letters* 20:452–460.

339        Zohner, C. M., B. M. Benito, J. C. Svenning, and S. S. Renner. 2016. Day length unlikely to  
340        constrain climate-driven shifts in leaf-out times of northern woody plants. *Nature Climate*  
341        Change 6:1120–+.



## Figures




Figure 1: Effects of forcing temperature, photoperiod, chilling and site on budburst (a) and leafout (b) days across 28 species. Dots and bars show mean and 50% credible intervals from a Bayesian hierarchical model that also incorporated species-level variations (see Tables S2-S3; Figs. 1, S2-S3). Advances in phenology are shown by negative numbers, while delays are shown as positive. Forcing temperatures and photoperiods were two levels each (see Methods), and chilling treatments were applied for 33 days. Budburst and leafout images from ?.




Figure 2: Coordinated effects of photoperiod, temperature and chilling across species: Crosses and bars show mean and 50% credible intervals from a Bayesian hierarchical model (see Tables S2-S3; Figs. 1, S2-S3). For visualization purposes, species names are represented by the first three letters of the genus and first three letters of the species epithet (see Table S1 for full species names and Fig. S5 for additional version of figure).