

# Supplemental Materials

## Temperature and photoperiod interactively drive spring phenology across all species in a temperate forest community

Flynn & Wolkovich

### Supplemental Methods

#### *Chilling calculations*

The cuttings were harvested in late January 2015, and thus experienced substantial natural chilling by the time they were harvested. Using weather station data from the Harvard Forest and St. Hippolyte site, chilling hours (below 7.2°C), Utah Model chill portions (hours below 7.2°C and between 0°C and 7.2°C) and Dynamic Model (Erez et al., 1988) chill portions were calculated both for the natural chilling experienced by harvest and the chilling experienced in the 4°C and 1.5°C treatments. The Utah Model and Dynamic Model of chill portions account for variation in the amount of chilling accumulated at different temperatures, with the greatest chilling occurring approximately between 5-10°C, and fewer chill portions accumulating at low temperatures and that higher temperatures can reduce accumulated chilling effects. The two differ in the parameters used to determine the shape of the chilling accumulation curve, with the Dynamic Model being shown to be the most successful in predicting phenology for some woody species (Luedeling et al., 2009). With both the Utah and Dynamic model, the more severe chilling treatment resulted in fewer calculated chilling portions.

#### *Non-leafouts*

Across all treatments, 20.2% of the cuttings did not break bud or leaf out. Across species, there was no overall predictive effect of temperature, photoperiod, chilling, or site on the propensity to fail to leaf out. Species ranged from complete leaf out (*Hamamelis*) to only 50% leaf out (*Fagus grandifolia*, *Acer saccharum*) across all treatments. The percent of nonleaf outs by site was similar, with 20.6% of Harvard Forest and 19.7% of St. Hippolyte samples failing to leaf out. Examining individual species, there was an interaction of temperature by day length for selected species, with greater failure to leaf out in cool, short-day conditions for *Acer pensylvanicum* and *Acer saccharum*. Site effects were inconsistent, with greater failure to leaf out for cuttings from St. Hippolyte in *Acer rubrum* and *Fagus grandifolia*, and from Harvard Forest in *Acer saccharum*.

#### *Additional method details*

In our figures we show 50% credible intervals because our interest is in the most likely value for the parameter and because 50% intervals are computationally stable. For those interested in a more traditional significance-testing approach, we provide 95% credible intervals in Table S2-S3.

### References

- Basler, D., and C. Körner. 2014. Photoperiod and temperature responses of bud swelling and bud burst in four temperate forest tree species. *Tree Physiology* 34:377–388.
- Erez, A., S. Fishman, Z. Gat, and G. A. Couvillon. 1988. Evaluation of winter climate for breaking bud rest using the dynamic model. *Acta Horticulturae* pages 76–89.
- Luedeling, E., M. Zhang, G. McGranahan, and C. Leslie. 2009. Validation of winter chill models using historic records of walnut phenology. *Agricultural and Forest Meteorology* 149:1854–1864.

## Supplemental Figures and Tables

Table S1: Mean leafout and budburst days (across all treatments, based on raw data) for the 28 species at both Harvard Forest (HF), USA and St. Hippolyte (SH), Canada

Species	Budburst.HF	Budburst.SH	Leafout.HF	Leafout.SH
<i>Acer pensylvanicum</i>	16.40	18.33	40.88	46.94
<i>Acer rubrum</i>	22.40	25.15	40.59	44.40
<i>Acer saccharum</i>	44.96	36.48	57.07	46.88
<i>Alnus incana subsp. rugosa</i>	32.91	25.36	45.15	44.36
<i>Aronia melanocarpa</i>	13.62		29.83	
<i>Betula alleghaniensis</i>	19.67	20.77	33.51	34.64
<i>Betula lenta</i>	29.83		50.57	
<i>Betula papyrifera</i>	16.89	18.04	28.71	35.63
<i>Corylus cornuta</i>	24.86	19.04	33.95	30.38
<i>Fagus grandifolia</i>	41.82	43.13	48.54	46.90
<i>Fraxinus nigra</i>	38.00	38.00	52.28	46.91
<i>Hamamelis virginiana</i>	43.67		47.38	
<i>Ilex mucronatus</i>	15.80	15.49	26.97	25.15
<i>Kalmia angustifolia</i>	30.25	32.48	37.80	42.20
<i>Lonicera canadensis</i>	16.91	15.75	28.26	25.08
<i>Lyonia ligustrina</i>	30.87		49.50	
<i>Nyssa sylvatica</i>	31.65		52.87	
<i>Populus grandidentata</i>	33.43	31.23	46.21	45.17
<i>Prunus pensylvanica</i>	17.81	16.21	32.13	29.65
<i>Quercus alba</i>	45.23		52.91	
<i>Quercus rubra</i>	36.43	33.57	45.02	42.80
<i>Quercus velutina</i>	52.09		59.16	
<i>Rhamnus frangula</i>	32.38		37.29	
<i>Rhododendron prinophyllum</i>	29.25		52.14	
<i>Spiraea alba</i>	18.00	20.21	25.94	24.62
<i>Vaccinium myrtilloides</i>	13.12	17.27	27.00	28.95
<i>Viburnum cassinoides</i>	15.41	18.46	16.80	18.71
<i>Viburnum lantanoides</i>	31.25	27.54	32.02	26.41

Table S2: Summary of mixed-effects model of budburst day. See also Figure S2.

	mean	sd	2.5%	50%	97.5%	Rhat
Forcing Temperature	-8.82	1.05	-10.87	-8.82	-6.73	1.00
Photoperiod	-4.53	0.90	-6.28	-4.53	-2.73	1.00
Chilling 4 °C	-15.82	2.13	-20.06	-15.82	-11.59	1.00
Chilling 1.5 °C	-13.13	2.00	-17.04	-13.14	-9.08	1.00
Site	1.31	1.08	-0.84	1.31	3.45	1.00
Forcing Temperature $\times$ Photoperiod	-0.62	0.79	-2.14	-0.63	0.95	1.00
Forcing Temperature $\times$ Chilling 4 °C	9.09	1.09	6.94	9.09	11.23	1.00
Forcing Temperature $\times$ Chilling 1.5 °C	9.78	1.17	7.50	9.79	12.07	1.00
Photoperiod $\times$ Chilling 4 °C	-0.26	1.11	-2.48	-0.26	1.98	1.00
Photoperiod $\times$ Chilling 1.5 °C	-0.14	1.25	-2.63	-0.13	2.27	1.00
Forcing Temperature $\times$ Site	-1.51	0.85	-3.13	-1.51	0.16	1.00
Photoperiod $\times$ Site	-0.08	0.81	-1.70	-0.08	1.49	1.00
Site $\times$ Chilling 4 °C	-2.26	1.21	-4.64	-2.25	0.13	1.00
Site $\times$ Chilling 1.5 °C	-3.47	1.34	-6.10	-3.48	-0.85	1.00

Table S3: Summary of mixed-effects model of leafout day. See also Figure S3.

	mean	sd	2.5%	50%	97.5%	Rhat
Forcing Temperature	-19.06	1.04	-21.10	-19.05	-17.05	1.00
Photoperiod	-11.19	0.86	-12.91	-11.18	-9.53	1.00
Chilling 4 °C	-17.44	2.07	-21.55	-17.42	-13.34	1.00
Chilling 1.5 °C	-15.84	2.05	-20.02	-15.80	-11.88	1.00
Site	1.34	1.24	-1.12	1.35	3.76	1.00
Forcing Temperature $\times$ Photoperiod	3.68	0.85	2.06	3.66	5.39	1.00
Forcing Temperature $\times$ Chilling 4 °C	10.30	1.17	8.01	10.30	12.62	1.00
Forcing Temperature $\times$ Chilling 1.5 °C	11.22	1.32	8.61	11.24	13.81	1.00
Photoperiod $\times$ Chilling 4 °C	0.79	1.18	-1.47	0.79	3.16	1.00
Photoperiod $\times$ Chilling 1.5 °C	2.35	1.32	-0.32	2.37	4.87	1.00
Forcing Temperature $\times$ Site	-0.50	0.83	-2.11	-0.50	1.15	1.00
Photoperiod $\times$ Site	-0.87	0.83	-2.51	-0.87	0.75	1.00
Site $\times$ Chilling 4 °C	-1.75	1.28	-4.26	-1.76	0.77	1.00
Site $\times$ Chilling 1.5 °C	-3.35	1.51	-6.35	-3.34	-0.41	1.00

Table S4: Chill units in field and field and growth chamber conditions.

Site	Treatment	Chilling Hours	Utah Model	Chill portions
Harvard Forest	Field chilling	892	814.50	56.62
	4.0 °C x 30 d	2140	2062.50	94.06
	1.5 °C x 30 d	2140	1702.50	91.17
St. Hippolyte	Field chilling	682	599.50	44.63
	4.0 °C x 30 d	1930	1847.50	82.06
	1.5 °C x 30 d	1930	1487.50	79.18




Figure S1: Responses of 28 woody plant species to photoperiod and temperature cues for leafout. Color of circle reflect unmodeled data on average leaf out day across treatments, across sites of origin, while size of circle represents the total number of clippings in the experiment—this varies mainly based on whether the species was found at both sites and whether it was exposed to all three chilling treatments.




Figure S2: Model estimates of effects of each predictor on budburst, including species-level effects. Dots and bars show mean and 50% credible intervals.




Figure S3: Model estimates of leafout, including species-level effects. Dots and bars show mean and 50% credible intervals.




Figure S4: Model estimates of sensitivity to warming, photoperiod, and chilling, compared to day of budburst (upper panels) or leafout (lower panels) across all experimental conditions.






Figure S5: Effects of photoperiod, temperature and chilling across species: Similar to Fig. 2 from main text, but designed to make species names easier to read by adjusting axes (note that axes vary across rows and columns) and removing bars showing credible intervals.




Figure S6: Effects of photoperiod, temperature and chilling across species: Similar to Fig. 2 from main text, but showing results with 1.5°C chilling treatment, and including relationship between chilling and photoperiod. Crosses and bars show mean and 50% credible intervals.