

Supplemental materials for Phenological sequences: How early-season events define those that follow

A.K. Ettinger, S. Gee, and E.M. Wolkovich

May 28, 2018

Supplemental Tables

Table S1: Summary of linear models for relationships between later phenophases and earlier phenophases, as shown in Figure 3 in the main text. Two types of linear models were fit: those with the intercept only estimated and a forced slope of one, and those with both the slope and intercept estimated (i.e., a standard regression model). All models were fit with the species-level mean day-of-year of the later phenological stages as the response variable, and mean day-of-year of earlier phenostage as the explanatory variable.

previous phenostage model	forced slope model			standard regression model				
	intercept	r ²	aic	intercept	slope	p	r ²	aic
leafout vs. budburst	8.94	0.10	164.78	65.84	0.53	<0.001	0.44	155.01
flowering vs. budburst	23.83	0.17	225.55	3.18	1.17	0.039	0.17	227.45
fruiting vs. budburst	140.71	0.13	260.44	-32.03	2.42	0.029	0.19	260.48
senescence vs. budburst	158.84	-0.12	210.97	243.88	0.30	0.427	0.03	209.43
flowering vs. leafout	14.90	0.23	223.60	-105.28	1.92	0.005	0.30	223.26
fruiting vs. leafout	131.77	0.08	261.78	-42.56	2.33	0.097	0.12	262.74
senescence vs. leafout	149.90	-0.07	209.74	237.39	0.33	0.484	0.02	209.58
fruiting vs. flowering	116.87	0.29	255.39	109.80	1.05	0.006	0.29	257.37
senescence vs. flowering	135.00	-1.79	233.74	261.65	0.13	0.332	0.04	209.08
senescence vs. fruiting	111.97	-5.43	254.66	235.89	0.17	0.008	0.27	202.28

Table S2: Summary of linear models for relationships between later phenophases and interphase duration, as shown in Figure 4 in the main text. Linear models were fit with the species-level mean day-of-year of the later phenological stages as the response variable, and the number of days in each previous interphase duration as the explanatory variable. The random.slopes column gives the range in which 95 percent of slopes in the randomization occur (between 2.5 percent and 97.5 percent).

interphase model	intercept	slope	r ²	p	aic	random.slopes
leafout vs. leafout-budburst	128.98	0.20	0.04	0.374	168.58	0.45-1.57
flowering vs. flowering-budburst	121.18	1.03	0.88	<0.001	180.24	0.83-1.17
fruiting vs. fruiting-budburst	114.31	1.05	0.97	<0.001	178.39	0.92-1.08
senescence vs. senescence-budburst	151.87	0.81	0.74	<0.001	176.81	0.77-1.21
flowering vs. flowering-leafout	129.28	1.10	0.93	<0.001	167.11	0.85-1.13
fruiting vs. fruiting-leafout	126.69	1.03	0.98	<0.001	168.41	0.93-1.07
senescence vs. fruiting-leafout	149.22	0.88	0.82	<0.001	167.30	0.82-1.18
fruiting vs. fruiting-flowering	143.56	1.02	0.74	<0.001	232.15	0.72-1.22
senescence vs. senescence-flowering	247.41	0.25	0.17	0.041	205.51	0.59-1.37
senescence vs. senescence-fruiting	282.11	-0.08	0.05	0.296	208.91	0.53-1.45

Table S3: Summary of linear models for relationships between later phenophases and interphase duration, with interphase calculated as the time between number of days between the end date of the previous phase and the start date of the later phase. Linear models were fit with the species-level mean day-of-year of the later phenological stages as the response variable, and the number of days in each previous interphase duration as the explanatory variable. The random.slopes column gives the range in which 95 percent of slopes in the randomization occur (between 2.5 percent and 97.5 percent).

interphase model	intercept	slope	r ²	p	aic	random.slopes
leafout vs. leafout-budburst	130.69	-0.00	0.00	0.987	169.46	0.67-1.34
flowering vs. flowering-budburst	145.59	0.89	0.75	<0.001	197.42	0.84-1.16
fruiting vs. fruiting-budburst	145.89	1.00	0.93	<0.001	198.54	0.92-1.08
senescence vs. senescence-budburst	196.43	0.62	0.58	<0.001	188.22	0.81-1.2
flowering vs. flowering-leafout	132.82	1.09	0.96	<0.001	150.15	0.86-1.14
fruiting vs. fruiting-leafout	129.33	1.03	0.99	<0.001	151.28	0.93-1.07
senescence vs. fruiting-leafout	134.86	0.99	0.90	<0.001	153.89	0.8-1.21
fruiting vs. fruiting-flowering	148.23	0.99	0.72	<0.001	234.21	0.75-1.25
senescence vs. senescence-flowering	249.92	0.23	0.16	0.047	205.75	0.62-1.35
senescence vs. senescence-fruiting	280.53	-0.17	0.19	0.029	204.81	0.43-1.49

Supplemental Figures




Figure S1: **Individual tree phenology during the 2015 growing season, ordered by species-level mean first-flower dates.** Growth phenology is shown for budburst (from its mean start day-of-year to the mean start day-of-year for leafout, across all individuals within a species), leafout (from the mean day-of-year when fully-expanded leaves were first observed through the start of senescence), and senescence (from the mean day-of-year when leaves first began changing color through the mean day-of-year when more than 95% of leaves on the tree had changed color). Reproductive phenology is shown for flowering (from the mean day-of-year when flowers first appeared to the mean day-of-year when fruits first appeared, across all individuals within a species) and fruiting (from the mean day-of-year when fruits first appeared to the mean day-of-year when more than 95% of fruits were first observed as ripe).