

Soil moisture interacts with temperature to affect plant phenology

A.K. Ettinger^{1,2,a}, J.S. Dukes^{3,b}, M.R. Johnston^{4,c}, C.R. Rollinson^{5,d}, and E.M.
Wolkovich^{1,4,6,e}

¹Arnold Arboretum of Harvard University, Boston, Massachusetts 02131, USA

²Tufts University, Medford, Massachusetts 02155, USA

³Department of Forestry & Natural Resources and Department of Biological Sciences,
Purdue University, West Lafayette, Indiana 47907, USA

⁴Department of Organismic & Evolutionary Biology, Harvard University, Cambridge,
Massachusetts 02138, USA

⁵The Morton Arboretum, Lisle, Illinois 60532, USA

⁶Forest & Conservation Sciences, Faculty of Forestry, University of British Columbia,
Vancouver, BC, Canada

^aCorresponding author; email: aettinger@fas.harvard.edu; phone: 781-296-4821; mailing
address: 1300 Centre Street, Boston, Massachusetts 02140, USA

August 11, 2020

Author contributions: All authors conceived of this manuscript, which began at a Radcliffe Exploratory Seminar in 2016, and all authors contributed to manuscript revisions. AKE and EMW conceived of the idea for the literature review, database compilation, and related Radcliffe Exploratory Seminar. AKE compiled the datasets; AKE analyzed the data and created the figures.

Data Accessibility The data reported in this paper are from the MC3E and ExPhen databases, which are

available at KNB (Ettinger and Wolkovich, 2018; ?)

Running title Soil moisture affects phenology

Key words global warming, warming experiment, microclimate, phenology, bud-burst, leaf-out, flowering, fruiting, senescence

Paper type ‘Brief communication’ for *American Journal of Botany*

Abstract

Past studies of phenology responses to recent climate change have focused on temperature as a driver of observed shifts. However, soil moisture is also affected by climate change and likely to alter biological responses. Here we synthesize microclimate and phenology data from climate change experiments to show how soil moisture interacts with temperature to affect plant phenology. We find that soil drying generally delays plant phenology, especially for budburst, a rate of 0.42 days per percent reduction in soil VWC, on average. These effects vary widely across sites and species. This rate suggests that, on average, climate change-induced shifts soil moisture will play only a small role in altering future phenology, compared to shifts in temperature. This is both because of the strong sensitivity of plant phenology to temperature and because of the large magnitude of projected shifts in temperature, compared to shifts in soil moisture. Nonetheless, although effects of soil moisture are comparably small on average, sensitivity to soil moisture varied dramatically by species, and soil moisture levels differed by site and among years. Thus, soil moisture is likely to play a role in phenological shifts with climate change for some species, in some locations and years. Quantifying phenological sensitivity to changes in soil moisture will be useful for improving forecasts of shifts in phenology with future climate change.

INTRODUCTION

Climate change is affecting organisms by altering temperature and soil moisture around the world (Parmesan, 2006; ?). One of most widespread biological responses to climate change is a shift in phenology, the timing of recurring biological events, which has occurred at rates of 2.3-5.1 days per decade (Parmesan, 2006; Poloczanska et al., 2013; Root et al., 2003). Shifts in plant phenology are the most widely documented, with spring phenology (budburst, leaf-out, and flowering) occurring earlier in recent years (Wolkovich et al., 2013), and senescence occurring later (Taylor et al., 2008; Delpierre et al., 2009).

Phenological shifts are typically attributed to warming temperature, a known and well-studied driver of plant phenology. The timing of spring budburst, for example, depends on temperature through both chilling (the prolonged exposure to cold temperatures after growth cessation in the fall) and forcing (exposure to warm temperatures). Recent trends of advancing phenology may be due to increases in both/either chilling and/or forcing with global warming (Fujisawa and Kobayashi, 2010; Ibanez et al., 2010; Cook et al., 2012). In places where delays in spring phenology have occurred, reductions in winter chilling are thought to be the cause (Yu et al., 2010).

Effects of altered precipitation and soil moisture on phenology have received less attention, but are likely to be important drivers of plant phenology. For example, budburst, flowering, and leaf drop are affected by tree water status in dry ecosystems (Essiamah and Eschrich, 1986; Reich and Borchert, 1984; van Schaik et al., 1993). Budburst can be slowed by water stress through inhibiting cell elongation (Essiamah and Eschrich, 1986), and growing season start may be delayed by drought in grasslands Cui et al. (2017). Flowering phenology, on the other hand, can be advanced by drought conditions (Hamann et al., 2018).

Here we conduct a meta-analysis of climate change experiments to test whether and how soil moisture interacts with temperature to affect plant phenology. Field-based climate change experiments that warm plots to different levels offer valuable tools to study climate change impacts on plant phenology. Experiments can combine temperature and precipitation treatments to create the “no-analog” climate scenarios forecasted for the future, particularly when they employ active-warming methods, such as forced air heaters, soil warming cables, or infrared heaters (Shaver et al., 2000; Williams et al., 2007; Aronson and McNulty, 2009). Climate change experiments often monitor daily soil moisture, as well as daily air and soil temperature, at the plot-

level, allowing detailed quantification of how microclimate affects plant phenology.

Previous meta-analyses of phenology in climate change experiments have focused primarily on effects of temperature (Wolkovich et al., 2013). We expected that soil moisture may also affect phenology, with drier soils delaying budburst and leafout phenology and advancing flowering and fruiting phenology. We use measured microclimate and phenology data from two databases of climate change experiments: MicroClimate from Climate Change Experiments (MC3E) and Experimental Phenology (ExPhen) to quantify effects of soil moisture and above-ground temperature on plant phenology (bud-burst, leaf-out, flowering, fruiting, senescence; see Materials and Methods). We also use forecasted changes in temperature and soil moisture to investigate how including soil moisture alters expected future shifts in phenology.

MATERIALS AND METHODS

Data— To investigate how soil moisture interacts with temperature to affect phenology, we used two databases that compile data from climate change experiments. Microclimate data came from the MicroClimate from Climate Change Experiments (MC3E) database Ettinger and Wolkovich (2018). Phenology data came from a ExPhen, a new database of phenology from climate change experiments ?.

To create both databases, we first identified published, active-warming field experiments, many of which included precipitation manipulations. We focused on *in situ* active-warming manipulations because recent analyses indicate that active-warming methods are the most controlled and consistent methods available for experimental warming (Kimball, 2005; Kimball et al., 2008; Aronson and McNulty, 2009; Wolkovich et al., 2013). We carried out a full literature review to identify potential active-warming field experiments, following the methods and search terms of Wolkovich et al. (2013) for their Synthesis of Timings Observed in iNcrease Experiments (STONE) database (Wolkovich et al., 2013), but restricting our focus to active-warming experiments. Further, because our goal was to tease out variation in microclimate (including temperature and soil moisture), we focused on warming studies that included both/either multiple levels of warming and/or precipitation treatments. These additional restrictions constrained the list to 11 new studies published after the STONE database, as well as six of the 37 studies in the STONE database. We contacted authors to obtain daily microclimate and phenological data for these 17 studies and received data (or obtained publicly

available data) for 10 of them, as well as datasets from five additional sites offered or suggested to us over the course of our literature review and data analysis. The daily temperature and soil moisture data from these 15 experiments comprise the MC3E database, which is available at KNB (Ettinger and Wolkovich, 2018). The phenology data from the same 15 experiments comprise the ExPhen database of experimental phenology, which is also available at KNB (?).

Analysis— To understand how soil moisture interacts with temperature to affect phenology, we fit models with measured soil moisture, measured temperature, and their interaction to phenology response data (budburst, leafout, flowering, fruiting, senescence). Microclimate data came from the MC3E database, and phenology data came from the ExPhen database. We excluded conifers from the analysis, because their phenology has distinct differences from angiosperm phenology Polgar et al. (2014) and conifer data existed from only one site in the database. For all phenology variables, the response variable was day-of-year of the phenological event. Predictors for our primary models were measured air temperature, soil moisture, and their interaction. We also fit model(s) with soil temperature in place of air temperature. Random effects for all phenology models were species (with random slopes and intercepts), site (random intercept), and year nested within site (random intercept). Equations for these models can be found in the Supplemental Methods. To better understand the interactive effects of measured temperature and soil moisture, we conducted follow-up analyses in which we fit the same phenology models subsets of the data (controls only, low temperature treatments only, and high temperature treatments only).

CUT THIS, RIGHT? To quantify how climate manipulations affect temperature and soil moisture, we used microclimate data from the 4 sites in the MC3E database that manipulated both precipitation and temperature, and measured both above-ground temperature and soil moisture data (exp01,exp05,exp07,exp12). (OR use all 8 studies that measured soil moisture and above-ground temp?) We then fit two groups of hierarchical models to microclimate data from these sites: one group with temperature response variables (including mean annual, and mean seasonal temperatures), and the other group with soil moisture as response variables (including mean annual, and mean seasonal soil moisture). For both groups of models, explanatory variables were temperature treatment, precipitation treatment, and their interaction. The models included a random effect of site, with a random slope and intercept structure, to allow effects of experimental treatments to vary across sites. Models also included random effects of day-of-year, nested within year, on the intercept only, to

account for non-independence of measurements taken on the same day within the same year. Equations for these models can be found in the Supplemental Methods.

RESULTS

How do soil moisture and temperature affect phenology?– We find that soil drying delays phenology and warming temperatures advance phenology, for most phenophases. The magnitude of these effects varies across phenophases, species, and sites.

Effects of soil moisture were strongest for budburst and leafout, and affected all phenophases except fruiting (Figures 1, 1S). Soil drying delays spring budburst at a rate of 0.42 days per percent reduction in soil VWC. Thus, if soil moisture is reduced by 10% of its current state (mean across all sites= XX), as is expected over the next 50 years in the northeastern US (Berg et al., 2017) budburst would be delayed by approximately XX days on average, due to changes in soil moisture alone.

Increasing air temperature advanced phenology for all phenophases except senescence (Figure 1S). Warming advances budburst phenology at a rate of 3.42 days per °C, a rate that is consistent with estimates from previous meta-analyses (Wolkovich et al., 2013).

DISCUSSION

Soil moisture affects phenology, budburst, and leaf-out (not previously a focus of experiments and meta-analyses).

Variable responses to moisture (and precip) may be caused by temporal and spatial variation in the most limiting resource (temperature vs moisture), as well as differences across species

As global warming reduces temperature limitation, importance of moisture limitation in plant phenology may increase.

Conclusions

References

- Aronson, E. L., and S. G. McNulty. 2009. Appropriate experimental ecosystem warming methods by ecosystem, objective, and practicality. *Agricultural and Forest Meteorology* 149:1791–1799.
- Berg, A., J. Sheffield, and P. C. Milly. 2017. Divergent surface and total soil moisture projections under global warming. *Geophysical Research Letters* 44:236–244.
- Cook, B. I., E. M. Wolkovich, T. J. Davies, T. R. Ault, J. L. Betancourt, J. M. Allen, K. Bolmgren, E. E. Cleland, T. M. Crimmins, N. J. B. Kraft, L. T. Lancaster, S. J. Mazer, G. J. McCabe, B. J. McGill, C. Parmesan, S. Pau, J. Regetz, N. Salamin, M. D. Schwartz, and S. E. Travers. 2012. Sensitivity of spring phenology to warming across temporal and spatial climate gradients in two independent databases. *Ecosystems* 15:1283–1294.
- Cui, T., L. Martz, and X. Guo. 2017. Grassland phenology response to drought in the canadian prairies. *Remote Sensing* 9:1258.
- Delpierre, N., E. Dufrêne, K. Soudani, E. Ulrich, S. Cecchini, J. Boé, and C. François. 2009. Modelling interannual and spatial variability of leaf senescence for three deciduous tree species in france. *Agricultural and Forest Meteorology* 149:938–948.
- Essiamah, S., and W. Eschrich. 1986. Water uptake in deciduous trees during winter and the role of conducting tissues in spring reactivation. *IAWA Journal* 7:31–38.
- Ettinger, A., and E. Wolkovich. 2018. Microclimate from climate change experiments (MC3E). doi:10.5063/F1QV3JQR.
- Fujisawa, M., and K. Kobayashi. 2010. Apple (*malus pumila* var. *domestica*) phenology is advancing due to rising air temperature in northern japan. *Global Change Biology* 16:2651–2660.
- Hamann, E., A. E. Weis, and S. J. Franks. 2018. Two decades of evolutionary changes in *brassica rapa* in response to fluctuations in precipitation and severe drought. *Evolution* 72:2682–2696.

- Ibanez, I., R. B. Primack, A. J. Miller-Rushing, E. Ellwood, H. Higuchi, S. D. Lee, H. Kobori, and J. A. Silander. 2010. Forecasting phenology under global warming. *Philosophical Transactions of the Royal Society B-Biological Sciences* 365.
- Kimball, B. 2005. Theory and performance of an infrared heater for ecosystem warming. *Global Change Biology* 11:2041–2056.
- Kimball, B. A., M. M. Conley, S. Wang, X. Lin, C. Luo, J. Morgan, and D. Smith. 2008. Infrared heater arrays for warming ecosystem field plots. *Global Change Biology* 14:309–320.
- Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology Evolution and Systematics* 37:637–669.
- Polgar, C. A., R. B. Primack, J. S. Dukes, C. Schaaf, Z. Wang, and S. S. Hoeppepner. 2014. Tree leaf out response to temperature: comparing field observations, remote sensing, and a warming experiment. *International journal of biometeorology* 58:1251–1257.
- Poloczanska, E. S., C. J. Brown, W. J. Sydeman, W. Kiessling, D. S. Schoeman, P. J. Moore, K. Brander, J. F. Bruno, L. B. Buckley, M. T. Burrows, et al. 2013. Global imprint of climate change on marine life. *Nature Climate Change* 3:919.
- Reich, P. B., and R. Borchert. 1984. Water stress and tree phenology in a tropical dry forest in the lowlands of costa rica. *The Journal of Ecology* pages 61–74.
- Root, T. L., J. T. Price, K. R. Hall, S. H. Schneider, C. Rosenzweig, and J. A. Pounds. 2003. Fingerprints of global warming on wild animals and plants. *Nature* 421:57–60. PT: J.
- Shaver, G. R., J. Canadell, F. S. Chapin, J. Gurevitch, J. Harte, G. Henry, P. Ineson, S. Jonasson, J. Melillo, L. Pitelka, et al. 2000. Global warming and terrestrial ecosystems: A conceptual framework for analysis ecosystem responses to global warming will be complex and varied. Ecosystem warming experiments hold great potential for providing insights on ways terrestrial ecosystems will respond to upcoming decades of climate change. Documentation of initial conditions provides the context for understanding and predicting ecosystem responses. *BioScience* 50:871–882.

- 167 Taylor, G., M. J. Tallis, C. P. Giardina, K. E. Percy, F. Miglietta, P. S. Gupta, B. Gioli, et al. 2008. Future
168 atmospheric CO₂ leads to delayed autumnal senescence. *Global Change Biology* 14:264–275.
- 169 van Schaik, C. P., J. W. Terborgh, and S. J. Wright. 1993. The phenology of tropical forests: adaptive
170 significance and consequences for primary consumers. *Annual Review of ecology and Systematics* 24:353–
171 377.
- 172 Williams, J. W., S. T. Jackson, and J. E. Kutzbacht. 2007. Projected distributions of novel and disappearing
173 climates by 2100 AD. *Proceedings of the National Academy of Sciences of the United States of America*
174 104:5738–5742.
- 175 Wolkovich, E. M., T. J. Davies, H. Schaefer, E. E. Cleland, B. I. Cook, S. E. Travers, C. G. Willis, and C. C.
176 Davis. 2013. Temperature-dependent shifts in phenology contribute to the success of exotic species with
177 climate change. *American Journal of Botany* 100:1407–1421.
- 178 Yu, H., E. Luedeling, and J. Xu. 2010. Winter and spring warming result in delayed spring phenology on the
179 tibetan plateau. *Proceedings of the National Academy of Sciences* 107:22151–22156.

Figures




Figure 1: Model coefficients from budburst, leafout, and flowering models (with centered predictors).