

# Problem Solutions

## Chapter 2

*Pierre Paquay*

### Problem 2.1

Let us begin by extracting the value of  $N$  from the  $\epsilon(M, N, \delta)$  expression. We have that

$$\sqrt{\frac{1}{2N} \ln \frac{2M}{\delta}} \leq \epsilon \Leftrightarrow N \geq \frac{1}{2\epsilon^2} \ln \frac{2M}{\delta}.$$

(a) So for  $M = 1$  and  $\delta = 0.03$ , to have  $\epsilon \leq 0.05$  we need

$$N \geq \frac{1}{2 \cdot 0.05^2} \ln \frac{2}{0.03} = 839.9410156.$$

(b) For  $M = 100$  and  $\delta = 0.03$ , to have  $\epsilon \leq 0.05$  we need

$$N \geq \frac{1}{2 \cdot 0.05^2} \ln \frac{2 \cdot 100}{0.03} = 1760.9750528.$$

(c) And for  $M = 10000$  and  $\delta = 0.03$ , to have  $\epsilon \leq 0.05$  we need

$$N \geq \frac{1}{2 \cdot 0.05^2} \ln \frac{2 \cdot 10000}{0.03} = 2682.00909.$$

### Problem 2.2

For  $N = 4$ , if we consider four non aligned points, this  $\mathcal{H}$  shatters these points (you only have to effectively enumerate them to see that all dichotomies can be generated), so in this case we have  $m_{\mathcal{H}}(4) = 2^4$ .

However, for  $N = 5$ , no matter how you place your five points, some point will be inside a rectangle defined by others. In this case, we are not able to generate all dichotomies and consequently  $m_{\mathcal{H}}(5) < 2^5$ .

From these two observations, we may conclude that, for positive rectangles, we have  $d_{VC} = 4$ , thus

$$m_{\mathcal{H}}(N) \leq N^4 + 1.$$

### Problem 2.3

(a) We already know that the growth function for positive rays is equal to  $N + 1$ . If we enumerate the dichotomies added by negative rays, we get  $N - 1$  new dichotomies (you get the opposite of the ones from positive rays and you have to subtract the two dichotomies where all points are  $+1$  and where all points are  $-1$ ). So in total, we get that

$$m_{\mathcal{H}}(N) = 2N.$$

As the largest value of  $N$  for which  $m_{\mathcal{H}}(N) = 2^N$  is 2 ( $m_{\mathcal{H}}(3) = 6$ ), we have that  $d_{VC} = 2$ .

(b) Here, we already know that the growth function for positive intervals is equal to  $N^2/2 + N/2 + 1$ . If we add the new dichotomies generated by negative intervals, we get  $N - 2$  new ones (for example for  $N = 3$ , we only add the  $(+1, -1, +1)$  dichotomy, and for  $N = 4$ , we add the  $(+1, -1, +1, +1)$  and  $(+1, +1, -1, +1)$

dichotomies). Of course, this only holds if  $N > 1$ , in the case where  $N = 1$  we already generate the two dichotomies with the positive intervals alone. In conclusion, we may write that

$$m_{\mathcal{H}}(N) = \frac{N^2}{2} + \frac{3N}{2} - 1 \text{ if } N > 1 \text{ and } 2 \text{ if } N = 1.$$

As the largest value of  $N$  for which  $m_{\mathcal{H}}(N) = 2^N$  is 3 ( $m_{\mathcal{H}}(4) = 13$ ), we have that  $d_{VC} = 3$ .

(c) To determine the growth function for concentric circles, we have to map the problem from  $\mathbb{R}^d$  to  $[0, +\infty[$ . To do this, we use the map  $\phi$  defined as

$$\phi : (x_1, \dots, x_d) \mapsto r = \sqrt{x_1^2 + \dots + x_d^2}.$$

By doing that, we may see that the problem of concentric circles in  $\mathbb{R}^d$  is equivalent to the problem of positive intervals in  $\mathbb{R}$  (it is easy to see that  $\phi$  maps points with the same radius to a unique point in  $[0, +\infty[$ ), and consequently we may write that

$$m_{\mathcal{H}}(N) = \frac{N^2}{2} + \frac{N}{2} + 1$$

which is independent of  $d$ . As the largest value of  $N$  for which  $m_{\mathcal{H}}(N) = 2^N$  is 2 ( $m_{\mathcal{H}}(3) = 7$ ), we have that  $d_{VC} = 2$ .

## Problem 2.4

We proceed by constructing a specific set of dichotomies for  $N$  points so that among the  $2^N$  possible dichotomies on  $N$  points, we select those that contain at most  $k - 1$  points labelled  $(-1)$ . More precisely, we consider the following dichotomies.

- The dichotomies that contain no  $(-1)$ . We have only  $1 = \binom{N}{0}$  of those.
- The dichotomies that contain a unique  $(-1)$ . We have  $N = \binom{N}{1}$  of those.
- The dichotomies that contain exactly two  $(-1)$ s. We have  $\binom{N}{2}$  of those.
- ...
- The dichotomies that contain exactly  $k - 1$   $(-1)$ s. We have  $\binom{N}{k-1}$  of those.

In total, we have exactly  $\sum_{i=0}^{k-1} \binom{N}{i}$  such dichotomies. Moreover, these dichotomies do not shatter any subset of  $k$  variables because to do that, we would need one dichotomy that contains  $k$   $(-1)$ s, which is not the case in our set. So, we may conclude that

$$B(N, k) \geq \sum_{i=0}^{k-1} \binom{N}{i}$$

and with Sauer's lemma, we get

$$B(N, k) = \sum_{i=0}^{k-1} \binom{N}{i}.$$

## Problem 2.5

To prove the inequality, we begin with the case  $D = 0$ . Here, it is easy to see that

$$1 = \binom{N}{0} \leq N^0 + 1 = 2.$$

Now, we assume the result is correct for  $D$  ( $D \geq 1$ ), and we will prove it for  $D + 1$ . We may write that

$$\begin{aligned}
\sum_{i=0}^{D+1} \binom{N}{i} &= \sum_{i=0}^D \binom{N}{i} + \binom{N}{D+1} \\
&\leq N^D + 1 + \binom{N}{D+1} \\
&\leq N^D + 1 + \frac{N!}{(D+1)!(N-D-1)!}.
\end{aligned}$$

To continue, we have to prove that

$$\frac{N!}{(N-D-1)!} \leq N^{D+1},$$

which is equivalent to

$$\frac{1}{N^{D+1}} \cdot \frac{N!}{(N-D-1)!} \leq 1.$$

To see this, it suffices to note that

$$\frac{1}{N^{D+1}} \cdot \frac{N!}{(N-D-1)!} = \frac{1}{N^{D+1}} \prod_{i=0}^D (N-i) = \prod_{i=0}^D \frac{N-i}{N^{D+1}} \leq 1.$$

So, we are now able to write that

$$\begin{aligned}
\sum_{i=0}^{D+1} \binom{N}{i} &\leq N^D + 1 + \frac{N!}{(D+1)!(N-D-1)!} \\
&\leq N^D + 1 + \frac{N^{D+1}}{(D+1)!}.
\end{aligned}$$

As  $D \geq 1$ , we have  $(D+1)! \geq 2$ , and consequently

$$\frac{1}{(D+1)!} \leq \frac{1}{2},$$

which enables us to write that

$$\begin{aligned}
\sum_{i=0}^{D+1} \binom{N}{i} &\leq N^D + 1 + \frac{N^{D+1}}{(D+1)!} \\
&\leq N^D + 1 + \frac{N^{D+1}}{2}.
\end{aligned}$$

Moreover, as we assumed  $N \geq D+1$  (if not, we trivially have the result, as in this case  $\binom{N}{D+1} = 0$ ), we get  $N \geq 2$  and consequently

$$\frac{1}{N} < \frac{1}{2} \Leftrightarrow \frac{N^D}{N^{D+1}} < \frac{1}{2} \Leftrightarrow N^D < \frac{N^{D+1}}{2},$$

which allows us to get our result as we now have

$$\begin{aligned}
\sum_{i=0}^{D+1} \binom{N}{i} &\leq N^D + 1 + \frac{N^{D+1}}{2} \\
&\leq \frac{N^{D+1}}{2} + 1 + \frac{N^{D+1}}{2} = N^{D+1} + 1.
\end{aligned}$$

## Problem 2.6

As we have  $N \geq d$ , we may write that  $N/d \geq 1$ , and also that  $(N/d)^{d-i} \geq 1$  for  $i = 0, \dots, d$ . Now, we have that

$$\begin{aligned}
 \sum_{i=0}^d \binom{N}{i} &= \sum_{i=0}^d \binom{N}{i} \cdot 1 \\
 &\leq \sum_{i=0}^d \binom{N}{i} \left(\frac{N}{d}\right)^{d-i} \\
 &\leq \left(\frac{N}{d}\right)^d \sum_{i=0}^d \binom{N}{i} \left(\frac{d}{N}\right)^i \\
 &\leq \left(\frac{N}{d}\right)^d \sum_{i=0}^N \binom{N}{i} \left(\frac{d}{N}\right)^i.
 \end{aligned}$$

Moreover, we also have that

$$\begin{aligned}
 \sum_{i=0}^N \binom{N}{i} \left(\frac{d}{N}\right)^i &= \sum_{i=0}^N \binom{N}{i} 1^{N-i} \cdot \left(\frac{d}{N}\right)^i \\
 &= \left(1 + \frac{d}{N}\right)^N \leq e^d.
 \end{aligned}$$

In conclusion, we have proven that

$$\sum_{i=0}^d \binom{N}{i} \leq \left(\frac{N}{d}\right)^d \cdot e^d = \left(\frac{eN}{d}\right)^d.$$

As we already know that

$$m_{\mathcal{H}}(N) \leq \sum_{i=0}^{d_{VC}} \binom{N}{i},$$

we immediately get that

$$m_{\mathcal{H}}(N) \leq \left(\frac{eN}{d_{VC}}\right)^{d_{VC}}$$

for  $N \geq d_{VC}$ .