

News from the frontline: recent insights into PAMP-triggered immunity in plants

Benjamin Schwessinger and Cyril Zipfel

Plants have developed a complex defence network to fight off invading pathogens. In recent years, the full importance of pathogen-associated molecular pattern (PAMP)-triggered immunity (PTI) within this network became apparent. Several new PAMPs have been isolated and new pattern-recognition receptors (PRRs) identified. The discovery of the PRR-interacting protein BAK1 sheds light on the immediate downstream signalling events. Further, transcriptomic analyses identified a core set of ~100 PAMP-responsive genes. These studies also revealed a significant overlap with genes regulated during effector-triggered immunity (ETI). Strikingly, ETI seems to operate by alleviating the negative feedback regulation of PTI, leading to stronger defences. This review discusses recent findings in PTI recognition and signalling, and illustrates the need to discover new regulators of PTI responses for a full understanding of plant innate immunity.

Addresses

The Sainsbury Laboratory, Colney Lane, Norwich NR4 7UH, UK

Corresponding author: Zipfel, Cyril (cyril.zipfel@tsl.ac.uk)

Current Opinion in Plant Biology 2008, **11**:389–395

This review comes from a themed issue on
 Biotic Interactions
 Edited by Murray Grant and Sophien Kamoun

Available online 3rd July 2008

1369-5266/\$ – see front matter
 © 2008 Elsevier Ltd. All rights reserved.

DOI [10.1016/j.pbi.2008.06.001](https://doi.org/10.1016/j.pbi.2008.06.001)

Plants are constantly surrounded and attacked by a plethora of microorganisms including plant pathogens. The first line of active defence relies on the recognition of pathogen-associated (or microbe-associated) molecular patterns (PAMPs/MAMPs) by PRRs leading to PTI [1^{••},2^{••}]. The importance of PTI becomes apparent by enhanced susceptibility of mutants in PTI recognition or signalling components to adapted or non-adapted bacteria. Furthermore, successful pathogens need to suppress PTI using effectors or toxins to achieve their full virulent potential [3[•],4^{••}] (reviews by Alfano, Hogenhout, Birch and Hussey, this issue). PTI suppression together with other effector functions leads to effector-triggered susceptibility (ETS) [1^{••}]. Some plants have evolved to sense specific pathogen effectors by R-proteins leading to ETI. Often ETI is quantitatively stronger than PTI and is

accompanied by a hypersensitive response form of programmed cell death [1^{••}].

In this review we will focus on recent progress in the identification of new molecules activating PTI, new insights into PAMP recognition and downstream signalling. We will also discuss the role of negative feedback regulation and the interconnection of PTI and ETI signalling.

PAMPs/MAMPs/DAMPs: the thing that gives you away

PAMPs/MAMPs are defined as invariant epitopes within molecules that are fundamental to the pathogens' fitness, widely distributed among different microbes, absent in the host and recognised by a wide array of potential hosts.

Filamentous oomycetes and fungi

Pep13 was the first molecule clearly defined as a PAMP. This is a short 13-amino acid peptide of a conserved surface-exposed fragment within a calcium-dependent cell-wall transglutaminase from the oomycete *Phytophthora sojae* that elicits defence responses in *Solanaceae* [5]. Recently, the cellulose-binding elicitor lectin from *Phytophthora parasitica* var. *nicotianae* was shown to trigger defence responses in *Arabidopsis* and tobacco [6]. The two cellulose-binding domains, which are probably important for cell adhesion during infection, are sufficient for defence induction. Additionally, many plant-species-specific elicitors from oomycetes are known such as elicitors and heptagluconosides [5].

Other more broadly recognised fungal PAMPs are derived from cell-wall components, such as β -glucan, ergosterol or chitin [5].

Bacteria

Plants recognise many different bacterial PAMPs. Lipopolysaccharides (LPSs) are perceived by a range of plant species [7]. The main PAMP of LPS is the widely conserved lipid A. However, the core oligosaccharide and O-antigen structure are also recognised, potentially by a different perception system than lipid A in *Arabidopsis* [7]. Peptidoglycans (PGNs) are a major cell-wall component of Gram-positive bacteria. PGN sugar chains longer than the disaccharide, but not the protein moieties, are recognised as a PAMP in *Arabidopsis* [8]. The highly conserved N-terminal domain of flagellin (flg22) is another extracellular PAMP which is recognised by most plant species [9]. Tomato is able to recognise a shorter version of the same epitope (flg15) and rice

seems insensitive to flg22 but can recognise full-length flagellin [10,11*].

In addition, several bacterial pathogens can evade flagellin recognition. For example, *Xanthomonas campestris* pv. *campestris* (*Xcc*) has within-species polymorphism in flg22, which leads to the inability of *Arabidopsis* to recognise the 'disguised' epitope [12*], though the biological relevance of this relative recognition on *Xcc*-adapted hosts needs further investigation. Therefore, PAMPs are more variable than previously anticipated and the recognised sub-molecular epitopes can vary between plant species.

Notably, not all PAMPs are unambiguously part of extracellular entities. The bacterial cold shock protein (CSP) and the translation elongation factor Tu (EF-Tu) are best known as intracellular proteins. Nevertheless, the core 22-amino acid (csp22) of the RNA-binding domain-1 and the *N*-acetylated N-terminal 18 amino acids of EF-Tu (elf18) are recognised by *Solanaceae* and *Brassicaceae*, respectively [9].

Overall the variability of PAMP recognition between different plant species demonstrates a diversity of PRR specificities. This diversity can be potentially exploited by pyramiding PRRs or chimeric PRRs of orthologues with new recognition capacity to overcome local microbial adaptation.

Flagellin, LPS and PGN are also classical PAMPs recognised by other higher eukaryotes; however, the actual recognised epitopes vary greatly between different kingdoms [7–9]. This clearly reflects the outcome of convergent evolution in innate immunity.

Beyond PAMPs

In addition to sensing invading microbes by means of PAMPs (infectious non-self), plants and animals can also sense infectious-self or modified-self via danger-associated molecular patterns (DAMPs) [13]. For example, plants can recognise oligo- α -galacturonides released from plant cell walls by fungal hydrolytic enzymes [14]. Furthermore, Nep1 (necrosis and ethylene-inducing peptide 1)-like proteins secreted by many pathogens are recognised by plants [15*]. This recognition results from their phytotoxic modulation of the host, and can therefore be referred to as toxin-mediated immunity. Also, recognition of the PAMP-inducible endogenous AtPep peptides was proposed to amplify PTI via a positive feedback loop, and can be seen as part of the infectious-self [16*].

Update on PAMP recognition

In recent years, the first plant PRRs have been cloned. Most identified PRRs are receptor-like kinases (RLKs) or receptor-like proteins (RLPs), except for the extracellular glucan-binding protein (GBP) that binds and hydrolyses heptagluconide from *P. sojae* [17*]. The tomato xylanase

receptors LeEIX1/2 are RLPs. Intriguingly, both GBP and LeEIX1/2 lack obvious intracellular signalling domains. Therefore, other membrane-associated proteins are probably required for signal relay.

Until recently only two *Arabidopsis* RLKs involved in PAMP perception were known; the leucine-rich repeat (LRR)-RLKs FLS2 and EFR that recognise bacterial flagellin and EF-Tu, respectively [18**,19*]. Interestingly, the *Arabidopsis* LysM-RLK CERK1/LysM-RLK1 was recently shown to be required for chitin responses [20*,11*]. However, it is not known if CERK1/LysM-RLK1 is required for chitin binding, and therefore if it is a PRR *per se*.

Notably, in rice the LysM-containing transmembrane protein CEBiP is required for chitin binding and responses [21*]. As CEBiP lacks intracellular-signalling domains, it might interact with the rice CERK1/LysM-RLK1 orthologue to transduce the chitin signal. Conversely, CEBiP orthologues might be required for chitin binding in *Arabidopsis*.

FLS2 and BAK1: update on flagellin recognition and immediate downstream signalling

FLS2 is the best studied plant PRR. AtFLS2 orthologues have been recently identified in *Nicotiana benthamiana* and tomato, based on sequence homology and requirement for flg22 perception [22*,23**]. In tomato the minimal recognised flagellin epitope is smaller and the recognised epitope more diverse than in *Arabidopsis*, demonstrating the evolutionary adaptation and flexibility of PAMP recognition [22*]. In the AtFLS2 extracellular domain the LRRs 9–15 were identified as contributing significantly to flg22 binding [24*]. However, the specific flg22-binding site could not be identified and the authors proposed that a limited number of amino acids quantitatively contributed to flg22 binding.

Flg22 binding leads to AtFLS2 endocytosis from the plasma membrane into intracellular mobile vesicles [25]. The internalisation relies on kinase activity and requires the ubiquitination-related PEST motif in the AtFLS2 C-terminus [25]. AtFLS2 endocytosis is not mediated by homodimerisation as AtFLS2 does not form homodimers in protoplasts [26]. However, flg22-mediated endocytosis of FLS2 was not observed in the protoplast system [26], which raises the question of protoplasts as a suitable system for these studies. Nevertheless, flg22 binding reduces AtFLS2 membrane mobility most probably through heterodimerisation [26].

The LRR-RLK BAK1/SERK3 rapidly forms complexes with AtFLS2 upon flg22 binding and positively regulates flg22 responses [27**,28**]. BAK1 also interacts with and phosphorylates the LRR-RLK brassinosteroid hormone

receptor BRI1 [29]. Additionally, defence responses triggered by *elf18* in *Arabidopsis*, or *osp22* and the *Phytophthora infestans* elicitor INF1 in *N. benthamiana* are also partially compromised in plants defective in *BAK1* expression [27^{••},30^{••}]. However, *BAK1* is not involved in flg22 binding [27^{••}]. Therefore *BAK1* does not act as a co-receptor but rather as a signal transducer most probably relying on its kinase activity.

Interestingly, *bak1 Arabidopsis* mutants are not more susceptible to bacteria [30^{••}]. By contrast, *N. benthamiana* silenced for *NbBAK1* are more susceptible to bacterial pathogens [28^{••}]. The discrepancy in the requirement of *BAK1* for bacterial resistance could be explained by silencing of close *BAK1* paralogues in *N. benthamiana*, which in *Arabidopsis* could partially substitute for *BAK1* loss. The best candidate is *BKK1*, a *BAK1* paralogue that acts redundantly with *BAK1* in cell death control and brassinosteroid signalling in *Arabidopsis* [30^{••},31^{••}]. This cell death control probably contributes to the hypersusceptibility of *bak1* mutants to necrotrophic pathogens [30^{••}].

Overall, *BAK1* appears to be a general regulator of LRR-RLKs, with newly defined roles in PTI signalling and cell death control that are independent of brassinosteroid signalling. It is probable that *BAK1* interacts with and regulates the activity of yet other unknown RLKs.

PTI responses, signalling and regulation: many read-outs, little genetic proof

PAMP perception triggers many different traceable molecular, physiological and pathogenesis-related changes. Seconds to minutes after PAMP treatment there is ion-flux across the plasma membrane, increased intracellular Ca^{2+} concentration, oxidative burst, MAP kinase (MAPK) activation, protein phosphorylation, receptor endocytosis and protein–protein interaction, for example [5,32[•]]. By 30 min transcriptional changes are induced. They comprise up to 3% of the *Arabidopsis* genome and significantly overlap with the ETI transcriptome [18^{••},33]. *Elf18* and flg22 induce the same set of genes, and ~30% or ~50% of those are also regulated by PGN or chitin, respectively [8,18^{••},11[•]]. This highly significant overlap in transcriptional regulation by different PAMPs clearly suggests signal convergence after PAMP recognition. Furthermore, comparison of *Arabidopsis* whole-genome transcriptional changes following infection with non-adapted, adapted and non-virulent bacteria identified over 800 PAMP-regulated genes [34,35]. Ninety-six were up-regulated over a prolonged time period of 12 hours post-infection by *Pseudomonas syringae* pv. *tomato* DC3000 mutants defective in their type-III secretion system, as well as by four different PAMPs (flg22, *elf18*, LPS and HrpZ). Thus, these genes could be the core primary PTI response [35]. Recently, expression profiling experiments using mini-arrays designed to

analyse pathogen-responsive genes revealed the partial dependence of PTI-mediated gene induction on SA-signalling [36]. PAMPs are able to increase SA levels and mutations in the SA-signalling network abate PTI-induced defence [37,38].

Other responses that occur within hours are the enhancement of ethylene biosynthesis, stomatal closure and callose deposition [32[•]]. Despite the plethora of biological read-outs the signalling, regulation and interrelationship between these events and their genetic basis are only emerging slowly.

MAP kinases and WRKYs: positive and negative regulation of PTI


Different members of the MAPK and WRKY transcription factor families are components of downstream PTI signalling. The MAPK cascade AtMEKK1, AtMKK4/5 and AtMPK3/6 leading to the activation of WRKY22/29 was previously proposed as a positive regulatory module of flg22 signalling using constitutively active kinases in protoplasts [39]. However, recent genetic studies concluded that AtMEKK1 does not regulate AtMPK3/6 activity but rather acts upstream of the negative defence regulator AtMPK4 [40–42]. Furthermore, the positive defence regulator AtMKK1 activates both AtMPK3/6 and AtMPK4 [43]. It is therefore apparent that PTI not only up-regulates defence responses but also initiates its own negative feedback regulation (Figure 1). Together this demonstrates that interconnected MAPK networks regulate PTI signalling. Additionally, the specificity of individual MAPKs is questionable, as MPK3/6 is also involved in stomatal patterning and jasmonic acid signalling, for example [44,45].

Let us take on the negative regulators

Plants have to strike a balance between fast and efficient defence activation and penalising inappropriate expression of defence genes and uncontrolled cell death [46]. In animal innate immunity it is of paramount importance to control the strength and duration of innate immune responses to prevent over-reaction of the immune system (e.g. sepsis) and autoimmune diseases [47]. There has been much recent progress in identifying negative regulators of mammal innate immunity. These range from TAM (Tyro3/Axl/Mer) family members and Toll-like receptors, inactive signalling mimics, phosphatases, ubiquitin lyases and ligases, prolyl isomerases, transcription factors, to micro-RNAs [48,49,50[•],51^{••},52]. Negative regulators can either be constitutively expressed or transcriptionally induced, and/or additionally regulated post-translationally, leading to a regulatory network of interconnected positive and negative feedback loops [47,51^{••}].

Several lines of evidence suggest a similar regulatory network exists in plant defence. Firstly, treatment with the protein synthesis inhibitor cycloheximide elicits

Figure 1



Conceptual model of plant innate immunity. (a) Hypothetical constitutive negative regulation of PTI. Plants tightly regulate their defence responses in the absence of microbes by constitutive negative regulators. (b) Activation of PTI by recognition of infectious-self and non-self. Plants recognise invading microbes (e.g. bacteria) by sensing PAMPs/MAMPs or DAMPs via PRRs. This triggers diminution of the negative regulation and consequently activation of downstream signalling cascades leading to PTI. This activation is in turn dampened and/or enhanced by negative and/or positive feedback loops, respectively. (c) Suppression of PTI by pathogen effectors. Pathogens can suppress PTI via effector molecules. This includes binding to and modification of PRRs and signalling components. Additionally, effectors could mimic or stabilise negative regulators. (d) Activation of ETI by recognition of pathogen effectors. Some plants recognise individual effector molecules leading to ETI. This could be partially achieved by sequestering PTI negative regulators and possibly by reinforcing PTI-induced transcriptional changes. For definitions and identities of molecular players involved, please refer to main text.

similar gene expression to untreated seedlings following flg22 application [53], suggesting the presence of unstable negative regulators, which are rapidly degraded after defence elicitation. Secondly, three E3-ligases involved in cell death control are rapidly induced after flg22 treatment and ETI [54,55]. These are primary candidates to degrade constitutively present negative regulators. Thirdly, it became a paradigm that plant hormones de-repress signalling pathways that are normally constitutively suppressed [56].

Our knowledge about negative regulation of PTI is limited but it is highly probable that post-translational modification of host proteins are required to suppress PTI (see Block *et al.*, this issue). The kinase-associated protein phosphatase binds to AtFLS2 and abrogates downstream signalling [32^{*}]. AtNUDT7, a nucleoside diphosphate hydrolase, negatively regulates basal defence via an unknown mechanism [57,58]. An expression profile study with four *Pseudomonas* strains carrying different effectors triggering ETI supports the idea of the importance of ADP-ribosylation in plant defence. Furthermore,

NUDT7 also negatively regulates ETI, strengthening the notion of interconnected PTI and ETI signalling networks [59]. Interestingly, the *Pseudomonas* effector HopU1 ADP-ribosylates the glycine-rich RNA-binding protein GPR7 most probably interfering with RNA binding and disturbing the amount of PTI-related transcripts [60^{**}]. Plants expressing *HopU1* or mutated in *GRP7* are indeed more susceptible to *Pseudomonas* infection.

RIN4, an interactor of the R-proteins RPM1 and RPT2, was also suggested to be a negative regulator of PTI. However, it is still an unsolved question how this proposed function can be reconciled with its degradation by the effector AvrRpt2 [61]. The *Arabidopsis* homologue of the human transcription factor NF-X1 negatively regulates toxin-mediated immunity induced by trichothecene [62]. Further, several WRKY transcription factors, such as WRKY11/17 and WRKY18/40/60 are negative regulators of basal defence and with the exception of *WRKY60* are all transcriptionally up-regulated after PAMP treatment [63^{*},64^{*},65^{**},66]. Intriguingly, the nuclear localisation of the activated MLA10 R-protein

leads to physical interaction with the barley orthologue of *AtWRKY18/40/60* [65**]. Therefore it is conceivable that the quantitative difference between PTI and ETI is mediated by de-repressing negative feedback regulation of PTI. This is at least partially achieved by sequestration of negative regulators of PTI (Figure 1). Thus effectors might also mimic or stabilise negative regulators to suppress PTI.

Conclusion

There has been much recent progress in understanding plant defences, and PTI is a central research focus. However, not insignificant challenges lie ahead of us. We have to identify key negative and positive PTI regulators to further elucidate the complex signalling network underlying the different layers of plant innate immunity. Furthermore, the role of small RNAs in PTI signalling and the role of PTI in symbiosis are interesting and valuable areas to investigate. Finally we have to demonstrate the transferability of our work on plant innate immunity to generate broad-spectrum resistance in crops.

Acknowledgements

We apologise to colleagues who could not be cited because of strict space limitations. Our research is supported by the Gatsby Charitable Foundation. We would like to thank John Rathjen for comments on the manuscript and stimulating discussions.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Jones JD, Dangl JL: **The plant immune system.** *Nature* 2006, **444**:323-329.
Two excellent reviews providing a comprehensive overview of plant immunity. The authors introduce the concepts of PTI and ETI, and integrate them in the so-called 'zig-zag' model that underlies the arms-race occurring between plants and their pathogens.
2. Chisholm ST, Coaker G, Day B, Staskawicz BJ: **Host-microbe interactions: shaping the evolution of the plant immune response.** *Cell* 2006, **124**:803-814.
Two excellent reviews providing a comprehensive overview of plant immunity. The authors introduce the concepts of PTI and ETI, and integrate them in the so-called 'zig-zag' model that underlies the arms-race occurring between plants and their pathogens.
3. Bray Speth E, Lee YN, He SY: **Pathogen virulence factors as molecular probes of basic plant cellular functions.** *Curr Opin Plant Biol* 2007, **10**:580-586.
A comprehensive review of effector functions within plant cells. It introduces the interesting idea of using effectors as molecular probes to investigate basic cellular functions.
4. Xiang T, Zong N, Zou Y, Wu Y, Zhang J, Xing W, Li Y, Tang X, Zhu L, Chai J *et al.*: ***Pseudomonas syringae* effector AvrPto blocks innate immunity by targeting receptor kinases.** *Curr Biol* 2008, **18**:74-80.
The authors showed that the PRRs FLS2 and EFR are directly targeted by the bacterial effector AvrPto. The physical interactions of AvrPto and the PRRs require similar, but not identical, motifs in AvrPto to those involved in binding to the tomato resistance protein Pto. This physical interaction leads to the blockage of PTI and enhances susceptibility of the host plant.
5. Nurnberger T, Brunner F, Kemmerling B, Piater L: **Innate immunity in plants and animals: striking similarities and obvious differences.** *Immunol Rev* 2004, **198**:249-266.

6. Gaulin E, Drame N, Lafitte C, Torto-Alalibo T, Martinez Y, Ameline-Torregrosa C, Khatib M, Mazarguil H, Villalba-Mateos F, Kamoun S *et al.*: **Cellulose binding domains of a *Phytophthora* cell wall protein are novel pathogen-associated molecular patterns.** *Plant Cell* 2006, **18**:1766-1777.
7. Newman M-A, Dow JM, Molinaro A, Parrilli M: **Invited review: priming, induction and modulation of plant defence responses by bacterial lipopolysaccharides.** *J Endotoxin Res* 2007, **13**:69-84.
8. Gust AA, Biswas R, Lenz HD, Rauhut T, Ranf S, Kemmerling B, Gotz F, Glawischnig E, Lee J, Felix G *et al.*: **Bacteria-derived peptidoglycans constitute pathogen-associated molecular patterns triggering innate immunity in *Arabidopsis*.** *J Biol Chem* 2007, **282**:32338-32348.
9. Zipfel C, Felix G: **Plants and animals: a different taste for microbes?** *Curr Opin Plant Biol* 2005, **8**:353-360.
10. Takai R, Kaneda T, Isogai A, Takayama S, Che FS: **A new method of defense response analysis using a transient expression system in rice protoplasts.** *Biosci Biotechnol Biochem* 2007, **71**:590-593.
11. Wan J, Zhang XC, Neece D, Ramonell KM, Clough S, Kim SY, Stacey MG, Stacey G: **A LysM receptor-like kinase plays a critical role in chitin signaling and fungal resistance in *Arabidopsis*.** *Plant Cell* 2008, **20**:471-481.
Here CERK1/LysM-RLK1 is identified as a positive regulator of chitin signalling in *Arabidopsis*. *Arabidopsis* CERK1/LysM-RLK1 mutants are blocked in chitin-induced transcriptional reprogramming, which resembles transcriptional changes induced by other PAMPs. These mutants are also more susceptible to fungal pathogen. Furthermore, pre-treatment of plants with chitin induces resistance to fungal and bacterial pathogens.
12. Sun W, Dunning FM, Pfund C, Weingarten R, Bent AF: **Within-species flagellin polymorphism in *Xanthomonas campestris* pv. *campestris* and its impact on elicitation of *Arabidopsis* FLAGELLIN SENSING2-dependent defenses.** *Plant Cell* 2006, **18**:764-779.
Sequence analysis of the flg22 epitope in different *Xanthomonas campestris* pv. *campestris* strains combined with elicitor activity studies showed that flg22 is more variable than previously anticipated.
13. Matzinger P: **Friendly and dangerous signals: is the tissue in control?** *Nat Immunol* 2007, **8**:11-13.
14. Ferrari S, Galletti R, Denoux C, De Lorenzo G, Ausubel FM, Dewdney J: **Resistance to *Botrytis cinerea* induced in *Arabidopsis* by elicitors is independent of salicylic acid, ethylene, or jasmonate signaling but requires PHYTOALEXIN DEFICIENT3.** *Plant Physiol* 2007, **144**:367-379.
15. Qutob D, Kemmerling B, Brunner F, Kufner I, Engelhardt S, Gust AA, Luberacki B, Seitz HU, Stahl D, Rauhut T *et al.*: **Phytotoxicity and innate immune responses induced by Nep1-like proteins.** *Plant Cell* 2006, **18**:3721-3744.
The phytotoxin Nep1-like proteins of oomycetes trigger defence responses resembling those induced by flg22. This observation supports the idea of DAMP recognition and toxin-mediated immunity.
16. Ryan CA, Huffaker A, Yamaguchi Y: **New insights into innate immunity in *Arabidopsis*.** *Cell Microbiol* 2007, **9**:1902-1908.
This review summarises recent findings that recognition of the PAMP-induced endogenous peptides AtPep1 and its paralogues by the LRR-RLK PEPR1 activates plant defence and could provide a positive feedback loop mechanism to amplify PAMP-induced responses.
17. Zipfel C: **Pattern-recognition receptors in plant innate immunity.** *Curr Opin Immunol* 2008, **20**:10-16.
A recent review providing a detailed overview on known plant PRRs.
18. Zipfel C, Kunze G, Chinchilla D, Caniard A, Jones JD, Boller T, Felix G: **Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts *Agrobacterium*-mediated transformation.** *Cell* 2006, **125**:749-760.
This paper reports the identification and characterisation of the LRR-RLK EFR as the EF-Tu receptor. Comparison of transcription profiles after elf18 or flg22 treatment showed a substantial overlap and led the authors to suggest signalling convergence downstream of individual PAMP recognition. It also showed that transformation of plant cells by the bacterium *Agrobacterium tumefaciens* is normally restricted by plant defences.

19. Chinchilla D, Bauer Z, Regenass M, Boller T, Felix G: **The *Arabidopsis* receptor kinase FLS2 binds flg22 and determines the specificity of flagellin perception.** *Plant Cell* 2006, **18**:465-476.
The authors demonstrated that FLS2 is the flg22-binding site and provided recognition specificity between different flg22-derived peptides, proving that FLS2 is the flg22 PRR *per se*.
20. Miya A, Albert P, Shinya T, Desaki Y, Ichimura K, Shirasu K, Narusaka Y, Kawakami N, Kaku H, Shibuya N: **CERK1, a LysM receptor kinase, is essential for chitin elicitor signaling in *Arabidopsis*.** *Proc Natl Acad Sci U S A* 2007, **104**:19613-19618.
Adopting a reverse genetic approach studying LysM RLKs in *Arabidopsis* the authors identified the receptor-like kinase CERK1/LysM-RLK1 as an essential signalling component for chitin responses in *Arabidopsis*. Furthermore, they showed that CERK1/LysM-RLK1 localises to the plasma membrane and possesses *de novo* kinase activity.
21. Kaku H, Nishizawa Y, Ishii-Minami N, Akimoto-Tomiyama C, Dohmae N, Takio K, Minami E, Shibuya N: **Plant cells recognize chitin fragments for defense signaling through a plasma membrane receptor.** *Proc Natl Acad Sci U S A* 2006, **103**:11086-11091.
In an exhaustive biochemical study the LysM containing receptor-like protein CEBiP was identified as the chitin-binding PRR in rice. Plants silenced for CEBiP has reduced chitin binding and responsiveness. This is the first and only identified PRR in monocots.
22. Robatzek S, Bittel P, Chinchilla D, Kochner P, Felix G, Shi SH, Boller T: **Molecular identification and characterization of the tomato flagellin receptor LeFLS2, an orthologue of *Arabidopsis* FLS2 exhibiting characteristically different perception specificities.** *Plant Mol Biol* 2007, **64**:539-547.
Tomato is a species extremely sensitive to flagellin and with distinct recognition specificities compared to *Arabidopsis*. This paper reports the cloning of the tomato LeFLS2 based on sequence similarity with AtFLS2, complementation of *Arabidopsis fls2* mutant and binding to assays.
23. Hann DR, Rathjen JP: **Early events in the pathogenicity of *Pseudomonas syringae* on *Nicotiana benthamiana*.** *Plant J* 2007, **49**:607-618.
The authors identified the FLS2 orthologue in *N. benthamiana* and showed that NbFLS2 silencing enhances susceptibility to a range of adapted and non-adapted bacteria. Furthermore, the two bacterial effectors AvrPto and AvrPtoB suppress PTI when transiently expressed in *N. benthamiana* leading to enhanced bacterial susceptibility.
24. Dunning FM, Sun W, Jansen KL, Helft L, Bent AF: **Identification and mutational analysis of *Arabidopsis* FLS2 leucine-rich repeat domain residues that contribute to flagellin perception.** *Plant Cell* 2007, **19**:3297-3313.
Combining mutational, phylogenetic and structural modelling approaches, the authors showed that the AtFLS2 LRR domains 9-15 contributed significantly to flg22 binding.
25. Robatzek S, Chinchilla D, Boller T: **Ligand-induced endocytosis of the pattern recognition receptor FLS2 in *Arabidopsis*.** *Genes Dev* 2006, **20**:537-542.
26. Ali GS, Prasad KV, Day I, Reddy AS: **Ligand-dependent reduction in the membrane mobility of FLAGELLIN SENSITIVE2, an *Arabidopsis* receptor-like kinase.** *Plant Cell Physiol* 2007, **48**:1601-1611.
27. Chinchilla D, Zipfel C, Robatzek S, Kemmerling B, Nurnberger T, Jones JD, Felix G, Boller T: **A flagellin-induced complex of the receptor FLS2 and BAK1 initiates plant defence.** *Nature* 2007, **448**:497-500.
In a reverse-genetic approach focussed on LRR-RLK genes rapidly induced by flg22, the authors found that plants mutated in BAK1, a gene involved in brassinosteroid signalling, are strongly affected in flg22 and partially in elf18 responses. Although, BAK1 is dispensable for flg22 binding, BAK1 and FLS2 rapidly interact in a ligand-dependent manner.
28. Heese A, Hann DR, Gimenez-Ibanez S, Jones AM, He K, Li J, Schroeder JI, Peck SC, Rathjen JP: **The receptor-like kinase SERK3/BAK1 is a central regulator of innate immunity in plants.** *Proc Natl Acad Sci U S A* 2007, **104**:12217-12222.
Using a proteomic approach, the authors also identified BAK1 as a FLS2 interactor and positive regulator of flg22 signalling. Interestingly, silencing of NbBAK1 in *Nicotiana benthamiana* decreases responsiveness not only to flg22 but also to the PAMPs CSP22 and INF1. NbBAK1-silenced plants are also more susceptible to a range of bacteria and a biotrophic oomycete.
29. Gendron JM, Wang Z-Y: **Multiple mechanisms modulate brassinosteroid signaling.** *Curr Opin Plant Biol* 2007, **10**:436-441.
30. Kemmerling B, Schwedt A, Rodriguez P, Mazzotta S, Frank M, Qamar SA, Mengiste T, Betsuyaku S, Parker JE, Mussig C *et al.*: **The BRI1-associated kinase 1, BAK1, has a brassinolide-independent role in plant cell-death control.** *Curr Biol* 2007, **17**:1116-1122.
The authors showed that BAK1 is a negative regulator of cell death. BAK1 mutants have enhanced runaway cell death upon bacterial infection with Pto DC3000 without changes in the disease susceptibility. However, this loss of cell death control enhances susceptibility to necrotrophic fungi. Similar to the above studies, the effects of BAK1 are brassinosteroid-independent, revealing that BAK1 controls signalling triggered by several different LRR-RLKs.
31. He K, Gou X, Yuan T, Lin H, Asami T, Yoshida S, Russell SD, Li J: **BAK1 and BKK1 regulate brassinosteroid-dependent growth and brassinosteroid-independent cell-death pathways.** *Curr Biol* 2007, **17**:1109-1115.
This paper showed that BKK1, the closest homologue of BAK1 in the SERK family, has a redundant role in cell death and brassinosteroid signalling.
32. Altenbach D, Robatzek S: **Pattern recognition receptors: from the cell surface to intracellular dynamics.** *Mol Plant Microbe Interact* 2007, **20**:1031-1039.
A comprehensive review on PAMP-triggered immunity.
33. Navarro L, Zipfel C, Rowland O, Keller I, Robatzek S, Boller T, Jones JD: **The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis.** *Plant Physiol* 2004, **135**:1113-1128.
34. Thilmony R, Underwood W, He SY: **Genome-wide transcriptional analysis of the *Arabidopsis thaliana* interaction with the plant pathogen *Pseudomonas syringae* pv. tomato DC3000 and the human pathogen *Escherichia coli* O157:H7.** *Plant J* 2006, **46**:34-53.
35. Truman W, Zabala MT, Grant M: **Type III effectors orchestrate a complex interplay between transcriptional networks to modify basal defence responses during pathogenesis and resistance.** *Plant J* 2006, **46**:14-33.
36. Sato M, Mitra RM, Collier J, Wang D, Spivey NW, Dewdney J, Denoux C, Glazebrook J, Katagiri F: **A high-performance, small-scale microarray for expression profiling of many samples in *Arabidopsis*-pathogen studies.** *Plant J* 2007, **49**:565-577.
37. Mishina TE, Zeier J: **Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in *Arabidopsis*.** *Plant J* 2007, **50**:500-513.
38. Tsuda K, Sato M, Glazebrook J, Cohen JD, Katagiri F: **Interplay between MAMP-triggered and SA-mediated defense responses.** *Plant J* 2008, **53**:763-775.
39. Asai T, Tena G, Plotnikova J, Willmann MR, Chiu WL, Gomez-Gomez L, Boller T, Ausubel FM, Sheen J: **MAP kinase signalling cascade in *Arabidopsis* innate immunity.** *Nature* 2002, **415**:977-983.
40. Ichimura K, Casais C, Peck SC, Shinozaki K, Shirasu K: **MEKK1 is required for MPK4 activation and regulates tissue-specific and temperature-dependent cell death in *Arabidopsis*.** *J Biol Chem* 2006, **281**:36969-36976.
41. Nakagami H, Soukupova H, Schikora A, Zarsky V, Hirt H: **A mitogen-activated protein kinase kinase kinase mediates reactive oxygen species homeostasis in *Arabidopsis*.** *J Biol Chem* 2006, **281**:38697-38704.
42. Suarez-Rodriguez MC, Adams-Phillips L, Liu Y, Wang H, Su SH, Jester PJ, Zhang S, Bent AF, Krysan PJ: **MEKK1 is required for flg22-induced MPK4 activation in *Arabidopsis* plants.** *Plant Physiol* 2007, **143**:661-669.
43. Meszaros T, Helfer A, Hatzimasoura E, Magyar Z, Serazetdinova L, Rios G, Bardoczky V, Teige M, Koncz C, Peck S *et al.*: **The *Arabidopsis* MAP kinase kinase MKK1 participates in defence responses to the bacterial elicitor flagellin.** *Plant J* 2006, **48**:485-498.

44. Takahashi F, Yoshida R, Ichimura K, Mizoguchi T, Seo S, Yonezawa M, Maruyama K, Yamaguchi-Shinozaki K, Shinozaki K: **The mitogen-activated protein kinase cascade MKK3-MPK6 is an important part of the jasmonate signal transduction pathway in *Arabidopsis*.** *Plant Cell* 2007, **19**:805-818.
45. Wang H, Ngwenyama N, Liu Y, Walker JC, Zhang S: **Stomatal development and patterning are regulated by environmentally responsive mitogen-activated protein kinases in *Arabidopsis*.** *Plant Cell* 2007, **19**:63-73.
46. Hofius D, Tsitsigiannis DI, Jones JDG, Mundy J: **Inducible cell death in plant immunity.** *Semin Cancer Biol* 2007, **17**:166-187.
47. Liew FY, Xu D, Brint EK, O'Neill LAJ: **Negative regulation of Toll-like receptor-mediated immune responses.** *Nat Rev Immunol* 2005, **5**:446-458.
48. Saitoh T, Tun-Kyi A, Ryo A, Yamamoto M, Finn G, Fujita T, Akira S, Yamamoto N, Lu KP, Yamaoka S: **Negative regulation of interferon-regulatory factor 3-dependent innate antiviral response by the prolyl isomerase Pin1.** *Nat Immunol* 2006, **7**:598-605.
49. Taganov KD, Boldin MP, Baltimore D: **MicroRNAs and immunity: tiny players in a big field.** *Immunity* 2007, **26**:133-137.
50. Carmody RJ, Ruan Q, Palmer S, Hilliard B, Chen YH: **Negative regulation of Toll-like receptor signaling by NF- κ B p50 ubiquitination blockade.** *Science* 2007, **317**:675-678.
This paper showed that B-cell leukemia 3 (Bcl-3) negatively regulates animal innate immunity. Bcl3 blocks the ubiquitination and degradation of the transcriptional repressor p50, a member of the NF- κ B family.
51. Rothlin CV, Ghosh S, Zuniga EI, Oldstone MBA, Lemke G: **TAM receptors are pleiotropic inhibitors of the innate immune response.** *Cell* 2007, **131**:1124-1136.
The authors identified TAM receptors as pleiotropic negative regulators of animal innate immunity. This negative regulation requires new gene expression and is mediated by the positive regulator of the cytokine feed-forward loop STAT1. Interestingly this secondary negative feedback loop mediated by TAM receptors also requires STAT1. A wonderful study.
52. Whitmore MM, Iparraguirre A, Kubelka L, Weninger W, Hai T, Williams BRG: **Negative regulation of TLR-signaling pathways by activating transcription factor-3.** *J Immunol* 2007, **179**:3622-3630.
53. Navarro L, Zipfel C, Rowland O, Keller I, Robatzek S, Boller T, Jones JDG: **The transcriptional innate immune response to flg22. Interplay and overlap with Avr gene-dependent defense responses and bacterial pathogenesis.** *Plant Physiol* 2004, **135**:1113-1128.
54. Yang C-W, Gonzalez-Lamothe R, Ewan RA, Rowland O, Yoshioka H, Shenton M, Ye H, O'Donnell E, Jones JDG, Sadanandom A: **The E3 ubiquitin ligase activity of *Arabidopsis* PLANT U-BOX17 and its functional tobacco homolog ACRE276 are required for cell death and defense.** *Plant Cell* 2006, **18**:1084-1098.
55. Gonzalez-Lamothe R, Tsitsigiannis DI, Ludwig AA, Panicot M, Shirasu K, Jones JDG: **The U-Box protein CMPG1 is required for efficient activation of defense mechanisms triggered by multiple resistance genes in tobacco and tomato.** *Plant Cell* 2006, **18**:1067-1083.
56. Chow B, McCourt P: **Plant hormone receptors: perception is everything.** *Genes Dev* 2006, **20**:1998-2008.
57. Bartsch M, Gobbato E, Bednarek P, Debey S, Schultze JL, Bautor J, Parker JE: **Salicylic acid-independent ENHANCED DISEASE SUSCEPTIBILITY1 signaling in *Arabidopsis* immunity and cell death is regulated by the monooxygenase FMO1 and the Nudix hydrolase NUDT7.** *Plant Cell* 2006, **18**:1038-1051.
58. Ge X, Li GJ, Wang SB, Zhu H, Zhu T, Wang X, Xia Y: **AtNUDT7, a negative regulator of basal immunity in *Arabidopsis*, modulates two distinct defense response pathways and is involved in maintaining redox homeostasis.** *Plant Physiol* 2007, **145**:204-215.
59. Adams-Phillips L, Wan J, Tan X, Dunning FM, Meyers BC, Michelmore RW, Bent AF: **Discovery of ADP-ribosylation and other plant defense pathway elements through expression profiling of four different *Arabidopsis*-*Pseudomonas* R-avr interactions.** *Mol Plant Microbe Interact* 2008, **21**:646-657.
60. Fu ZQ, Guo M, Jeong BR, Tian F, Elthon TE, Cerny RL, Staiger D, Alfano JR: **A type III effector ADP-ribosylates RNA-binding proteins and quells plant immunity.** *Nature* 2007, **447**:284-288.
The authors set out to elucidate the function of HopU1 an effector with similarity to mono-ADP-ribosyltransferases. This enzymatic activity is important for suppressing PTI host defences in *Arabidopsis* and *N. benthamiana*. Furthermore, the glycine-rich RNA-binding protein GRP7 was identified as *in vivo* target and as being ADP-ribosylated in its RNA-binding domain. *Gpr7* mutant plants show reduced PAMP responsiveness and basal resistance to *Pseudomonas*.
61. Kim MG, da Cunha L, McFall AJ, Belkhadir Y, DebRoy S, Dangl JL, Mackey D: **Two *Pseudomonas syringae* type III effectors inhibit RIN4-regulated basal defense in *Arabidopsis*.** *Cell* 2005, **121**:749-759.
62. Asano T, Masuda D, Yasuda M, Nakashita H, Kudo T, Kimura M, Yamaguchi K, Nishiuchi T: **AtNFXL1, an *Arabidopsis* homologue of the human transcription factor NF-X1, functions as a negative regulator of the trichothecene phytotoxin-induced defense response.** *Plant J* 2008, **53**:450-464.
63. Journot-Catalino N, Somssich IE, Roby D, Kroj T: **The transcription factors WRKY11 and WRKY17 act as negative regulators of basal resistance in *Arabidopsis thaliana*.** *Plant Cell* 2006, **18**:3289-3302.
The authors employed a reverse-genetic approach based on the transcriptional up-regulation of WRKY family members during the infection with avirulent *Pto* DC3000 bacteria to identify WRKY11 and 17 as negative regulators of basal defence with partially redundant roles.
64. Xu X, Chen C, Fan B, Chen Z: **Physical and functional interactions between pathogen-induced *Arabidopsis* WRKY18, WRKY40, and WRKY60 transcription factors.** *Plant Cell* 2006, **18**:1310-1326.
The authors showed that WRKY18/40/60 form homo-dimeric and hetero-dimeric complexes *in planta* and that the composition of these complexes significantly influences the DNA binding capacity. Furthermore, *wrky18/40*, *wrky18/60* and *wrky18/40/60* plants have enhanced basal resistance to the bacterial pathogen *Pto* DC3000, but enhanced susceptibility to the necrotrophic fungal pathogen *Botrytis cinerea*. This correlated with their regulation of SA-induced and JA-induced genes.
65. Shen Q-H, Saijo Y, Mauch S, Biskup C, Bieri S, Keller B, Seki H, Ulker B, Somssich IE, Schulze-Lefert P: **Nuclear activity of MLA immune receptors links isolate-specific and basal disease-resistance responses.** *Science* 2007, **315**:1098-1103.
The barley R-protein MLA10 requires nuclear localisation for full function. After effector recognition, MLA10 physically interacts with the barley orthologues of the *Arabidopsis* WRKY18/40/60 in the nucleus. These WRKY transcription factors were shown to be negative regulators of basal defence (see Ref. [64]). This study suggests that ETI could function by de-repressing the negative feedback loop regulation of PTI.
66. Zipfel C, Robatzek S, Navarro L, Oakeley EJ, Jones JD, Felix G, Boller T: **Bacterial disease resistance in *Arabidopsis* through flagellin perception.** *Nature* 2004, **428**:764-767.