



QR Code for
report verification

Case ID : 2600136977	Sample Type : EDTA PERIPHERAL BLOOD
Name : MR. JABEDUL ISLAM (KOL2605010084)	Date & Time Collected : 01-May-2026 12:58 PM
Sex/Age : Male/20 Years	Date & Time Received : 06-May-2026 03:55 PM
Bill. Loc. : OncQuest Laboratories Ltd., New Delhi	Date & Time Reported : 18-May-2026 06:55 PM
Ref. By :	Report Version : 1
Indication :	

Heamseq Comprehensive Leukemia Panel On Illumina Novaseq 6000 Platform

Clinical Indication:

Acute leukemia

Positive

(Clinically significant CNV and Fusion detected)

Key Findings:

Tier	Gene & Transcript	Variant	Exon	Coverage/VAF
-Negative-				

Fusion Genes	Breakpoints	%Total Reads
BCR::ABL1 t(9;22)(q34;q11)	chr22:23,182,239 -chr9:130,854,064	41.73%

CNV Type	Chromosome	Copy Number	Significance
Deletion	Monosomy7 (-7) (Loss)	1	Epidemiology: -7 is the most frequent abnormality in secondary myeloid disorders, found in 51% of the cases in a series of 246 cases, while del(7q) was found in 7%, and a partial monosomy 7 as a result of an unbalanced translocation in 8% of cases; in contrast, -7/del(7q) is found in 10% of de novo myeloid disorders; the sex ratio is 1.5 male for 1 female; the proportion of adults with a -7 myeloid disorder grows dramatically after 60 years Clinics: characterized by infectious susceptibility, quick aggravation, and treatment resistance Prognosis: monosomy 7 is classified as a poor prognostic criterium by the International Prognostic Scoring System; the actuarial relapse rate at one year is 82 %, and the 7-yr actuarial event-free survival is 6 %; after an allogeneic bone marrow transplantation, -7 is predictive of an unfavorable outcome

Page 1 of 93

Dr. Nirmal A. Vaniawala
MD (Path. & Bact.)

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist



MC-7414



QR Code for
report verification

Case ID	: 2600136977	Sample Type	: EDTA PERIPHERAL BLOOD
Name	: MR. JABEDUL ISLAM (KOL2605010084)	Date & Time Collected	: 01-May-2026 12:58 PM
Sex/Age	: Male/20 Years	Date & Time Received	: 06-May-2026 03:55 PM
Bill. Loc.	: OncQuest Laboratories Ltd., New Delhi	Date & Time Reported	: 18-May-2026 06:55 PM
Ref. By	:	Report Version	: 1
Indication	:		

Test Description

HaemSeq Comprehensive Leukemia Panel is advancing precision hematological malignancies research through the analysis of multiple relevant biomarkers in a single next-generation sequencing (NGS) test.

The HaemSeq Comprehensive Leukemia Panel assay analyses **513** unique genes (**262** DNA + **251** RNA) for single gene and multiple gene biomarker insights, detects single gene biomarkers from all variant types including **SNVs, indels, CNVs, known and novel fusions, and splice variants**, analyze complex multi-gene biomarkers for mutational signatures and Detects **ITD** variants in **FLT3** gene.

The performance of the panel was verified using both commercially sourced reference controls and PB & BM samples.

Cytogenetics and CNV Analysis: Karyotyping analysis for the identification of complex variants in co-relation to molecular analysis of Small/Large Deletion and Duplication.

Test Methodology

Extraction & Library Preparation: DNA & RNA extracted from PB/BM sample is used for targeted gene capture using a custom designed hybrid capture panel. In process quality controls were determined for prepared library.

Sequencing: The libraries were sequenced at mean depth: >1000x on Illumina Novaseq- next generation sequencing platform.

Data Analysis: Sequencing data are analyzed using a customized and validated bioinformatics pipeline capable of detecting various classes of genomic alterations, including single nucleotide variants (**SNVs**), **Indels**, copy number variations (**CNVs**) and **fusions**. As part of the streamlined bioinformatics workflow. Reportable genomic alterations and fusions are prioritized, classified, and reported based on AMP-ASCO-CAP guidelines and NCCN guidelines.

Limit of Detection (LOD): The LOD for SNVs and InDels is 5% Variant allele Frequency (VAF) and for Fusions is ≥ 10 supporting reads.

Notes

Treatment decisions should not be based solely on the results of this test. Findings must be interpreted in the context of the patient's complete clinical profile—including pathological, radiological, and family history—to guide diagnosis, prognosis, and therapeutic decisions.

Therapy-related information in this report is derived from multiple sources including FDA-approved drug data, NCCN guidelines, peer-reviewed literature, standard clinical databases, and biomarker evidence strength. However, therapy options listed may not be suitable or beneficial for all patients. This report aims to provide a summary of potentially

Page 2 of 93

Dr. Nirmal A. Vaniawala
MD (Path. & Bact.)

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist



MC-7414



QR Code for
report verification

Case ID : 2600136977	Sample Type : EDTA PERIPHERAL BLOOD
Name : MR. JABEDUL ISLAM (KOL2605010084)	Date & Time Collected : 01-May-2026 12:58 PM
Sex/Age : Male/20 Years	Date & Time Received : 06-May-2026 03:55 PM
Bill. Loc. : OncQuest Laboratories Ltd., New Delhi	Date & Time Reported : 18-May-2026 06:55 PM
Ref. By :	Report Version : 1
Indication :	

effective treatments, therapies that may pose increased risks, and those that may have limited benefit, based on genomic alterations identified. Prognostic and diagnostic biomarkers relevant to the disease context may also be reported.

SN Genelab Pvt Ltd does not guarantee completeness, accuracy, or eligibility for trial enrollment. Physicians must verify eligibility criteria for specific trials independently.

Identification of a genomic alteration does not guarantee therapeutic efficacy or predict ineffectiveness. Final decisions on treatment should rest with the treating physician. The absence of reported therapeutic options does not imply ineffectiveness or lack of safety of any medication selected independently.

All variant classifications and related clinical information are based on the latest evidence from peer-reviewed publications, public clinical databases, and medical guidelines (e.g., NCCN, ASCO, AMP) available at the time of report generation. However, literature is continuously evolving, and classification may change with emerging evidence.

The assay is designed and validated for somatic variant detection, it does not differentiate between somatic and germline variants. If indicated, germline testing and genetic counselling are recommended.

Poor-quality DNA or RNA may fail QC thresholds at various stages such as extraction, library preparation, or bioinformatics, and may result in assay failure or compromised results, such as low gene coverage or shallow variant depth.

Variants with allele frequencies between 3–5% may be reported if they are of strong or potential clinical relevance, or if previously detected in the same patient. False positives or negatives below assay sensitivity cannot be ruled out.

CNVs/gene amplifications have >80% detection sensitivity in this assay. Confirmation using orthogonal methods such as FISH/DDPCR/MLPA is recommended when clinically indicated.

Variants in non-coding regions or deep intronic sequences are not assessed in this assay.

This assay is under the scope of NABL as Heamseq comprehensive leukemia panel.

DNA Gene List

ABL1	ACD	AKT1	ALK	ANKRD26
ARAF	ARID1A	ARID1B	ARID2	ASXL1
ASXL2	ATM	ATRX	B2M	BCL10
BCL11A	BCL2	BCL6	BCL7A	BCOR

Page 3 of 93

Dr. Nirmal A. Vaniawala
MD (Path. & Bact.)

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist



MC-7414



QR Code for
report verification


Case ID : 2600136977
Name : **MR. JABEDUL ISLAM (KOL2605010084)**
Sex/Age : Male/20 Years
Bill. Loc. : OncQuest Laboratories Ltd., New Delhi
Ref. By :
Indication :


Sample Type : EDTA PERIPHERAL BLOOD
Date & Time Collected : 01-May-2026 12:58 PM
Date & Time Received : 06-May-2026 03:55 PM
Date & Time Reported : 18-May-2026 06:55 PM
Report Version : 1

BCORL1	BIRC2	BIRC3	BLM	BRAF
BTG1	BTG2	BTk	CALR	CARD11
CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CCR6	CD22	CD274
CD28	CD58	CD70	CD79A	CD79B
CD83	CDK4	CDK6	CDKN1B	CDKN2A
CDKN2B	CDKN2C	CEBPA	CHEK2	CIITA
CKS1B	CNOT3	CRBN	CREBBP	CSF1R
CSF3R	CTCF	CTNNB1	CUX1	CXCR4
CYLD	DDX3X	DDX41	DIS3	DNMT3A
DTX1	EBF1	EGFR	EGR1	EGR2
ELANE	EP300	ERBB2	ERBB3	ETNK1
ETV6	EZH2	FAS	FAT1	FAT3
FAT4	FBXO11	FBXW7	FGFR1	FGFR2
FGFR3	FLT3	FOXO1	GATA1	GATA2
GATA3	GNA13	GNAS	GPR34	GTF2I
HIST1H1B	HIST1H1C	HIST1H1D	HIST1H1E	HIST1H2AC
HIST1H2AG	HIST1H2AM	HIST1H2BC	HIST1H2BD	HIST1H2BG
HIST1H2BK	HIST1H3B	HIST1H3G	HLA-A	HLA-B
HLA-C	HRAS	ID3	IDH1	IDH2
IGH	IKBKB	IKZF1	IKZF3	IL2RG
IL7R	IRF1	IRF4	IRF8	ITPKB
JAK1	JAK2	JAK3	KDM6A	KIK3R1
KIT	KLF2	KLHL6	KMT2A	KMT2C
KMT2D	KRAS	LCK	LMO2	LTB
MAP2K1	MAP2K4	MAP3K1	MCL1	MECOM
MED12	MEF2B	MET	MGA	MPEG1
MPL	MSH2	MSH6	MTOR	MYB
MYC	MYD88	NF1	NF2	NFE2
NFKB1	NFKB2	NFKBIA	NFKBIE	NFKBIZ
NOTCH1	NOTCH2	NPM1	NRAS	NSD2
NTRK1	P2RY8	PAX5	PDCD1LG2	PDGFRA
PDGFRB	PDS5B	PHF6	PIGA	PIK3CA
PIK3CB	PIK3CD	PIM1	PLCG2	POT1
POU2AF1	POU2F2	PPM1D	PRDM1	PRKCB
PRPF8	PTCH1	PTEN	PTPN11	PTPN2
PTPN6	PTPRC	PTPRD	RAD21	RAD51
RASA2	RB1	RCOR1	REL	RHOA
ROS1	RPS15	RUNX1	SAMD9	SAMD9L
SETBP1	SETD2	SETDB1	SF1	SF3B1

Page 4 of 93


Dr. Nirmal A. Vaniawala
MD (Path. & Bact.)


Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist


Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist





QR Code for
report verification


Case ID : 2600136977	Sample Type : EDTA PERIPHERAL BLOOD
Name : MR. JABEDUL ISLAM (KOL2605010084)	Date & Time Collected : 01-May-2026 12:58 PM
Sex/Age : Male/20 Years	Date & Time Received : 06-May-2026 03:55 PM
Bill. Loc. : OncQuest Laboratories Ltd., New Delhi	Date & Time Reported : 18-May-2026 06:55 PM
Ref. By :	Report Version : 1
Indication :	


SGK1	SH2B3	SMARCA4	SMARCB1	SMC1A
SMC3	SOCS1	SPEN	SPIB	SRC
SRSF2	STAG2	STAT3	STAT5B	STAT6
STK11	SUZ12	SYK	TBL1XR1	TCF3
TENT5C	TERC	TERT	TET2	THRAP3
TLR2	TMEM30A	TMSB4X	TNF	TNFAIP2
TNFAIP3	TNFRSF14	TP53	TRAF3	U2AF1
U2AF2	UBR5	VAV1	WT1	XPO1
ZFP36L1	ZRSR2			

RNA Gene list				
ABI1	ABI2	ABL1	ABL2	ACSL6
ACTN4	ADGRG7	AFDN	AFF1	AFF3
AFF4	AGGF1	ALK	ANLN	ARHGAP26
ARHGEF12	ASNS	ATF7IP	ATIC	ATXN2L
AUTS2	BAIAP2L1	BCL11A	BCL11B	BCL2
BCL6	BCL9	BCOR	BCR	BIRC3
BRAF	BTBD18	CBFA2T3	CBFB	CBL
CCDC6	CCDC88C	CCND1	CCND3	CD28
CD74	CDK6	CEP170B	CEP43	CHD1
CHIC2	CIITA	CIP2A	CLTC	CNTRL
CREBBP	CRLF2	CSF1R	CT45A2	CTLA4
DAB2IP	DCTN1	DDX10	DDX3X	DEK
DHH	DIAPH2	DUSP22	EBF1	EGFR
EIF4A1	ELL	ELN	EML1	EML4
EP300	EPOR	EPS15	ERG	ESRRB
ETV6	EWSR1	FGFR1	FGFR3	FIP1L1
FLNA	FLT3	FNBP1	FOXO3	FOXO4
FOXP1	FRYL	FUS	GAS7	GATA1
GLIS2	GOT1	GPHN	GTF2H3	HAUS1
HLF	HNRNPUL1	HOXA11	HOXA13	HOXA9
HOXC11	HOXC13	HOXD13	IGLL5	IKZF2
IKZF3	INPP5D	IRF2BP2	ITK	JAK2
KANK1	KAT6A	KDM5A	KIF5B	KLF2
KMT2A	KMT2E	KNL1	LASP1	LMNA
LNP1	LPP	LYN	MALT1	MAML2
MAPK13	MAPRE1	MECOM	MEF2D	MET
MLF1	MLLT1	MLLT10	MLLT11	MLLT3
MLLT6	MN1	MNX1	MRTFA	MSN
MYB	MYC	MYH11	MYO1F	NCKIPSD

Page 5 of 93


Dr. Nirmal A. Vaniawala
MD (Path. & Bact.)


Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist


Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist



MC-7414



QR Code for
report verification

Case ID : 2600136977
Name : MR. JABEDUL ISLAM (KOL2605010084)
Sex/Age : Male/20 Years
Bill. Loc. : OncQuest Laboratories Ltd., New Delhi
Ref. By :
Indication :
Sample Type : EDTA PERIPHERAL BLOOD
Date & Time Collected : 01-May-2026 12:58 PM
Date & Time Received : 06-May-2026 03:55 PM
Date & Time Reported : 18-May-2026 06:55 PM
Report Version : 1

NCOA2	NCOA4	NDE1	NF1	NFKB2
NOTCH1	NPM1	NSD1	NSD3	NTRK1
NTRK3	NUMA1	NUP214	NUP98	P2RY8
PAG1	PAN3	PAX5	PAX8	PBX1
PCM1	PDCD1LG2	PDGFRA	PDGFRB	PDS5A
PHF23	PICALM	PIM1	PLEKHA6	PML
PPARG	PRDM16	PRKG2	PRRX1	PSIP1
PSMA2	PTK2B	PTPRR	RANBP2	RAP1GDS1
RARA	RBM15	RCSD1	REL	RET
RHEBL1	ROS1	RPL22	RPN1	RUNX1
RUNX1T1	S100A7	SALL2	SARNP	SEC31A
SEMA6A	SEPTIN14	SEPTIN2	SEPTIN5	SEPTIN6
SEPTIN9	SET	SETD2	SFPQ	SH3GL1
SLC34A2	SLC45A3	SORBS2	SPI1	SQSTM1
SSBP2	STAT5B	STIL	STMN1	SYK
TACC3	TAF15	TAL1	TBL1XR1	TCF3
TCF7	TET1	TFG	THAP4	TMPRSS2
TNIP1	TOMM40	TOP1	TOP3A	TP63
TPM3	TPM4	TRIP11	TYK2	UBA2
USP42	USP9X	VAV1	WTIP	ZBTB16
ZCCHC7	ZFYVE19	ZMIZ1	ZMYM2	ZNF384
ZNF521				

Tier	Interpretation
Tier IA	Variants of Strong Clinical Significance: Biomarkers that predict response or resistance to US FDA-approved therapies for a specific type of tumor or have been included in professional guidelines as therapeutic, diagnostic, and/or prognostic biomarkers for specific types of tumors.
Tier IB	Variants of Strong Clinical Significance: Biomarkers that predict response or resistance to a therapy based on well-powered studies with consensus from experts in the field, or have diagnostic and/or prognostic significance of certain diseases based on well-powered studies with expert consensus.
Tier IIC	Variants of Potential Clinical Significance: biomarkers that predict response or resistance to therapies approved by FDA or professional societies for a different tumor type (i.e., off-label use of a drug), serve as inclusion criteria for clinical trials, or have diagnostic and/or prognostic significance based on the results of multiple small studies.
Tier IID	Variants of Potential Clinical Significance: Biomarkers that show plausible therapeutic significance based on preclinical studies or may assist disease diagnosis and/or prognosis themselves or along with other biomarkers based on small studies or multiple case reports with no consensus.

Page 6 of 93

Nirmal A. Vaniawala

Dr.Nirmal A. Vaniawala
MD (Path. & Bact.)

Salil Vaniawala

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist

Salil Vaniawala

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist



MC-7414



QR Code for
report verification

Case ID	: 2600136977	Sample Type	: EDTA PERIPHERAL BLOOD
Name	: MR. JABEDUL ISLAM (KOL2605010084)	Date & Time Collected	: 01-May-2026 12:58 PM
Sex/Age	: Male/20 Years	Date & Time Received	: 06-May-2026 03:55 PM
Bill. Loc.	: OncQuest Laboratories Ltd., New Delhi	Date & Time Reported	: 18-May-2026 06:55 PM
Ref. By	:	Report Version	: 1
Indication	:		

Tier III	Variants of Unknown clinical significance (VUS): Not observed at a significant allele frequency in the general or specific subpopulation or pancancer or tumor specific variant databases. No convincing published evidence of cancer Association
-----------------	---

----- End Of Report -----

Dr.Nirmal A. Vaniawala
MD (Path. & Bact.)

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics) Consulting
Geneticist

Dr. Salil Vaniawala
M.Sc Ph.D.(Human Genetics)
Consulting Geneticist



MC-7414



Sample Cancer Type: Acute Lymphoblastic Leukemia

Table of Contents	Page	Report Highlights
Relevant Therapy Summary	3	1 Relevant Biomarkers
Relevant Therapy Details	12	50 Therapies Available
Prognostic Details	79	61 Clinical Trials
Diagnostic Details	81	
Clinical Trials Summary	82	
Alert Details	86	

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	BCR::ABL1 fusion	dasatinib ^{1, 2} dasatinib + chemotherapy ^{1, 2} imatinib ^{† 1, 2} imatinib [†] + chemotherapy ^{1, 2} ponatinib ^{1, 2} ponatinib + chemotherapy ¹ allogeneic stem cells asciminib + chemotherapy blinatumomab bosutinib bosutinib + chemotherapy bosutinib + inotuzumab ozogamicin dasatinib + inotuzumab ozogamicin imatinib + inotuzumab ozogamicin inotuzumab ozogamicin inotuzumab ozogamicin + nilotinib inotuzumab ozogamicin + ponatinib nilotinib nilotinib + chemotherapy tisagenlecleucel-t brexucabtagene autoleucel dasatinib + inotuzumab ozogamicin + chemotherapy dasatinib + tisagenlecleucel-t dasatinib + venetoclax + chemotherapy imatinib + inotuzumab ozogamicin + chemotherapy imatinib + tisagenlecleucel-t imatinib + venetoclax + chemotherapy inotuzumab ozogamicin + ponatinib + chemotherapy	bosutinib ^{1, 2} dasatinib ^{1, 2} imatinib ^{† 1, 2} nilotinib ^{1, 2} asciminib ^{1, 2} ponatinib allogeneic stem cells azacitidine cladribine + cytarabine + daunorubicin cladribine + cytarabine + mitoxantrone + filgrastim cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + daunorubicin + fludarabine cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine gemtuzumab ozogamicin glasdegib + chemotherapy liposomal cytarabine-daunorubicin CPX-351 venetoclax + chemotherapy asciminib + chemotherapy bosutinib + chemotherapy dasatinib + chemotherapy imatinib + chemotherapy	61

* Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO
* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO
† Includes biosimilars/generics
Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

For queries please call on 9586177755

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
		obecabtagene autoleucel ponatinib + tisagenlecleucel-t ponatinib + venetoclax + chemotherapy venetoclax + chemotherapy	nilotinib + chemotherapy ponatinib + chemotherapy	
	Prognostic significance: NCCN: Unfavorable to Standard Diagnostic significance: Acute Lymphoblastic Leukemia			

* Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO
* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO
† Includes biosimilars/generics
Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

Relevant Therapy Summary

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
dasatinib					
imatinib					
ponatinib					(I/II)
dasatinib + chemotherapy					
imatinib + chemotherapy					
ponatinib + chemotherapy					
bosutinib					
nilotinib					
asciminib					
dasatinib + blinatumomab					
ponatinib + blinatumomab					
inotuzumab ozogamicin					(II)
obecabtagene autoleucel					(I)
blinatumomab					
bosutinib + blinatumomab					
bosutinib + CALGB 10701 regimen					
bosutinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone					
bosutinib + cyclophosphamide + daunorubicin + vincristine + prednisone					
bosutinib + cytarabine + HyperCVAD + methotrexate					
bosutinib + cytarabine + methotrexate + methylprednisolone					
bosutinib + EsPhALL regimen					
bosutinib + EWALL regimen					
bosutinib + inotuzumab ozogamicin					
bosutinib + mercaptopurine + methotrexate + vincristine + prednisone					
bosutinib + steroid					
bosutinib + vincristine + dexamethasone					

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
bosutinib + vincristine + prednisone	×	●	×	×	×
brexucabtagene autoleucel	×	●	×	×	×
dasatinib + ALL-REZ BFM 90 regimen	×	●	×	×	×
dasatinib + calaspargase pegol + cytarabine	×	●	×	×	×
dasatinib + CALGB 10701 regimen	×	●	×	×	×
dasatinib + clofarabine + cyclophosphamide + etoposide	×	●	×	×	×
dasatinib + COG AALL01P2 regimen	×	●	×	×	×
dasatinib + COG AALL0622 regimen	×	●	×	×	×
dasatinib + COG AALL0622 regimen + blinatumomab	×	●	×	×	×
dasatinib + COG AALL07P1 regimen	×	●	×	×	×
dasatinib + COG AALL1131 regimen + blinatumomab	×	●	×	×	×
dasatinib + COG AALL1331 regimen	×	●	×	×	×
dasatinib + COG AALL1631 regimen	×	●	×	×	×
dasatinib + COG AALL1631 regimen + blinatumomab	×	●	×	×	×
dasatinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone	×	●	×	×	×
dasatinib + cyclophosphamide + daunorubicin + vincristine + prednisone	×	●	×	×	×
dasatinib + cytarabine + fludarabine + filgrastim	×	●	×	×	×
dasatinib + cytarabine + HyperCVAD + methotrexate	×	●	×	×	×
dasatinib + cytarabine + methotrexate + methylprednisolone	×	●	×	×	×
dasatinib + cytarabine + PEG-L-asparaginase	×	●	×	×	×
dasatinib + DFCI-ALL Protocol 16-001 regimen	×	●	×	×	×
dasatinib + DFCI-ALL Protocol 16-001 regimen + blinatumomab	×	●	×	×	×
dasatinib + EsPhALL regimen	×	●	×	×	×
dasatinib + EWALL regimen	×	●	×	×	×
dasatinib + FLAG-IDA	×	●	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
dasatinib + inotuzumab ozogamicin	×	●	×	×	×
dasatinib + inotuzumab ozogamicin + cyclophosphamide + vincristine + dexamethasone	×	●	×	×	×
dasatinib + mercaptopurine + methotrexate + vincristine + prednisone	×	●	×	×	×
dasatinib + steroid	×	●	×	×	×
dasatinib + tisagenlecleucel-t	×	●	×	×	×
dasatinib + Total Therapy XVII regimen	×	●	×	×	×
dasatinib + Total Therapy XVII regimen + blinatumomab	×	●	×	×	×
dasatinib + UKALL R3 regimen	×	●	×	×	×
dasatinib + venetoclax + calaspargase pegol + vincristine + dexamethasone	×	●	×	×	×
dasatinib + venetoclax + calaspargase pegol + vincristine + prednisone	×	●	×	×	×
dasatinib + venetoclax + PEG-L-asparaginase + vincristine + dexamethasone	×	●	×	×	×
dasatinib + venetoclax + PEG-L-asparaginase + vincristine + prednisone	×	●	×	×	×
dasatinib + vincristine + dexamethasone	×	●	×	×	×
dasatinib + vincristine + prednisone	×	●	×	×	×
imatinib + ALL-REZ BFM 90 regimen	×	●	×	×	×
imatinib + blinatumomab	×	●	×	×	×
imatinib + calaspargase pegol + cytarabine	×	●	×	×	×
imatinib + CALGB 10701 regimen	×	●	×	×	×
imatinib + clofarabine + cyclophosphamide + etoposide	×	●	×	×	×
imatinib + COG AALL01P2 regimen	×	●	×	×	×
imatinib + COG AALL07P1 regimen	×	●	×	×	×
imatinib + COG AALL1331 regimen	×	●	×	×	×
imatinib + COG AALL1631 regimen	×	●	×	×	×
imatinib + COG AALL1631 regimen + blinatumomab	×	●	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
imatinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone	×	●	×	×	×
imatinib + cyclophosphamide + daunorubicin + vincristine + prednisone	×	●	×	×	×
imatinib + cytarabine + fludarabine + filgrastim	×	●	×	×	×
imatinib + cytarabine + HyperCVAD + methotrexate	×	●	×	×	×
imatinib + cytarabine + methotrexate + methylprednisolone	×	●	×	×	×
imatinib + cytarabine + PEG-L-asparaginase	×	●	×	×	×
imatinib + EsPhALL regimen	×	●	×	×	×
imatinib + EWALL regimen	×	●	×	×	×
imatinib + FLAG-IDA	×	●	×	×	×
imatinib + inotuzumab ozogamicin	×	●	×	×	×
imatinib + inotuzumab ozogamicin + cyclophosphamide + vincristine + dexamethasone	×	●	×	×	×
imatinib + mercaptopurine + methotrexate + vincristine + prednisone	×	●	×	×	×
imatinib + steroid	×	●	×	×	×
imatinib + tisagenlecleucel-t	×	●	×	×	×
imatinib + UKALL R3 regimen	×	●	×	×	×
imatinib + venetoclax + calaspargase pegol + vincristine + dexamethasone	×	●	×	×	×
imatinib + venetoclax + calaspargase pegol + vincristine + prednisone	×	●	×	×	×
imatinib + venetoclax + PEG-L-asparaginase + vincristine + dexamethasone	×	●	×	×	×
imatinib + venetoclax + PEG-L-asparaginase + vincristine + prednisone	×	●	×	×	×
imatinib + vincristine + dexamethasone	×	●	×	×	×
imatinib + vincristine + prednisone	×	●	×	×	×
inotuzumab ozogamicin + nilotinib	×	●	×	×	×
inotuzumab ozogamicin + ponatinib	×	●	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
inotuzumab ozogamicin + ponatinib + cyclophosphamide + vincristine + dexamethasone	×	●	×	×	×
nilotinib + blinatumomab	×	●	×	×	×
nilotinib + CALGB 10701 regimen	×	●	×	×	×
nilotinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone	×	●	×	×	×
nilotinib + cyclophosphamide + daunorubicin + vincristine + prednisone	×	●	×	×	×
nilotinib + cytarabine + HyperCVAD + methotrexate	×	●	×	×	×
nilotinib + cytarabine + methotrexate + methylprednisolone	×	●	×	×	×
nilotinib + EsPhALL regimen	×	●	×	×	×
nilotinib + EWALL regimen	×	●	×	×	×
nilotinib + mercaptopurine + methotrexate + vincristine + prednisone	×	●	×	×	×
nilotinib + steroid	×	●	×	×	×
nilotinib + vincristine + dexamethasone	×	●	×	×	×
nilotinib + vincristine + prednisone	×	●	×	×	×
ponatinib + ALL-REZ BFM 90 regimen	×	●	×	×	×
ponatinib + calaspargase pegol + cytarabine	×	●	×	×	×
ponatinib + CALGB 10701 regimen	×	●	×	×	×
ponatinib + clofarabine + cyclophosphamide + etoposide	×	●	×	×	×
ponatinib + COG AALL01P2 regimen	×	●	×	×	×
ponatinib + COG AALL07P1 regimen	×	●	×	×	×
ponatinib + COG AALL1331 regimen	×	●	×	×	×
ponatinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone	×	●	×	×	×
ponatinib + cyclophosphamide + daunorubicin + vincristine + prednisone	×	●	×	×	×
ponatinib + cytarabine + fludarabine + filgrastim	×	●	×	×	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
ponatinib + cytarabine + HyperCVAD + methotrexate	×	●	×	×	×
ponatinib + cytarabine + methotrexate + methylprednisolone	×	●	×	×	×
ponatinib + cytarabine + PEG-L-asparaginase	×	●	×	×	×
ponatinib + EsPhALL regimen	×	●	×	×	×
ponatinib + EWALL regimen	×	●	×	×	×
ponatinib + FLAG-IDA	×	●	×	×	×
ponatinib + mercaptopurine + methotrexate + vincristine + prednisone	×	●	×	×	×
ponatinib + steroid	×	●	×	×	×
ponatinib + tisagenlecleucel-t	×	●	×	×	×
ponatinib + UKALL R3 regimen	×	●	×	×	×
ponatinib + venetoclax + calaspargase pegol + vincristine + dexamethasone	×	●	×	×	×
ponatinib + venetoclax + calaspargase pegol + vincristine + prednisone	×	●	×	×	×
ponatinib + venetoclax + PEG-L-asparaginase + vincristine + dexamethasone	×	●	×	×	×
ponatinib + venetoclax + PEG-L-asparaginase + vincristine + prednisone	×	●	×	×	×
ponatinib + vincristine + dexamethasone	×	●	×	×	×
ponatinib + vincristine + prednisone	×	●	×	×	×
tisagenlecleucel-t	×	●	×	×	×
venetoclax + calaspargase pegol + vincristine + dexamethasone	×	●	×	×	×
venetoclax + calaspargase pegol + vincristine + prednisolone	×	●	×	×	×
venetoclax + PEG-L-asparaginase + vincristine + dexamethasone	×	●	×	×	×
venetoclax + PEG-L-asparaginase + vincristine + prednisone	×	●	×	×	×
allogeneic hematopoietic stem cell transplantation	×	○	×	●	×
asciminib + chemotherapy	×	○	×	●	×

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
bosutinib + chemotherapy	×	○	×	●	×
nilotinib + chemotherapy	×	○	×	●	×
cytarabine + daunorubicin	×	○	×	○	×
azacitidine	×	○	×	×	×
cladribine + cytarabine + mitoxantrone + filgrastim	×	○	×	×	×
cytarabine	×	○	×	×	×
cytarabine + daunorubicin + etoposide	×	○	×	×	×
cytarabine + etoposide + idarubicin	×	○	×	×	×
cytarabine + fludarabine + idarubicin + filgrastim	×	○	×	×	×
cytarabine + idarubicin	×	○	×	×	×
cytarabine + mitoxantrone	×	○	×	×	×
decitabine	×	○	×	×	×
gemtuzumab ozogamicin	×	○	×	×	×
glasdegib + cytarabine	×	○	×	×	×
liposomal cytarabine-daunorubicin CPX-351	×	○	×	×	×
venetoclax + azacitidine	×	○	×	×	×
venetoclax + cladribine + cytarabine	×	○	×	×	×
venetoclax + cytarabine	×	○	×	×	×
venetoclax + decitabine	×	○	×	×	×
imatinib (Accord)	×	×	●	×	×
imatinib (Accord) + chemotherapy	×	×	●	×	×
cladribine + cytarabine + daunorubicin	×	×	×	○	×
cytarabine + daunorubicin + fludarabine	×	×	×	○	×
chemotherapy, nilotinib	×	×	×	×	● (III)
imatinib, dasatinib	×	×	×	×	● (III)
olverembatinib, imatinib, chemotherapy	×	×	×	×	● (III)
ponatinib, dasatinib, chemotherapy, blinatumomab, steroid	×	×	×	×	● (III)
allogeneic stem cells, chemotherapy, radiation therapy	×	×	×	×	● (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
allopurinol, chemotherapy, sirolimus, radiation therapy, mycophenolate mofetil	×	×	×	×	● (II)
blinatumomab, asciminib	×	×	×	×	● (II)
blinatumomab, ponatinib, chemotherapy	×	×	×	×	● (II)
CART-CD19	×	×	×	×	● (II)
CART-CD19B	×	×	×	×	● (II)
CART-CD7	×	×	×	×	● (II)
CLIC-1901	×	×	×	×	● (II)
flumatinib, allogeneic stem cells	×	×	×	×	● (II)
flumatinib, chemotherapy	×	×	×	×	● (II)
inotuzumab ozogamicin, chemotherapy	×	×	×	×	● (II)
inotuzumab ozogamicin, dasatinib, ponatinib, steroid, chemotherapy, allogeneic stem cells	×	×	×	×	● (II)
natural killer cell therapy, allogeneic stem cells	×	×	×	×	● (II)
olverembatinib, chemotherapy, steroid	×	×	×	×	● (II)
olverembatinib, venetoclax, chemotherapy	×	×	×	×	● (II)
olverembatinib, venetoclax, steroid	×	×	×	×	● (II)
venetoclax, chemotherapy, olverembatinib	×	×	×	×	● (II)
α/β CD3+ T-cell and CD19+ B-cells	×	×	×	×	● (II)
CART-CD19, chemotherapy, aldesleukin	×	×	×	×	● (I/II)
CART-CD19, rimiducid, chemotherapy	×	×	×	×	● (I/II)
CART-CD19/CD22	×	×	×	×	● (I/II)
CART-CD19/CD22, chemotherapy	×	×	×	×	● (I/II)
CN-201	×	×	×	×	● (I/II)
IM-19 (Imunopharm)	×	×	×	×	● (I/II)
rapcabtagene autoleucel	×	×	×	×	● (I/II)
steroid, rituximab, dasatinib, chemotherapy, venetoclax, blinatumomab	×	×	×	×	● (I/II)
YK-012	×	×	×	×	● (I/II)
B-019	×	×	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

For queries please call on 9586177755

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

BCR::ABL1 fusion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
CART cell therapy	×	×	×	×	● (I)
CART-CD19, chemotherapy	×	×	×	×	● (I)
CART-CD19A, chemotherapy	×	×	×	×	● (I)
CC-312	×	×	×	×	● (I)
dasatinib, CART-CD19/CD22, chemotherapy	×	×	×	×	● (I)
edetate calcium disodium, succimer	×	×	×	×	● (I)
GNR-084	×	×	×	×	● (I)
INT-2104	×	×	×	×	● (I)
lomab-B, CART-CD19	×	×	×	×	● (I)
JY-231	×	×	×	×	● (I)
META10-19, chemotherapy	×	×	×	×	● (I)
olverembatinib, lisaftoclax, steroid	×	×	×	×	● (I)
radiation therapy, chemotherapy, stem cell engraftment therapy (Orca Biosystems), tacrolimus	×	×	×	×	● (I)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Details

Current FDA Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

FDA information is current as of 2025-05-14. For the most up-to-date information, search www.fda.gov.

BCR::ABL1 fusion

☒ dasatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-12-03

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

PHYRAGO® is indicated for the treatment of adult patients with

- newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase.
- chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy including imatinib.
- Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) with resistance or intolerance to prior therapy.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/216099s003lbl.pdf

☒ dasatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-07-31

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

SPRYCEL® is a kinase inhibitor indicated for the treatment of

- newly diagnosed adults with Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase.
- adults with chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy including imatinib.
- adults with Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) with resistance or intolerance to prior therapy.
- pediatric patients 1 year of age and older with Ph+ CML in chronic phase.
- pediatric patients 1 year of age and older with newly diagnosed Ph+ ALL in combination with chemotherapy.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/021986s028lbl.pdf

BCR::ABL1 fusion (continued)**① imatinib**

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2022-08-19

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

GLEEVEC® is a kinase inhibitor indicated for the treatment of:

- Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.
- Patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in blast crisis (BC), accelerated phase (AP), or in chronic phase (CP) after failure of interferon-alpha therapy.
- Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).
- Pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy.
- Adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements.
- Adult patients with aggressive systemic mastocytosis (ASM) without the D816V c-Kit mutation or with c-Kit mutational status unknown.
- Adult patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL) who have the FIP1L1-PDGFRα fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFRα fusion kinase negative or unknown.
- Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP).
- Patients with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors (GIST).
- Adjuvant treatment of adult patients following resection of Kit (CD117) positive GIST.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/021588s062lbl.pdf

BCR::ABL1 fusion (continued)

① imatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-11-22

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:
Imkeldi™ is a kinase inhibitor indicated for the treatment of:

- Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.
- Patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in blast crisis (BC), accelerated phase (AP), or in chronic phase (CP) after failure of interferon-alpha therapy.
- Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).
- Pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy.
- Adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements.
- Adult patients with aggressive systemic mastocytosis (ASM) without the D816V c-Kit mutation or with c-Kit mutational status unknown.
- Adult patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL) who have the FIP1L1-PDGFRα fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFRα fusion kinase negative or unknown.
- Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP).
- Patients with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors (GIST).
- Adjuvant treatment of adult patients following resection of Kit (CD117) positive GIST.

Reference:
https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/219097s000lbl.pdf

● dasatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2024-07-31

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:
SPRYCEL® is a kinase inhibitor indicated for the treatment of

- newly diagnosed adults with Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase.
- adults with chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy including imatinib.
- adults with Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) with resistance or intolerance to prior therapy.
- pediatric patients 1 year of age and older with Ph+ CML in chronic phase.
- pediatric patients 1 year of age and older with newly diagnosed Ph+ ALL in combination with chemotherapy.

Reference:
https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/021986s028lbl.pdf

Page 21 of 93

BCR::ABL1 fusion (continued)

● imatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Label as of: 2022-08-19 Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Indications and usage:

GLEEVEC® is a kinase inhibitor indicated for the treatment of:

- Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.
- Patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in blast crisis (BC), accelerated phase (AP), or in chronic phase (CP) after failure of interferon-alpha therapy.
- Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).
- Pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy.
- Adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements.
- Adult patients with aggressive systemic mastocytosis (ASM) without the D816V c-Kit mutation or with c-Kit mutational status unknown.
- Adult patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL) who have the FIP1L1-PDGFRα fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFRα fusion kinase negative or unknown.
- Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP).
- Patients with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors (GIST).
- Adjuvant treatment of adult patients following resection of Kit (CD117) positive GIST.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/021588s062lbl.pdf

BCR::ABL1 fusion (continued)

● **imatinib + chemotherapy**

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2024-11-22

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

Imkeldi™ is a kinase inhibitor indicated for the treatment of:

- Newly diagnosed adult and pediatric patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.
- Patients with Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in blast crisis (BC), accelerated phase (AP), or in chronic phase (CP) after failure of interferon-alpha therapy.
- Adult patients with relapsed or refractory Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL).
- Pediatric patients with newly diagnosed Philadelphia chromosome positive acute lymphoblastic leukemia (Ph+ ALL) in combination with chemotherapy.
- Adult patients with myelodysplastic/myeloproliferative diseases (MDS/MPD) associated with platelet-derived growth factor receptor (PDGFR) gene re-arrangements.
- Adult patients with aggressive systemic mastocytosis (ASM) without the D816V c-Kit mutation or with c-Kit mutational status unknown.
- Adult patients with hypereosinophilic syndrome (HES) and/or chronic eosinophilic leukemia (CEL) who have the FIP1L1-PDGFRα fusion kinase (mutational analysis or fluorescence in situ hybridization [FISH] demonstration of CHIC2 allele deletion) and for patients with HES and/or CEL who are FIP1L1-PDGFRα fusion kinase negative or unknown.
- Adult patients with unresectable, recurrent and/or metastatic dermatofibrosarcoma protuberans (DFSP).
- Patients with Kit (CD117) positive unresectable and/or metastatic malignant gastrointestinal stromal tumors (GIST).
- Adjuvant treatment of adult patients following resection of Kit (CD117) positive GIST.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/219097s000lbl.pdf

● **ponatinib, ponatinib + chemotherapy**

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2024-03-19

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

ICLUSIG® is a kinase inhibitor indicated for the treatment of adult patients with:

Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia (Ph+ ALL)

- Newly diagnosed Ph+ ALL, in combination with chemotherapy. This indication is approved under accelerated approval based on minimal residual disease (MRD)-negative complete remission (CR) at the end of induction. Continued approval for this indication may be contingent upon verification of clinical benefit in a confirmatory trial(s)
- As monotherapy in Ph+ ALL for whom no other kinase inhibitors are indicated or T315I-positive Ph+ ALL.

Chronic Myeloid Leukemia (CML)

- Chronic phase (CP) CML with resistance or intolerance to at least two prior kinase inhibitors.
- Accelerated phase (AP) or blast phase (BP) CML for whom no other kinase inhibitors are indicated.
- T315I-positive CML (chronic phase, accelerated phase, or blast phase).

Limitations of Use: ICLUSIG® is not indicated and is not recommended for the treatment of patients with newly diagnosed CP-CML.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/203469s037lbl.pdf

BCR::ABL1 fusion (continued)

○ asciminib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-10-29

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

SCEMBLIX® is a kinase inhibitor indicated for the treatment of adult patients with:

- Newly diagnosed Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP). This indication is approved under accelerated approval based on major molecular response rate. Continued approval for this indication may be contingent upon verification of clinical benefit in a confirmatory trial(s).
- Previously treated Ph+ CML in CP.
- Ph+ CML in CP with the T315I mutation.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/215358s009lbl.pdf

○ bosutinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-12-03

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

BOSULIF® is a kinase inhibitor indicated for the treatment of

- adult and pediatric patients 1 year of age and older with chronic phase Ph+ chronic myelogenous leukemia (CML), newly-diagnosed or resistant or intolerant to prior therapy.
- adult patients with accelerated, or blast phase Ph+ CML with resistance or intolerance to prior therapy.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/203341s026lbl.pdf

○ nilotinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-11-07

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

DANZITEN™ is a kinase inhibitor indicated for the treatment of:

- Adult patients with newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.
- Adult patients with chronic phase (CP) and accelerated phase (AP) Ph+ CML resistant to or intolerant to prior therapy that included imatinib.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/219293s000lbl.pdf

BCR::ABL1 fusion (continued)

○ nilotinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-02-08

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Indications and usage:

TASIGNA® is a kinase inhibitor indicated for the treatment of:

- Adult and pediatric patients greater than or equal to 1 year of age with newly diagnosed Philadelphia chromosome positive chronic myeloid leukemia (Ph+ CML) in chronic phase.
- Adult patients with chronic phase (CP) and accelerated phase (AP) Ph+ CML resistant to or intolerant to prior therapy that included imatinib.
- Pediatric patients greater than or equal to 1 year of age with Ph+ CML-CP and CML-AP resistant or intolerant to prior tyrosine-kinase inhibitor (TKI) therapy.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/022068s041lbl.pdf

Current NCCN Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

NCCN information is current as of 2025-05-01. To view the most recent and complete version of the guideline, go online to [NCCN.org](https://www.nccn.org).
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

BCR::ABL1 fusion

● blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]
NCCN Recommendation category: 2A

Population segment (Line of therapy):
 ■ B-cell (Consolidation therapy)
 ■ B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]
NCCN Recommendation category: 2A

Population segment (Line of therapy):
 ■ B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● bosutinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]
NCCN Recommendation category: 2A

Population segment (Line of therapy):
 ■ B-cell (Consolidation therapy)
 ■ B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● bosutinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + CALGB 10701 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● **bosutinib + cyclophosphamide + daunorubicin + vincristine + prednisone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **bosutinib + cytarabine + HyperCVAD + methotrexate**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **bosutinib + cytarabine + methotrexate + methylprednisolone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **bosutinib + EsPhALL regimen**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● bosutinib + EWALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + mercaptopurine + methotrexate + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + steroid

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● bosutinib + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● bosutinib + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● brexucabtagene autoleucel

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● dasatinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + ALL-REZ BFM 90 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● dasatinib + calaspargase pegol + cytarabine

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + CALGB 10701 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + clofarabine + cyclophosphamide + etoposide

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + COG AALL01P2 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● dasatinib + COG AALL0622 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + COG AALL0622 regimen + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + COG AALL07P1 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + COG AALL1331 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + COG AALL1631 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

- dasatinib + COG AALL1631 regimen + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):
 - B-cell (Consolidation therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]
- dasatinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):
 - B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]
- dasatinib + cyclophosphamide + daunorubicin + vincristine + prednisone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):
 - B-cell (Induction therapy); Other recommended intervention
 - B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]
- dasatinib + cytarabine + fludarabine + filgrastim

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):
 - B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● dasatinib + cytarabine + HyperCVAD + methotrexate

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + cytarabine + methotrexate + methylprednisolone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + cytarabine + PEG-L-asparaginase

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + EsPhALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● dasatinib + EWALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + FLAG-IDA

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● **dasatinib + inotuzumab ozogamicin + cyclophosphamide + vincristine + dexamethasone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **dasatinib + mercaptopurine + methotrexate + vincristine + prednisone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **dasatinib + steroid**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **dasatinib + tisagenlecleucel-t**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **dasatinib + Total Therapy XVII regimen**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● dasatinib + Total Therapy XVII regimen + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + UKALL R3 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + venetoclax + calaspargase pegol + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + venetoclax + calaspargase pegol + vincristine + prednisone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + venetoclax + PEG-L-asparaginase + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● dasatinib + venetoclax + PEG-L-asparaginase + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● dasatinib + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● imatinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib + ALL-REZ BFM 90 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● **imatinib + calaspargase pegol + cytarabine**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **imatinib + CALGB 10701 regimen**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **imatinib + clofarabine + cyclophosphamide + etoposide**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **imatinib + COG AALL01P2 regimen**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● imatinib + COG AALL07P1 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + COG AALL1331 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + COG AALL1631 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + COG AALL1631 regimen + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● **imatinib + cyclophosphamide + daunorubicin + vincristine + prednisone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **imatinib + cytarabine + fludarabine + filgrastim**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **imatinib + cytarabine + HyperCVAD + methotrexate**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **imatinib + cytarabine + methotrexate + methylprednisolone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● imatinib + cytarabine + PEG-L-asparaginase

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + EsPhALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib + EWALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib + FLAG-IDA

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● imatinib + inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib + inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + inotuzumab ozogamicin + cyclophosphamide + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + mercaptopurine + methotrexate + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● imatinib + steroid

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● imatinib + tisagenlecleucel-t

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + UKALL R3 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + venetoclax + calaspargase pegol + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● imatinib + venetoclax + calaspargase pegol + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● **imatinib + venetoclax + PEG-L-asparaginase + vincristine + dexamethasone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **imatinib + venetoclax + PEG-L-asparaginase + vincristine + prednisone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● **imatinib + vincristine + dexamethasone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● **imatinib + vincristine + prednisone**

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● inotuzumab ozogamicin

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● inotuzumab ozogamicin + nilotinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● inotuzumab ozogamicin + ponatinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● nilotinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + CALGB 10701 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● nilotinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + cyclophosphamide + daunorubicin + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + cytarabine + HyperCVAD + methotrexate

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + cytarabine + methotrexate + methylprednisolone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● nilotinib + EsPhALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + EWALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + mercaptopurine + methotrexate + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + steroid

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● nilotinib + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● nilotinib + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● obecabtagene autoleucel

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● ponatinib

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + CALGB 10701 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + cyclophosphamide + cytarabine + doxorubicin + methotrexate + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● ponatinib + cyclophosphamide + daunorubicin + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + cytarabine + HyperCVAD + methotrexate

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + cytarabine + methotrexate + methylprednisolone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + EsPhALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● ponatinib + EWALL regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell (Consolidation therapy)
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + mercaptopurine + methotrexate + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + steroid

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● ponatinib + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention
- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

● ponatinib + vincristine + prednisone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● tisagenlecleucel-t

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

● tisagenlecleucel-t

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● venetoclax + calaspargase pegol + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● venetoclax + calaspargase pegol + vincristine + prednisolone

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● venetoclax + PEG-L-asparaginase + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● venetoclax + PEG-L-asparaginase + vincristine + prednisone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● inotuzumab ozogamicin + ponatinib

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● inotuzumab ozogamicin + ponatinib + cyclophosphamide + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + ALL-REZ BFM 90 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● ponatinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + calaspargase pegol + cytarabine

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + clofarabine + cyclophosphamide + etoposide

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + COG AALL01P2 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + COG AALL07P1 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● ponatinib + COG AALL1331 regimen

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + cytarabine + fludarabine + filgrastim

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + cytarabine + PEG-L-asparaginase

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + FLAG-IDA

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + tisagenlecleucel-t

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● ponatinib + UKALL R3 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + venetoclax + calaspargase pegol + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + venetoclax + calaspargase pegol + vincristine + prednisone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + venetoclax + PEG-L-asparaginase + vincristine + dexamethasone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● ponatinib + venetoclax + PEG-L-asparaginase + vincristine + prednisone

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell; Relapsed, Refractory (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

● tisagenlecleucel-t

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- B-cell (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + COG AALL1131 regimen + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: ABL1 fusion

Other criteria: CRLF2 fusion negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + DFCI-ALL Protocol 16-001 regimen

Cancer type: Acute Lymphoblastic LeukemiaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

● dasatinib + DFCI-ALL Protocol 16-001 regimen + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- B-cell (Consolidation therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

asciminib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

bosutinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

dasatinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

imatinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

nilotinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

asciminib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

☐ bosutinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

☐ cytarabine

Cancer type: Acute Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

☐ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

○ cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

○ cytarabine + idarubicin

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

○ cytarabine + idarubicin

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

○ dasatinib

Cancer type: Accelerated Phase Chronic Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

☐ decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ gemtuzumab ozogamicin

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: CD33 positive

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ glasdegib + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ imatinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

☐ imatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

☐ liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ nilotinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

☐ ponatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (First-line therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]

BCR::ABL1 fusion (continued)

○ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

○ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

○ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

○ venetoclax + cladribine + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

☐ venetoclax + cytarabine

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ venetoclax + decitabine

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ venetoclax + decitabine

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

☐ venetoclax + decitabine

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

cladribine + cytarabine + mitoxantrone + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cladribine + cytarabine + mitoxantrone + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

(Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

cytarabine + mitoxantrone

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

(Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: Deletion 17p negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

(Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: TP53 mutation negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

(Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]

BCR::ABL1 fusion (continued)

asciminib

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Chronic Phase, Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

asciminib + chemotherapy

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

bosutinib

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Chronic Phase, Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

bosutinib + chemotherapy

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

BCR::ABL1 fusion (continued)

○ dasatinib

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention
- Chronic Phase (Line of therapy not specified); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

○ dasatinib + chemotherapy

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

○ imatinib

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Chronic Phase, Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

○ imatinib + chemotherapy

Cancer type: Myeloid/Lymphoid Neoplasms with EosinophiliaVariant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

BCR::ABL1 fusion (continued)

nilotinib

Cancer type: Myeloid/Lymphoid Neoplasms with Eosinophilia

Variant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention
- Chronic Phase (Line of therapy not specified); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

nilotinib + chemotherapy

Cancer type: Myeloid/Lymphoid Neoplasms with Eosinophilia

Variant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

ponatinib

Cancer type: Myeloid/Lymphoid Neoplasms with Eosinophilia

Variant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Chronic Phase, Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

ponatinib + chemotherapy

Cancer type: Myeloid/Lymphoid Neoplasms with Eosinophilia

Variant class: ABL1 fusion

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Blast Phase (Line of therapy not specified); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Myeloid/Lymphoid Neoplasms with Eosinophilia and Tyrosine Kinase Gene Fusions [Version 2.2025]

Current EMA Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

EMA information is current as of 2025-05-14. For the most up-to-date information, search www.ema.europa.eu.

BCR::ABL1 fusion

☒ dasatinib

Cancer type: Acute Lymphoblastic Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-05-24

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Reference:
https://www.ema.europa.eu/en/documents/product-information/sprycel-epar-product-information_en.pdf

☒ imatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-12-11

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Reference:
https://www.ema.europa.eu/en/documents/product-information/glivec-epar-product-information_en.pdf

☒ imatinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-10-01

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Reference:
https://www.ema.europa.eu/en/documents/product-information/imatinib-teva-epar-product-information_en.pdf

☒ imatinib (Accord)

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Acute Lymphoblastic Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia

Label as of: 2025-05-06

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Reference:
https://www.ema.europa.eu/en/documents/product-information/imatinib-accord-epar-product-information_en.pdf

BCR::ABL1 fusion (continued)

- dasatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2024-05-24

Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/sprycel-epar-product-information_en.pdf
- imatinib (Accord) + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2025-05-06

Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/imatinib-accord-epar-product-information_en.pdf
- imatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2024-12-11

Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/glivec-epar-product-information_en.pdf
- imatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2024-10-01

Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/imatinib-teva-epar-product-information_en.pdf
- ponatinib

Cancer type: Acute Lymphoblastic Leukemia

Label as of: 2025-02-18

Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/iclusig-epar-product-information_en.pdf
- asciminib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Label as of: 2024-12-05

Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]

Reference:

https://www.ema.europa.eu/en/documents/product-information/asciminib-epar-product-information_en.pdf

BCR::ABL1 fusion (continued)

☐ bosutinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Blast Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia	Label as of: 2025-01-16	Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]
Reference: https://www.ema.europa.eu/en/documents/product-information/bosulif-epar-product-information_en.pdf		

☐ nilotinib

Cancer type: Accelerated Phase Chronic Myeloid Leukemia, Chronic Phase Chronic Myeloid Leukemia	Label as of: 2024-12-03	Variant class: BCR::ABL1 fusion [t(9;22) (q34;q11)]
Reference: https://www.ema.europa.eu/en/documents/product-information/tasigna-epar-product-information_en.pdf		

Current ESMO Information

☒ In this cancer type ☐ In other cancer type ☒ In this cancer type and other cancer types

ESMO information is current as of 2025-05-01. For the most up-to-date information, search www.esmo.org.

BCR::ABL1 fusion

● allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: 1 / A

Population segment (Line of therapy):
■ (Consolidation therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

● asciminib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):
■ (Induction therapy, Consolidation therapy, Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

● bosutinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):
■ (Induction therapy, Consolidation therapy, Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

● dasatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):
■ (Induction therapy, Consolidation therapy, Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

BCR::ABL1 fusion (continued)

● imatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):
■ (Induction therapy, Consolidation therapy, Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

● nilotinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):
■ (Induction therapy, Consolidation therapy, Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

● ponatinib + chemotherapy

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):
■ (Induction therapy, Consolidation therapy, Maintenance therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

● dasatinib + blinatumomab

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):
■ (Consolidation therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

BCR::ABL1 fusion (continued)

● ponatinib + blinatumomab

Cancer type: Acute Lymphoblastic LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: III / A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

○ dasatinib

Cancer type: Chronic Phase Chronic Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

○ imatinib

Cancer type: Chronic Phase Chronic Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

○ nilotinib

Cancer type: Chronic Phase Chronic Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: I / A

Population segment (Line of therapy):

- (First-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

BCR::ABL1 fusion (continued)

○ allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

○ cytarabine + daunorubicin

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

○ dasatinib

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

○ imatinib

Cancer type: Acute Myeloid LeukemiaVariant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

BCR::ABL1 fusion (continued)

nilotinib

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / A

Population segment (Line of therapy):

- (Line of therapy not specified)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

cladribine + cytarabine + daunorubicin

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / C

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

cytarabine + daunorubicin + fludarabine

Cancer type: Acute Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: II / C

Population segment (Line of therapy):

- (Induction therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

bosutinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Resistant, Refractory (Second-line therapy, Third-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

BCR::ABL1 fusion (continued)

☐ dasatinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Resistant, Refractory (Second-line therapy, Third-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

☐ imatinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Resistant, Refractory (Second-line therapy, Third-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

☐ nilotinib

Cancer type: Chronic Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

ESMO Level of Evidence/Grade of Recommendation: V / A

Population segment (Line of therapy):

- Resistant, Refractory (Second-line therapy, Third-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Chronic Myeloid Leukemia [Ann Oncol (2017) 28 (suppl 4): iv41–iv51. (Corrigendum: 03 October 2018)]

Prognostic Details

Current NCCN Information

NCCN information is current as of 2025-05-01. To view the most recent and complete version of the guideline, go online to NCCN.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

BCR::ABL1 fusion

Prognostic significance: NCCN: Standard

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: IKZF1 deletion negative, PAX5 deletion negative

NCCN Recommendation category: 2A

Summary:

- Cytogenetics risk groups for B-ALL

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

Prognostic significance: NCCN: Standard

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: CDKN2B deletion negative, IKZF1 deletion negative

NCCN Recommendation category: 2A

Summary:

- Cytogenetics risk groups for B-ALL

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

Prognostic significance: NCCN: Standard

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: CDKN2A deletion negative, IKZF1 deletion negative

NCCN Recommendation category: 2A

Summary:

- Cytogenetics risk groups for B-ALL

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

BCR::ABL1 fusion (continued)

Prognostic significance: NCCN: Standard

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Other criteria: F2R deletion negative, IKZF1 deletion negative

NCCN Recommendation category: 2A

- Summary:
- Cytogenetics risk groups for B-ALL

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

Prognostic significance: NCCN: Unfavorable

Cancer type: Acute Lymphoblastic Leukemia Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Reference: NCCN Guidelines® - NCCN-Pediatric Acute Lymphoblastic Leukemia [Version 3.2025]

Diagnostic Details

Current NCCN Information

NCCN information is current as of 2025-05-01. To view the most recent and complete version of the guideline, go online to NCCN.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

BCR::ABL1 fusion

Diagnostic significance: Acute Lymphoblastic Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

NCCN Recommendation category: 2A

Diagnostic notes:

- B-cell Lymphoblastic Leukemia/Lymphoma subtype based on molecular characterization

Reference: NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]

Current ESMO Information

ESMO information is current as of 2025-05-01. For the most up-to-date information, search www.esmo.org.

BCR::ABL1 fusion

Diagnostic significance: Acute Lymphoblastic Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Lymphoblastic Leukaemia [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2023.09.3112> (eUpdate 11 October 2023, published)]

Clinical Trials Summary

BCR::ABL1 fusion

NCT ID	Title	Phase
NCT02611492	A Phase III Study, Randomized, to Evaluate the Reduction of Chemotherapy Intensity in Association With Nilotinib (Tasigna®) in Philadelphia Chromosome-positive (Ph+) ALL of Young Adults (18-59 Years Old) (GRAAPH-2014)	III
No NCT ID	CCCG-ALL-2015 Protocol Based Multi-Center Trial Of Acute Lymphoblastic Leukemia	III
NCT06051409	A Pivotal Registrational Phase III Study of Olverembatinib Combined With Chemotherapy Versus Imatinib Combined With Chemotherapy in Patients With Newly Diagnosed Philadelphia Chromosome-positive Acute Lymphoblastic Leukemia (Ph+ALL)	III
NCT04530565	A Phase III Randomized Trial of Steroids + Tyrosine Kinase Inhibitor (TKI) Induction With Chemotherapy or Blinatumomab for Newly Diagnosed BCR-ABL-Positive Acute Lymphoblastic Leukemia (ALL) in Adults	III
NCT06308588	Phase II Study of the Combination of Blinatumomab and Asciminib in Patients With Philadelphia Chromosome-Positive Acute Lymphoblastic Leukemia	II
NCT03263572	Phase II Study of the Combination of Blinatumomab and Ponatinib in Patients With Philadelphia Chromosome (Ph)-Positive and/or BCR-ABL Positive Acute Lymphoblastic Leukemia (ALL)	II
NCT03984968	Evaluation of Safety and Efficacy of CD19 CAR-T Consolidation Therapy for Acute Lymphoblastic Leukemia Concomitant With Infusion of CD19+T Cells as Feeding Cells	II
NCT06179524	Phase II Clinical Study of CAR-T-19 (Anti-CD19 Single-chain Antibody Chimeric Antigen Receptor T Cell) Injection in the Treatment of CD19-positive Relapsed/Refractory B-cell Acute Lymphoblastic Leukemia(B-ALL) Under 25 Years of Age (Inclusive)	II
No NCT ID	Phase II clinical study of autologous reinfusion preparation targeting CD19 chimeric antigen receptor T lymphocytes (pCAR-19B) in the treatment of CD19-positive relapsed/refractory B-cell acute lymphoblastic leukemia (22-70 years old)	II
NCT05334823	Phase II clinical study of CD19-targeted chimeric antigen receptor T lymphocyte (pCAR-19B) autologous infusion preparation in the treatment of CD19-positive relapsed/refractory B-cell acute lymphoblastic leukemia	II
NCT05827835	A Study to Evaluate the Efficacy and Safety of CD7CAR-T Bridging to Allogeneic Hematopoietic Stem Cell Transplantation in Patients With Refractory or Relapsed CD7 Positive Malignant Hematologic Diseases	II
No NCT ID	CLIC-1901 CAR T-cells for treatment of patients with relapsed/refractory CD19-positive ALL and NHL (DAN-CART 1901)	II
No NCT ID	Clinical Study of Efficacy And Safety Of Flumatinib Combined With Autologous Hematopoietic Stem Cell Transplantation In The Treatment Of Ph+ Acute Lymphoblastic Leukemia	II
No NCT ID	Multicenter Clinical Study Of Flumatinib Combined With Chemotherapy In The Treatment Of Adults Patients (Aged 16-65 Years) Newly Diagnosed With Ph+ALL	II
NCT03441061	Phase II Study of Inotuzumab Ozogamicin in Patients with B-Cell Lineage Acute Lymphocytic Leukemia with Positive Minimal Residual Disease	II
NCT03610438	A Phase IIA Study of Feasibility and Effectiveness of Inotuzumab Ozogamicin (IO) in Adult Patients With B-Cell Acute Lymphoblastic Leukemia With Positive Minimal Residual Disease Before Any Hematopoietic Stem Cell Transplantation	II
NCT02981628	A Phase II Study of Inotuzumab Ozogamicin (InO) (NCT03977218) in Children and Young Adults With Relapsed or Refractory CD22+ B-Acute Lymphoblastic Leukemia (B-ALL)	II
NCT04747912	A Phase II Study of a Chemotherapy-Free Induction Regimen for Ph+ Acute Lymphoblastic Leukemia (ALL) Incorporating Inotuzumab Ozogamicin (InO)	II

For queries please call on 9586177755

Clinical Trials Summary (continued)

BCR::ABL1 fusion (continued)

NCT ID	Title	Phase
No NCT ID	Role of olverembatinib with vindesine or vincristine and prednisolone in the major molecular remission for newly diagnosed adult philadelphia chromosome-positive acute lymphoblastic leukemia in the induction therapy: A single-arm, multi-center, prospective study	II
NCT05594784	Olverembatinib Combined With Reduced-Intensity Chemotherapy and Venetoclax for Newly Diagnosed Philadelphia Chromosome-positive Acute Lymphoblastic Leukemia: A Prospective, Single-arm, Single-center Clinical Trial	II
NCT06754267	Venetoclax Combined With Olverembatinib and Prednisone in Treating Ph-positive Precursor B Cell Acute Lymphoblastic Leukemia: a Phase II, Single Arm and Multicenter Study	II
NCT06578546	A Single-arm, Open Label, Multicenter Clinical Study to Evaluate the Safety and Efficacy of VAO Regimen in Patients With Newly Diagnosed Ph-positive Acute Lymphoblastic Leukemia	II
No NCT ID	Safety and Feasibility of de-centralized CAR-T cell manufacturing and treatment of relapsed and refractory B cell acute leukemias and lymphomas	I/II
NCT04271410	CD19+Targeted CAR-T Cell Therapy for Relapsed/Refractory CD19+ B Cell Leukemia and Lymphoma	I/II
NCT04271800	Safety and Efficacy of CD19-Targeted CAR-T Therapy for Relapsed/Refractory CD19+ B Cell Leukemia and Lymphoma	I/II
NCT03666000	Phase I/Ib, Open-label, Dose-escalation, Dose-expansion, Parallel Assignment Study to Evaluate Safety and Clinical Activity of PBCAR0191 (azercabtagene Zapreleucel or "azer-cel") in Subjects with Relapsed/Refractory (r/r) Non-Hodgkin Lymphoma (NHL) and R/r B-cell Acute Lymphoblastic Leukemia (B-ALL)	I/II
NCT03016377	Administration of Autologous CAR-T Cells Targeting the CD19 Antigen and Containing the Inducible caspase9 Safety Switch in Patients With Relapsed/Refractory Acute Lymphoblastic Leukemia	I/II
NCT04781634	Study Evaluating Safety and Efficacy of CD19 and CD22 Targeted Prime CAR-T Cell in Patients With Relapsed/Refractory B-ALL	I/II
NCT06213636	T-cell Infusion Targeting CD19 and CD22 for Refractory/Relapsed Leukemia/Lymphoma Patients With or Without Central Nervous System Involvement	I/II
NCT05579132	An Open-label, Multi-center Phase Ib/II Study of CN201 in Adult Subjects With Precursor B-cell Acute Lymphoblastic Leukemia	I/II
NCT05309213	A Phase I/II Study to Evaluate the Safety and Efficacy of IM19 CAR-T Cells in Patients With Relapsed and Refractory (R/R) B-cell Acute Lymphoblastic Leukemia	I/II
NCT03104491	Inotuzumab Ozogamicin Post-Transplant For Acute Lymphocytic Leukemia	I/II
NCT01371630	Phase I/II Study of the Combination of Inotuzumab Ozogamycin (CMC-544) With Low-Intensity Chemotherapy in Patients With Acute Lymphoblastic Leukemia (ALL)	I/II
NCT03934372	An Open-Label, Single-Arm, Phase I/II Study Evaluating the Safety and Efficacy of Ponatinib for the Treatment of Recurrent or Refractory Leukemias, Lymphomas or Solid Tumors in Pediatric Participants	I/II
NCT03960840	Phase I/II, Open Label, Multicenter Study of Rapcabtagene Autoleucel in Adult Patients With CLL/SLL, 3L+ DLBCL, r/r ALL and 1L HR LBCL	I/II
NCT04872790	A Dose-Finding Phase Ib Study of the Oral BCL-2 Inhibitor Venetoclax (ABT-199) in Combination With Standard Induction Therapy, Dasatinib and Prednisone (and Rituximab in CD20+ Patients), in Adult Patients With Newly Diagnosed and Relapsed Philadelphia Chromosome Positive (Ph+) ALL and Ph+ MPAL	I/II
NCT06580301	An Open-Label, Multi-Center, Phase Ib/II Study to Evaluate the Safety, Tolerability, Pharmacokinetics and Preliminary Anti-tumor Activity of YK012 in Participants With B-cell Acute Lymphoblastic Leukemia	I/II

For queries please call on 9586177755

Clinical Trials Summary (continued)

BCR::ABL1 fusion (continued)

NCT ID	Title	Phase
NCT06927466	A Clinical Study to Evaluate the Safety, Tolerability, and Preliminary Efficacy of B019 Injection in Subjects With Relapsed or Refractory B-cell Acute Lymphoblastic Leukemia.	I
NCT05038696	Chimeric-Antigen Receptor (CAR) T-Cell Therapy Using Multiple CARs and Cell Marker Profiling in High Risk and Relapsed/ Refractory B-Lineage Acute Lymphoblastic Leukaemia	I
NCT06307600	The Safety and Efficacy of RD06-03 CART Cell Injection in Patients With Relapsed or Refractory B- lymphoblastic Leukemia	I
NCT06756321	An Exploratory Clinical Study of the Safety and Efficacy of Chimeric Antigen Receptor T-Cells (CAR T- Cells) in Subjects with Relapsed/Refractory Hematologic Malignancy	I
NCT03825731	A Study of GC022 CAR-T Cell Immunotherapy for Relapsed or Refractory B- ALL	I
NCT05350787	To Evaluate the Safety, Efficacy and Pharmacokinetics of ThisCART19A in Patients With Relapsed and Refractory Acute B-cell Leukemia	I
NCT05679687	A Study to Evaluate the Efficacy, Safety, and Pharmacokinetics of Allogeneic Anti CD19 CAR-T Bridging to Hematopoietic Stem Cell Transplantation in Patients With Refractory or Relapsed B Cell Acute Lymphoblastic Leukemia	I
NCT06037018	A Phase I Study to Evaluate the Safety, Pharmacokinetics, Pharmacodynamics, and Preliminary Anti-tumor Efficacy of CC312 in Adult Patients With Relapsed/Refractory CD19 Positive B-cell Hematologic Malignancies	I
NCT05523661	The Safety and Clinical Efficacy of Dasatinib Plus Human CD19/CD22 Bispecific CAR-T Cell Therapy for Elderly Subjects With Ph-positive Acute Lymphoblastic Leukemia	I
NCT03630991	Monitoring, Detoxifying, and Rebalancing Metals During Front Line Acute Myeloid Leukemia (AML) and Myelodysplastic Syndrome (MDS) Therapy	I
NCT04601584	Dose-escalation Sequention Cohort Study of Safety, Tolerability, Pharmacokinetics and Pharmacodynamics of GNR-084 in Patients With Refractory or Relapse Acute Lymphoblastic B-cell Precursor Leukemia.	I
NCT05480501	Clinical Trial to Evaluate the Safety and Efficacy of IM19 CAR-T Cells in Patients With Relapsed and Refractory (R/R) B-cell Acute Lymphoblastic Leukemia	I
NCT06539338	A Two-Part Open Label Phase I Multicentre Study Evaluating the Safety of INT2104 Infusions in Female and Male Participants Aged 18 Years of Age and Older With Refractory/Relapsing B-cell Malignancies	I
NCT04512716	Iomab-ACT: A Pilot Study of 131-I Apamistamab Followed by CD19-Targeted CAR T-Cell Therapy for Patients With Relapsed or Refractory B-Cell Acute Lymphoblastic Leukemia or Diffuse Large B-Cell Lymphoma	I
NCT06678282	JY231 Injection for the Treatment of Relapsed or Refractory B Cell Lymphoma/ Leukemia - A Safety, Tolerability, and Efficacy Study	I
NCT05747157	Safety and Efficacy of Metabolically Armed CD19 CAR-T Cells (Meta10- 19) in the Treatment of r/r B- ALL Clinical Research	I
NCT06393335	Safety and Efficacy of Metabolically Armed CD19 CAR-T Cells (Meta10-19) in the Treatment of Relapsed and/or Refractory CD19-positive B Cell Hematological Malignancies Clinical Research	I
NCT06173518	A Single-Arm, Open-Label, Multi-Centre, Phase Ib Study Evaluating the Safety and Preliminary Efficacy of AUTO1 in Pediatric Patients With CD19-Positive Relapsed/ Refractory (r/r) B Cell Acute Lymphoblastic Leukemia (B ALL) and Aggressive Mature B Cell Non-Hodgkin Lymphoma (B NHL)	I

Clinical Trials Summary (continued)

BCR::ABL1 fusion (continued)

NCT ID	Title	Phase
NCT05495035	Study for Safety and Efficacy Evaluation of Olverembatinib Combined With APG-2575 in Children With Relapsed/Refractory Philadelphia Chromosome-positive Acute Lymphoblastic Leukemia (R/R Ph + ALL#	I
NCT06195891	A Single Center, Non-Randomized, Phase 1b Study of Orca-T Following Escalated Dose of Total Marrow and Lymphoid Irradiation in Patients With Acute Leukemias and MDS	I
NCT06013423	Optimized Cord Blood Transplantation for the Treatment of High-Risk Hematologic Malignancies in Adults and Pediatrics	II
NCT05805605	Allogeneic Hematopoietic Stem Cell Transplantation Using Reduced Intensity Conditioning (RIC) With Post-Transplant Cytoxan (PTCy) for the Treatment of Hematological Diseases	II
NCT02727803	Personalized NK Cell Therapy in CBT	II
NCT05800210	Alpha/Beta T Cell and CD19+ B Cell Depletion in Allogeneic Stem Cell Transplantation in Patients With Malignant Diseases	II

Alerts Informed By Public Data Sources

Current NCCN Information

 Contraindicated

 Not recommended

 Resistance

 Breakthrough

 Fast Track

NCCN information is current as of 2025-05-01. To view the most recent and complete version of the guideline, go online to NCCN.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

BCR::ABL1 fusion

 **imatinib**

Cancer type: Accelerated Phase Chronic Myeloid Leukemia

Variant class: BCR::ABL1 fusion [t(9;22)(q34;q11)]

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Imatinib is not recommended for patients with disease progression on prior TKI therapy."

Reference: NCCN Guidelines® - NCCN-Chronic Myeloid Leukemia [Version 3.2025]