

2830. Maximize the Profit as the Salesman

Description

You are given an integer `n` representing the number of houses on a number line, numbered from `0` to `n - 1`.

Additionally, you are given a 2D integer array `offers` where `offers[i] = [starti, endi, goldi]`, indicating that `ith` buyer wants to buy all the houses from `starti` to `endi` for `goldi` amount of gold.

As a salesman, your goal is to **maximize** your earnings by strategically selecting and selling houses to buyers.

Return *the maximum amount of gold you can earn*.

Note that different buyers can't buy the same house, and some houses may remain unsold.

Example 1:

Input: `n = 5, offers = [[0,0,1],[0,2,2],[1,3,2]]`

Output: `3`

Explanation: There are 5 houses numbered from 0 to 4 and there are 3 purchase offers.

We sell houses in the range `[0,0]` to 1st buyer for 1 gold and houses in the range `[1,3]` to 3rd buyer for 2 golds.

It can be proven that 3 is the maximum amount of gold we can achieve.

Example 2:

Input: `n = 5, offers = [[0,0,1],[0,2,10],[1,3,2]]`

Output: `10`

Explanation: There are 5 houses numbered from 0 to 4 and there are 3 purchase offers.

We sell houses in the range `[0,2]` to 2nd buyer for 10 golds.

It can be proven that 10 is the maximum amount of gold we can achieve.

Constraints:

- `1 <= n <= 105`
- `1 <= offers.length <= 105`
- `offers[i].length == 3`
- `0 <= starti <= endi <= n - 1`
- `1 <= goldi <= 103`

