2172. Maximum AND Sum of Array

Description

You are given an integer array nums of length n and an integer numSlots such that 2 * numSlots >= n. There are numSlots slots numbered from 1 to numSlots.

You have to place all n integers into the slots such that each slot contains at **most** two numbers. The **AND sum** of a given placement is the sum of the **bitwise** AND of every number with its respective slot number.

• For example, the AND sum of placing the numbers [1, 3] into slot [1, 3] and [4, 6] into slot [2, 6] is equal to [1, 3] the property of [1, 3] into slot [2, 6] into slo

Return the maximum possible AND sum of nums given numSlots slots.

Example 1:

```
Input: nums = [1,2,3,4,5,6], numSlots = 3

Output: 9

Explanation: One possible placement is [1, 4] into slot \underline{1}, [2, 6] into slot \underline{2}, and [3, 5] into slot \underline{3}.

This gives the maximum AND sum of (1 AND \underline{1}) + (4 AND \underline{1}) + (2 AND \underline{2}) + (6 AND \underline{2}) + (3 AND \underline{3}) + (5 AND \underline{3}) = 1 + 0 + 2 + 2 + 3 + 1 = 9.
```

Example 2:

```
Input: nums = [1,3,10,4,7,1], numSlots = 9
Output: 24
Explanation: One possible placement is [1, 1] into slot \underline{1}, [3] into slot \underline{3}, [4] into slot \underline{4}, [7] into slot \underline{7}, and [10] into slot \underline{9}.
This gives the maximum AND sum of (1 AND \underline{1}) + (1 AND \underline{1}) + (3 AND \underline{3}) + (4 AND \underline{4}) + (7 AND \underline{7}) + (10 AND \underline{9}) = 1 + 1 + 3 + 4 + 7 + 8 = 24.
Note that slots 2, 5, 6, and 8 are empty which is permitted.
```

Constraints:

- n == nums.length
- 1 <= numSlots <= 9
- 1 <= n <= 2 * numSlots
- 1 <= nums[i] <= 15