

1655. Distribute Repeating Integers

Description

You are given an array of `n` integers, `nums`, where there are at most `50` unique values in the array. You are also given an array of `m` customer order quantities, `quantity`, where `quantity[i]` is the amount of integers the `ith` customer ordered. Determine if it is possible to distribute `nums` such that:

- The `ith` customer gets **exactly** `quantity[i]` integers,
- The integers the `ith` customer gets are **all equal**, and
- Every customer is satisfied.

Return `true` *if it is possible to distribute* `nums` *according to the above conditions*.

Example 1:

Input: `nums = [1,2,3,4]`, `quantity = [2]`
Output: `false`
Explanation: The `0th` customer cannot be given two different integers.

Example 2:

Input: `nums = [1,2,3,3]`, `quantity = [2]`
Output: `true`
Explanation: The `0th` customer is given `[3,3]`. The integers `[1,2]` are not used.

Example 3:

Input: `nums = [1,1,2,2]`, `quantity = [2,2]`
Output: `true`
Explanation: The `0th` customer is given `[1,1]`, and the `1st` customer is given `[2,2]`.

Constraints:

- `n == nums.length`
- `1 <= n <= 105`
- `1 <= nums[i] <= 1000`
- `m == quantity.length`
- `1 <= m <= 10`
- `1 <= quantity[i] <= 105`
- There are at most `50` unique values in `nums`.

