

2874. Maximum Value of an Ordered Triplet II

Description

You are given a **0-indexed** integer array `nums`.

Return **the maximum value over all triplets of indices** `(i, j, k)` **such that** `i < j < k`. If all such triplets have a negative value, return `0`.

The **value of a triplet of indices** `(i, j, k)` is equal to `(nums[i] - nums[j]) * nums[k]`.

Example 1:

Input: `nums = [12,6,1,2,7]`

Output: `77`

Explanation: The value of the triplet `(0, 2, 4)` is `(nums[0] - nums[2]) * nums[4] = 77`.

It can be shown that there are no ordered triplets of indices with a value greater than 77.

Example 2:

Input: `nums = [1,10,3,4,19]`

Output: `133`

Explanation: The value of the triplet `(1, 2, 4)` is `(nums[1] - nums[2]) * nums[4] = 133`.

It can be shown that there are no ordered triplets of indices with a value greater than 133.

Example 3:

Input: `nums = [1,2,3]`

Output: `0`

Explanation: The only ordered triplet of indices `(0, 1, 2)` has a negative value of `(nums[0] - nums[1]) * nums[2] = -3`. Hence, the answer would be `0`.

Constraints:

- `3 <= nums.length <= 105`
- `1 <= nums[i] <= 106`

