

# Anosov Diffeomorphisms on Tori (Summer 2022, Yi Shi)

Ajorda Jiao

## Contents

1 July 19

1

## §1 July 19

Basic settings

- $M$  a closed Riemannian manifold.
- $f : M \rightarrow M$  a diffeomorphism.
- $\text{Diff}^r(M) := \{C^r\text{-diffeomorphisms on } M\}$ , equipped with the  $C^r$ -topology

$$d_{C^r}(f, g) := \sup_{x \in M} \{d(f(x), g(x)), d(Df(x), Dg(x)), \dots, d(D^r f(x), D^r g(x))\}.$$

**Definition 1.1.** We say  $f \in \text{Diff}^r(M)$  is an Anosov, if there exists a continuous  $Df$ -invariant splitting  $TM = E^s \oplus E^u$  and constants  $C \geq 1, 0 < \lambda < 1$  such that

- (1)  $\|Df^n|_{E^s(x)}\| \leq C\lambda^n, \forall x \in M, \forall n \geq 0.$
- (2)  $\|Df^{-n}|_{E^u(x)}\| \leq C\lambda^n, \forall x \in M, \forall n \geq 0.$

**Remark 1.2** —  $Df$  is uniformly contracting on  $E^s$ , uniformly expanding on  $E^u$ .

**Remark 1.3** —  $E^s, E^u$  are two continuous plane fields on  $M$ .

### Example 1.4

$A = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$  acts on  $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$  linearly. Then  $A$  is an Anosov diffeomorphism.

### Example 1.5

Similarly,  $A \in \text{GL}(d, \mathbb{Z})$  has no eigenvalues with modules 1, then  $A : \mathbb{T}^d \rightarrow \mathbb{T}^d$  is Anosov.

For every  $C^r$ -Anosov diffeomorphisms,  $E^s$ ,  $E^u$  are Hölder continuous. But in general,  $E^s$  and  $E^u$  are not differentiable.

We also concern about if  $E^s$  and  $E^u$  are integrable. The answer is given by the famous **Stable Manifold Theorem**. It says that both  $E^s$  and  $E^u$  are locally uniquely integrable, i.e. for every  $x \in M$ , there exists an immersed  $C^r$ -smooth submanifold  $\mathcal{F}^s(x)$  such that

- (1)  $x \in \mathcal{F}^s(x)$  and  $\mathcal{F}^s(x)$  is tangent to  $E^s$  everywhere, i.e.

$$T_y \mathcal{F}^s(x) = E^s(y), \quad \forall y \in \mathcal{F}^s(x).$$

- (2) For every local curve  $\gamma$  containing  $x$  and tangent to  $E^s$  everywhere, then  $\gamma \subset \mathcal{F}^s(x)$ .

#### Theorem 1.6 (Stable Manifold Theorem)

There exists two family of continuous foliations  $\mathcal{F}^s$  and  $\mathcal{F}^u$  with  $C^r$ -leaves tangent to  $E^s$  and  $E^u$  respectively.

**Remark 1.7** — If  $\mathcal{F}^s$  has dimension 1, then every leaf  $\mathcal{F}^s(x)$  is diffeomorphic to  $\mathbb{R}$ . (no leaf is  $\mathbb{S}^1$ ) The intuitive is that if a leaf  $\Gamma \cong \mathbb{S}^1$ , then  $\text{diam}(f^k(\Gamma)) \rightarrow 0$ . Then the orientations of  $E^s$  on  $f^k(\Gamma)$  will contradict with the uniformly continuous of  $E^s$ .

**Fact 1.8.**  $\mathbb{T}^2$  is the only 2-dimensional manifold supports an Anosov diffeomorphism.

### Foliations

Let  $M$  be an  $m$ -dimensional  $C^\infty$  closed Riemannian manifold.

**Definition 1.9.** A  $C^r$  **foliation** of dimension  $n$  is a  $C^r$ -atlas  $\mathcal{F}$  on  $M$  which is maximal satisfying

- (1) For every  $(U, \varphi) \in \mathcal{F}$ ,  $U \subset M$  is an open subset such that  $\varphi(U) = U_1 \times U_2 \subset \mathbb{R}^n \times \mathbb{R}^{m-n}$  where  $U_1, U_2$  are open disks in  $\mathbb{R}^n$  and  $\mathbb{R}^{m-n}$ .
- (2) For every  $(U, \varphi), (V, \psi) \in \mathcal{F}$  such that  $U \cap V \neq \emptyset$ , then  $\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$  has the form  $\psi \circ \varphi^{-1}(x, y) = (h_1(x, y), h_2(y))$ .

We say  $M$  is **foliated** by  $\mathcal{F}$ .

**Remark 1.10** — Stable foliation of an Anosov diffeomorphism is a  $C^0$ -foliation with  $C^r$ -leaves. That is, for every  $(U, \varphi) \in \mathcal{F}$ , write

$$\varphi : U \rightarrow \mathbb{R}^n \times \mathbb{R}^{m-n}, \quad x \mapsto (\varphi_1(x), \varphi_2(x)),$$

then  $\varphi_1$  is  $C^r$ -smooth but  $\varphi_2$  is only  $C^0$ -continuous.

### The space of Anosov diffeomorphisms

**Claim 1.11.** If  $f : M \rightarrow M$  is Anosov, then there exists a  $C^1$ -neighborhood  $\mathcal{U}$  of  $f$  in  $\text{Diff}^1(M)$  such that  $\forall g \in \mathcal{U}$  is Anosov. That is, the space of Anosov diffeomorphisms is open in  $\text{Diff}^r(M)$ .

**Proof. Cone Arguments.** Take  $\gamma \in (0, 1)$ , denote

$$\mathcal{C}_\gamma^u(x) := \{v \in T_x M : v = v^s + v^u \in E^s(x) \oplus E^u(x), \|v^s\| \leq \gamma \|v^u\|\},$$

$\mathcal{C}_\gamma^u = \bigcup_{x \in M} \mathcal{C}_\gamma^u(x)$ . Similarly, we can define  $\mathcal{C}_\gamma^s(x)$  and  $\mathcal{C}_\gamma^s$ . For  $\gamma$  small enough, there exists  $\lambda \in (0, 1)$  such that

$$Df(\mathcal{C}_\gamma^u) \subset \mathcal{C}_{\lambda\gamma}^u, \quad Df^{-1}(\mathcal{C}_\gamma^s) \subset \mathcal{C}_{\lambda\gamma}^s.$$

Besides,  $Df$  expands vectors in  $\mathcal{C}_\gamma^u$  and  $Df^{-1}$  expands vectors in  $\mathcal{C}_\gamma^s$ . **This is a robust property.** Then for  $g$  which is  $C^1$ -closed to  $f$ , we have

$$Dg(\mathcal{C}_\gamma^u) \subset \mathcal{C}_{\sqrt{\lambda}\gamma}^u, \quad Dg^{-1}(\mathcal{C}_\gamma^s) \subset \mathcal{C}_{\sqrt{\lambda}\gamma}^s,$$

$Dg$  expands vectors in  $\mathcal{C}_\gamma^u$  and  $Dg^{-1}$  expands vectors in  $\mathcal{C}_\gamma^s$ . Take

$$E_g^u(x) = \bigcap_{n \geq 0} g^{-n}(\mathcal{C}_\gamma^u(g^n x)), \quad E_g^s(x) = \bigcap_{n \geq 0} g^n(\mathcal{C}_\gamma^s(g^{-n} x)),$$

which is an Anosov splitting. □

A question is whether our examples on tori are special. But up to now, all known Anosov diffeomorphisms are supported on infra-nilmanifolds.

### Open problems

1. Are there Anosov diffeomorphisms not supported on infra-nilmanifolds?
  - All expanding maps are supported on infra-manifold.
2. Are there Anosov diffeomorphisms on simply connected manifolds?
  - Gogolev, etc. Acta Math, 2015. They constructed a partially hyperbolic diffeomorphism on simply connected 6-manifolds.
  - $\mathbb{S}^3 \times \mathbb{S}^3$  has Anosov diffeomorphisms or not?
3. Is every Anosov diffeomorphism transitive?
  - Yes on infra-manifolds, but don't know on other manifolds.

The answer of both problems are unknown on even 4-dimensional manifolds.