

# **Harmonic Analysis**

## **(Spring 2022, Dongyi Wei)**

Ajorda Jiao

# Contents

<b>1</b>	<b>Fourier Series and Integrals</b>	<b>3</b>
1.1	Fourier series . . . . .	3
1.2	The pointwise convergence . . . . .	3
1.3	Fourier series of continuous functions . . . . .	4
1.4	Convergence in norm . . . . .	4
1.5	Summability method . . . . .	5
1.6	The Fourier transform of $L^1$ functions . . . . .	6
1.7	The Schwartz class and tempered distributions . . . . .	7
1.8	The Fourier transform on $L^p, 1 < p \leq 2$ . . . . .	8
1.9	The convergence and summability of Fourier integral . . . . .	9
<b>2</b>	<b>The Hardy-Littlewood Maximal Function</b>	<b>10</b>
2.1	Approximations of the identity . . . . .	10
2.2	Weak-type inequalities and almost everywhere convergence . . . . .	10
2.3	The Marcinkiewicz interpolation theorem . . . . .	11
2.4	The Hardy-Littlewood maximal function . . . . .	11
2.5	The dyadic maximal function . . . . .	14
2.6	Some other covering lemmas . . . . .	15

# 1 Fourier Series and Integrals

## §1.1 Fourier series

For  $f \in L^1(\mathbb{T})$ , define the **Fourier coefficients**

$$\widehat{f}(k) := \int_0^1 f(x) e^{-2\pi i k x} dx.$$

Let

$$\sum_{k \in \mathbb{Z}} \widehat{f}(k) e^{2\pi i k x}$$

be the **Fourier series** of  $f$ . When we discuss the convergence of Fourier series, we consider two types of sum:

$$S_N f = \sum_{k=-N}^N \widehat{f}(k) e^{2\pi i k x}, \quad \sigma_N f = \frac{1}{N+1} \sum_{k=0}^N S_k f.$$

We concern about the following questions:

**Question 1.1.1.** The pointwise convergence of  $S_N f$ .

**Question 1.1.2.** The  $L^p$  convergence of  $S_N f$ .

**Question 1.1.3.** The almost everywhere convergence of  $S_N f$ .

**Question 1.1.4.** The convergence of  $\sigma_N f$ .

## §1.2 The pointwise convergence

**Definition 1.2.1.** The **Dirichlet kernel**  $D_N$  is given by

$$D_N(t) := \sum_{k=-N}^N e^{2\pi i k t} = \frac{\sin(\pi(2N+1)t)}{\sin(\pi t)}.$$

It satisfies

$$\int_0^1 D_N(t) dt = 1.$$

### **Theorem 1.2.2** (Dini's Criterion)

For  $x \in \mathbb{T}$ , if  $\exists \delta > 0$ , such that

$$\int_{|t|<\delta} \left| \frac{f(x+t) - f(x)}{t} \right| dt < \infty,$$

then  $S_N f(x) \rightarrow f(x)$ .

**Theorem 1.2.3**

If  $f$  is bounded variation on a neighborhood of  $x$ , then

$$S_N f(x) \rightarrow \frac{f(x+) + f(x-)}{2}.$$

**Example 1.2.4**

$f_1(t) = |t|^{-\alpha} \mathbb{1}_{(0,1/2)}$ ,  $f_2(t) = t^\alpha \sin \frac{1}{t} \mathbb{1}_{(0,1/2)}$ , where  $\alpha \in (0, 1)$ .

**Theorem 1.2.5 (Riemann Localization Principle)**

If  $f$  is zero in a neighborhood of  $x$ , then  $S_N f(x) \rightarrow 0$ .

**Theorem 1.2.6 (Riemann-Lebesgue)**

If  $f \in L^1(\mathbb{T})$ , then  $\widehat{f}(k) \rightarrow 0 (|k| \rightarrow \infty)$ .

**§1.3 Fourier series of continuous functions****Theorem 1.3.1**

There exists  $f \in C(\mathbb{T})$  such that  $S_N f(0)$  diverges.

*Proof.* Consider  $T_N : C(\mathbb{T}) \rightarrow \mathbb{C}, f \mapsto S_N f(0)$ . By theorem 1.3.2, it suffices to show  $\sup \|T_N\| = \infty$ . Suppose  $L_N = \|D_N\|_1$ , we can prove that  $\|T_N\| = L_N$ . Consider the functions  $f_n(t) = \frac{n D_N(t)}{1+n|D_N(t)|}$  is enough. The statement follows by lemma 1.3.3.  $\square$

**Theorem 1.3.2 (Uniform Boundedness Principle)**

$X, Y$ , Banach Spaces.  $\{T_a\}_{a \in A}$  is a family of bounded linear operators from  $X$  to  $Y$ . Then one of the following holds:

1.  $\sup_{a \in A} \|T_a\| < \infty$ .
2.  $\exists x \in X$ , such that  $\sup_{a \in A} \|T_a x\| = \infty$ .

**Lemma 1.3.3**

$$L_N = \frac{4}{\pi^2} \ln N + O(1).$$

**§1.4 Convergence in norm**

**Question 1.4.1.** We can ask:

1. Does  $\|S_N f - f\|_p \rightarrow 0$  for  $f \in L^p(\mathbb{T})$ ?
2. Does  $S_N f \rightarrow f$  a.e. for  $f \in L^p(\mathbb{T})$ ?

**Lemma 1.4.2**

$S_N f$  convergence to  $f$  in  $L^p$  norm,  $1 \leq p < \infty$ , iff exists  $C_p$  such that

$$\|S_N f\|_p \leq C_p \|f\|_p.$$

## §1.5 Summability method

**Definition 1.5.1.** The **Fejér kernel** is given by

$$F_N(t) := \frac{1}{N+1} \sum_{k=0}^N D_k(t) = \frac{1}{N+1} \left( \frac{\sin(\pi(N+1)t)}{\sin(\pi t)} \right)^2$$

It satisfies

$$\int_0^1 F_N(t) dt = 1 \text{ and } F_N(t) \geq 0.$$

**Theorem 1.5.2**

If  $f \in L^p(\mathbb{T})$ ,  $1 \leq p < \infty$ , or  $f \in C(\mathbb{T})$  and  $p = \infty$ , then

$$\|\sigma_N f - f\|_p \rightarrow 0.$$

*Proof.* Applying Minkowski's inequality and it follows by Fejér kernel is a good kernel.  $\square$

**Corollary 1.5.3**

The following holds:

1. The trigonometric polynomials  $V = \left\{ \sum_{k=-N}^N c_k e^{2\pi i k x} : c_k \in \mathbb{C}, N \in \mathbb{Z}_+ \right\}$  is dense in  $L^p(\mathbb{T})$ .
2. If  $f \in L^1(\mathbb{T})$  and  $\hat{f}(k) = 0$  for every  $k \in \mathbb{Z}$ , then  $f = 0$  a.e. .

**Theorem 1.5.4**

$$\|f\|_2 = \sum_{k=-\infty}^{\infty} |\hat{f}(k)|^2 \text{ and } \|S_N f\|_2 \leq \|f\|_2.$$

Define the **Poisson kernel**

$$P_r(t) = \sum_{k=-\infty}^{\infty} r^{|k|} e^{2\pi i k t} = \frac{1-r^2}{1-2r \cos(2\pi t) + r^2} = \frac{1-|z|^2}{|1-z|^2}, \quad z = r e^{2\pi i t}.$$

Let

$$u(z) = \sum_{k=0}^{\infty} \widehat{f}(k) z^k + \sum_{k=-\infty}^{-1} \widehat{f}(k) \bar{z}^{|k|}$$

be the Poisson sum, then  $u(re^{2\pi i\theta}) = P_r * f(\theta)$ .

### Theorem 1.5.5

If  $f \in L^p(\mathbb{T})$ ,  $1 \leq p < \infty$ , or  $f \in C(\mathbb{T})$  and  $p = \infty$ , then

$$\|P_r * f - f\|_p \rightarrow 0 (r \rightarrow 1^-).$$

**Remark 1.5.6** —  $\Delta u = 0$  in  $D = \{z \in \mathbb{C} : |z| < 1\}$ , and  $\mathbb{T} \cong \partial D = \mathbb{S}^1$ . If  $f \in C(\mathbb{T})$ , then  $u \in C(\overline{D})$  and  $u = f$  on  $\partial D$ .

**Fact 1.5.7.**  $\sigma_N f \rightarrow f$  a.e. and  $P_r * f \rightarrow f$  a.e. . We will prove these in the next chapter.

## §1.6 The Fourier transform of $L^1$ functions

For  $f \in L^1(\mathbb{R}^n)$ , let

$$\widehat{f}(\xi) = \int_{\mathbb{R}^n} f(x) e^{2\pi i \xi \cdot x} dx = (\mathcal{F}f)(\xi).$$

### Proposition 1.6.1

The following holds:

1.  $\widehat{\alpha f + \beta g} = \alpha \widehat{f} + \beta \widehat{g}$ .
2.  $\|\widehat{f}\|_{\infty} \leq \|f\|_1$  and  $\widehat{f} \in C(\mathbb{R}^n)$ .
3.  $\lim_{|\xi| \rightarrow \infty} \widehat{f}(\xi) = 0$ .
4.  $\widehat{f * g} = \widehat{f} \widehat{g}$ .
5.  $\widehat{\tau_h f} = \widehat{f}(\xi) e^{2\pi i h \cdot \xi}$  where  $\tau_h f = f(\cdot + h)$ .  $\widehat{f e^{2\pi i h \cdot x}}(\xi) = \widehat{f}(\xi - h)$ .
6.  $\rho \in O_n$ , then  $\widehat{f(\rho \cdot)}(\xi) = \widehat{f}(\rho \xi)$ .
7. If  $g(x) = \lambda^{-n} f(\lambda^{-1} x)$ , then  $\widehat{g}(\xi) = \widehat{f}(\lambda \xi)$  for every  $\lambda > 0$ .
8.  $\widehat{\left(\frac{\partial f}{\partial x_j}\right)}(\xi) = 2\pi i \xi_j \widehat{f}(\xi)$ , if  $\frac{\partial f}{\partial x_j} \in L^1$ .
9.  $\widehat{(-2\pi i x_j f)}(\xi) = \frac{\partial \widehat{f}}{\partial \xi_j}(\xi)$ , if  $x_j f \in L^1$ .

## §1.7 The Schwartz class and tempered distributions

Define the **Schwartz class**

$$\mathcal{S}(\mathbb{R}^n) := \left\{ f \in C^\infty(\mathbb{R}^n) : p_{\alpha\beta}(f) = \sup_{x \in \mathbb{R}^n} |x^\alpha D^\beta f|, \forall \alpha, \beta \in \mathbb{N}^n \right\}.$$

Then  $C_c^\infty(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n)$ ,  $e^{-|x|^2} \in \mathcal{S}(\mathbb{R}^n)$ . Moreover  $\mathcal{S}(\mathbb{R}^n) \subset L^p(\mathbb{R}^n)$  and is dense in  $L^p(\mathbb{R}^n)$  ( $1 \leq p < \infty$ ). The topology on  $\mathcal{S}$  is defined as

$$f_k \rightarrow f \text{ in } \mathcal{S} \iff \lim_{k \rightarrow \infty} p_{\alpha,\beta}(f_k - f) = 0, \forall \alpha, \beta \in \mathbb{N}^n.$$

We can give a family of semi-norms on  $\mathcal{S}(\mathbb{R}^n)$  as

$$\|f\|_{(k)} = \sup \{ p_{\alpha,\beta}(f) : \alpha, \beta \in \mathbb{N}^n, |\alpha| + |\beta| \leq k \}$$

and a quasi-norm on  $\mathcal{S}(\mathbb{R}^n)$  as

$$\|f\|_{(*)} = \sum_{k=0}^{\infty} \min \{ \|f\|_{(k)}, 2^{-k} \}.$$

Let  $d(f, g) := \|f - g\|_{(*)}$ , which makes  $\mathcal{S}$  a metric space  $(\mathcal{S}, d)$  and the topology is identified.

### Theorem 1.7.1

The following holds:

1.  $\mathcal{F}(\mathcal{S}) \subset \mathcal{S}$  and  $\mathcal{F} : \mathcal{S} \rightarrow \mathcal{S}$  is continuous.
2.  $\int_{\mathbb{R}^n} f \widehat{g} = \int_{\mathbb{R}^n} \widehat{f} g$ .

### Lemma 1.7.2

If  $f(x) = e^{-\pi|x|^2}$ , then  $\widehat{f}(\xi) = e^{-\pi|\xi|^2}$ .

**Remark 1.7.3** —  $\widehat{e^{-\pi\lambda|x|^2}} = \lambda^{-\frac{n}{2}} e^{-\pi|\xi|^2/\lambda}$  for every  $\lambda \in \mathbb{C}$  with  $\operatorname{Re} \lambda > 0$ .

### Theorem 1.7.4

The following holds:

1. If  $f \in \mathcal{S}$  (or  $f \in L^1$  and  $\widehat{f} \in L^1$ ), then  $f(x) = \int_{\mathbb{R}^n} \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi$ .
2.  $\forall f, g \in \mathcal{S}$ ,  $\int_{\mathbb{R}^n} \widehat{f} \widetilde{g} = \int_{\mathbb{R}^n} f \overline{g}$ .

*Proof.* For  $f \in \mathcal{S}$ , let  $g_\lambda(x) = e^{-\pi\lambda|x|^2}$ , by DCT and the identity

$$\int_{\mathbb{R}^n} \widehat{f}(x) g(\lambda x) dx = \int_{\mathbb{R}^n} f(\lambda x) \widehat{g}(x) dx.$$

□

Let  $\overline{\mathcal{F}}f(\xi) = \int_{\mathbb{R}^n} f(x) e^{2\pi i \xi \cdot x} dx$ ,  $\sigma f(x) = \widetilde{f}(x) = f(-x)$ ,  $Cf(x) = \overline{f(x)}$ . Then  $\overline{\mathcal{F}} = C\mathcal{F}C$ ,  $\overline{\mathcal{F}} = \mathcal{F}^{-1}$ ,  $\mathcal{F}^4 = \operatorname{Id}$ .

**Corollary 1.7.5** (Plancherel)

$$\|f\|_2 = \|\mathcal{F}f\|_2, \forall f \in \mathcal{S}.$$

We define the family of **tempered distributions**  $\mathcal{S}'$  as the continuous linear function on  $\mathcal{S}$ . Then  $T \in \mathcal{S}'$  if and only if  $\exists m \in \mathbb{N}$ , such that  $|\langle T, f \rangle| \leq C \|f\|_{(m)}$  for every  $f \in \mathcal{S}$ . For every  $1 \leq p \leq \infty$ , we have a natural embedding  $j_p : L^p \hookrightarrow \mathcal{S}'$ .

**Definition 1.7.6.**  $\forall T \in \mathcal{S}'$ , define  $\widehat{T}(f) = T(\widehat{f}), \forall f \in \mathcal{S}$ .

Let  $\mathcal{F}_1 : T \mapsto \widehat{T}$ . Then  $\mathcal{F}_1$  maps  $\mathcal{S}'$  to  $\mathcal{S}'$  is continuous. Moreover,  $\mathcal{F}_1 \circ j_1 = j_\infty \circ \mathcal{F}$ .

**Proposition 1.7.7**

If  $T \in \mathcal{S}'$ ,  $\widehat{T} \in L^1$ , then  $T(x) = \int_{\mathbb{R}^n} \widehat{T}(\xi) e^{2\pi i \xi \cdot x} d\xi$  a.e. .

**§1.8 The Fourier transform on  $L^p, 1 < p \leq 2$** **Theorem 1.8.1**

For  $\forall f \in L^2(\mathbb{R}^n)$ , then  $\widehat{f} \in L^2$  and  $\|\widehat{f}\|_2 = \|f\|_2$ .

**Theorem 1.8.2**

It holds  $\widehat{f}(\xi) = \lim_{R \rightarrow \infty} \int_{|x| < R} f(x) e^{-2\pi i \xi \cdot x} dx$ ,  $f(x) = \lim_{R \rightarrow \infty} \int_{|\xi| < R} \widehat{f}(\xi) e^{-2\pi i \xi \cdot x} d\xi$ , both convergences is in the sense of  $L^2$  norm.

Because  $\mathcal{F} : L^1 \rightarrow L^\infty, L^2 \rightarrow L^2$ , then by  $L^p \subset L^1 + L^2$  for  $1 < p < 2$ , we have  $\mathcal{F} : L^p \rightarrow L^1 + L^\infty$ .

**Theorem 1.8.3** (Riesz-Thorin Interpolation Theorem)

$p_0, p_1, q_0, q_1 \in [1, \infty], 0 < \theta < 1$ , let  $\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}$ . If  $T : L^{p_0} + L^{p_1} \rightarrow L^{q_0} + L^{q_1}$  such that  $\|Tf\|_{q_0} \leq M_0 \|f\|_{p_0}, \|Tf\|_{q_1} \leq M_1 \|f\|_{p_1}$ , then  $\|Tf\|_q \leq M_0^{1-\theta} M_1^\theta \|f\|_p$ .

**Corollary 1.8.4**

If  $f \in L^p, 1 \leq p \leq 2$ , then  $\mathcal{F}f \in L^{p'}$  and  $\|\mathcal{F}f\|_{p'} \leq \|f\|_p$ .

**Corollary 1.8.5**

$f \in L^p, g \in L^q, p, q, r \in [1, \infty]$  with  $1 + \frac{1}{r} = \frac{1}{p} + \frac{1}{q}$ , then  $\|f * g\|_r \leq \|f\|_p \|g\|_q$ .



## §1.9 The convergence and summability of Fourier integral

Let  $B_R = R \cdot B$  where  $B$  is a neighborhood of origin.

**Question 1.9.1.**  $f(x) = \lim_{R \rightarrow \infty} \int_{B_R} \widehat{f}(\xi) e^{2\pi i \xi \cdot x} d\xi$ ?

Let  $\widehat{S_R f} = \chi_{B_R} \widehat{f}$ , then  $\lim_{R \rightarrow \infty} \|S_R f - f\|_p = 0$  iff  $\|S_R f\|_p \leq C_p \|f\|_p$ .

**Fact 1.9.2.**  $S_R : L^p \rightarrow L^p$  bounded iff  $n = 1, 1 < p < \infty$  or  $n = 1, p = 2 (B = B(0, 1))$  or  $n > 1, 1 < p < \infty (B = Q(0, 1))$ .

$n = 1, B = (-1, 1)$ , then  $S_R f = D_R * f$ , where  $D_R$  is the Dirichlet kernel

$$D_R(x) = \int_{-R}^R e^{2\pi i \xi \cdot x} d\xi = \frac{\sin(2\pi R x)}{R x}.$$

Then  $D_R \notin L^1$  but  $D_R \in L^q (1 < q \leq \infty)$ .

**Almost everywhere convergence** Now we consider the almost everywhere convergence, an argument (Carleson-Hunt) shows that

$$\left\| \sup_R |S_R f|_p \right\| \leq C_p \|f\|_p \implies \lim_{R \rightarrow \infty} S_R f(x) = f(x) \text{ a.e. }, \forall f \in L^p, 1 < p < \infty.$$

**Cesàro sum** Let  $\sigma_R f(x) = \frac{1}{R} \int_0^R S_t f(x) dt = F_R * f(x)$ , where  $F_R$  is the Fejér kernel

$$F_R(x) = \frac{1}{R} \int_0^R D_t(x) dt = \frac{\sin^2(\pi R x)}{R(\pi x)^2}.$$

Then  $F_R \in L^1$  and  $F_R \geq 0$ . We have  $\lim_{R \rightarrow \infty} \|\sigma_R f - f\|_p = 0 \forall p \in [1, \infty)$  and  $\sigma_R f \rightarrow f$  a.e. .

**Abel-Poisson sum** Let  $u(x, t) = \int_{\mathbb{R}^n} e^{-2\pi t |\xi|} \widehat{f}(\xi) e^{-2\pi i \xi \cdot x} d\xi = P_t * f(x)$ , where

$$P_t(x) = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\pi^{\frac{n+1}{2}}} \frac{t}{(t^2 + |x|^2)^{\frac{n+1}{2}}}, \quad \widehat{P}_t(\xi) = e^{-2\pi t |\xi|}.$$

We have  $\Delta_{t,x} P_t(x) = 0$ , then  $\Delta u = 0$  on  $\mathbb{R}_+ \times \mathbb{R}^n$ . We also have  $\lim_{t \rightarrow 0+} u(x, t) = f(x)$  a.e. ,  $\forall f \in L^p(\mathbb{R}^n)$ .

Conversely, if  $\Delta u = 0$  in  $\mathbb{R}_+^{n+1}$ ,  $\sup_{t>0} \int_{\mathbb{R}^n} |u(x, t)|^p dx < \infty, 1 < p \leq \infty$ . Then  $\exists f \in L^p(\mathbb{R}^n)$  such that  $u(x, t) = P_t * f(x)$ .

**Gauss sum** Let  $w(x, t) = \int_{\mathbb{R}^n} e^{-2\pi t^2 |\xi|^2} \widehat{f}(\xi) e^{-2\pi i \xi \cdot x} d\xi = W_t * f(x)$ , where  $W_t$  is the **Gauss kernel**  $W_t := \mathcal{F}(e^{-\pi t^2 |\xi|^2}) = E^n e^{-\pi |x|^2/t}$ . Let  $\widetilde{W}(x, t) = W(x, \sqrt{4\pi t})$ , then  $\frac{\partial \widetilde{W}}{\partial t} - \Delta \widetilde{W} = 0$  in  $\mathbb{R}_+^{n+1}$ . We have  $\lim_{t \rightarrow 0+} \widetilde{W}(x, t) = \lim_{t \rightarrow 0+} W(x, t) = f(x)$  a.e. ,  $\forall f \in L^p(\mathbb{R}^n)$ .

# 2 The Hardy-Littlewood Maximal Function

## §2.1 Approximations of the identity

$\phi \in L^1(\mathbb{R}^n)$ ,  $\int \phi = 1$ . For  $t > 0$ , let  $\phi_t = t^{-n}\phi(t^{-1}x)$ . Then  $\phi_t \rightarrow \delta(t \rightarrow 0)$  in  $\mathcal{S}'$ , hence  $\phi_t * g \rightarrow g(t \rightarrow 0)$ .

**Example 2.1.1** (Cesàro sum)

$\phi = F_1 = \frac{\sin^2 \pi x}{(\pi x)^2}$ , then  $F_R = \phi_{1/R}$ .

**Example 2.1.2** (Poisson kernel)

$\phi = P_1 = \frac{\Gamma(\frac{n+1}{2})}{\pi^{\frac{n+1}{2}}} \frac{1}{(1+|x|^2)^{\frac{n+1}{2}}}$ , then  $P_t = \phi_t$ .

**Example 2.1.3** (Gauss kernel)

$\phi = W_1 = e^{-\pi|x|^2}$ , then  $W_t = \phi_t$ .

**Theorem 2.1.4**

$\int_{\mathbb{R}^n} \phi = A$ ,  $f \in L^p$ ,  $1 \leq p < \infty$  or  $p = \infty$ ,  $f \in C_0(\mathbb{R}^n)$ , then  $\lim_{t \rightarrow 0+} \|\phi_t * f - Af\|_p \rightarrow 0$ .

**Remark 2.1.5** — Then  $\exists \{t_k\} \rightarrow 0$  such that  $\phi_{t_k} * f \rightarrow f$  a.e. . Hence,

$$\left| \left\{ x : \lim_{t \rightarrow 0} \phi_t * f(x) \text{ exists but not equal to } f(x) \right\} \right| = 0.$$

## §2.2 Weak-type inequalities and almost everywhere convergence

$(X, \mu), (Y, \nu)$  measure spaces.  $T : L^p(X, \mu) \rightarrow m(Y, \nu)$  the measurable functions on  $Y$ .

**Definition 2.2.1.** We say  $T$  is **weak**  $(p, q)$ ,  $q < \infty$  if  $\exists C > 0$ , such that

$$\nu(\{y \in Y : |Tf(y)| > \lambda\}) \leq \left( \frac{C \|f\|_p}{\lambda} \right)^q, \quad \forall \lambda > 0.$$

Recall strong  $(p, q)$ , strong  $(p, \infty)$ , and we define weak  $(p, \infty)$  same as strong  $(p, \infty)$ .

**Definition 2.2.2.** Define the weak space as

$$L^{p,\infty}(Y, \nu) := \left\{ f \in m(Y, \nu) : \|f\|_{p,\infty} := \sup_{\lambda > 0} \lambda \nu(|Tf| > \lambda)^{\frac{1}{p}} < \infty \right\}.$$

Then weak  $(p, q)$  means  $\|Tf\|_{q, \infty} \leq C \|f\|_p$ . Besides, there holds  $\|f\|_{p, \infty} \leq \|f\|_p$ .

### Theorem 2.2.3

$\{T_t\}$  are linear operators on  $L^p(X, \mu)$ , let  $T^*f(x) = \sup_t |T_t f(x)|$ . If  $T^*$  is weak  $(p, q)$ , then

$$V := \left\{ f \in L^p(X, \mu) : \lim_{t \rightarrow 0} T_t(x) = f(x) \text{ a.e. } \right\}$$

is closed in  $L^p(X, \mu)$ .

**Remark 2.2.4** — There are something ambiguous in the theorem. One should notice that the definition of  $T^*$  do **not** guarantee the measurability of  $T^*f$ . Besides, if  $f = g$  a.e., there still might be  $T^*f \neq T^*g$  on a set with positive measure.

If  $\phi \in L^1$ ,  $\int \phi = 1$ , let  $T_t f = \phi_t * f$ , then  $\mathcal{S} \subseteq V$ . Then for  $1 \leq p < \infty$ , if we can prove  $\sup_{t>0} |T_t f|$  is weak  $(p, q)$ , then  $V = L^p$ .

## §2.3 The Marcinkiewicz interpolation theorem

$f : X \rightarrow \mathbb{C}$  measurable, define  $a_f(\lambda) = \mu \{x \in X : |f(x)| > \lambda\}$ ,  $\forall \lambda > 0$ .

### Proposition 2.3.1

$\phi : [0, \infty) \rightarrow [0, \infty)$   $C^1$  increasing,  $\phi(0) = 0$ , then

$$\int_X \phi(|f(x)|) d\mu = \int_0^\infty \phi'(\lambda) a_f(\lambda) d\lambda.$$

### Theorem 2.3.2 (Marcinkiewicz Interpolation Theorem)

$1 \leq p_0 < p_1 \leq \infty$ ,  $T : L^{p_0}(X, \mu) + L^{p_1}(X, \mu) \rightarrow L^{p_0, \infty}(Y, \nu) + L^{p_1, \infty}(Y, \nu)$  sub-linear. If  $T$  is weak  $(p_0, p_0)$  and weak  $(p_1, p_1)$ , then  $T$  is strong  $(p, p)$  for every  $p_0 < p < p_1$ .

### Theorem 2.3.3

$T : L^p(X, \mu)$  weak  $(p, q)$ , then  $\{f : Tf = 0 \text{ a.e.}\}$  is closed in  $L^p$ .

## §2.4 The Hardy-Littlewood maximal function

Let  $B_r = \{x \in \mathbb{R}^n : |x| < r\}$ , for every  $f \in L^1_{loc}(\mathbb{R}^n)$ , define

$$Mf(x) = \sup_{r>0} \frac{1}{|B_r|} \int_{B_r} |f(x-y)| dy = \sup_{t>0} \phi_t * |f|(x), \quad \phi = \frac{1}{|B_1|} \mathbb{1}_{B_1}.$$

Besides, for  $Q = [-r, r]^n$ ,  $|Q_r| = (2r)^n$ , define

$$M'f(x) = \sup_{r>0} \frac{1}{(2r)^n} \int_{Q_r} |f(x-y)| dy = \sup_{t>0} \phi_t * |f|(x), \quad \phi = \frac{1}{2^n} \mathbb{1}_{Q_1}.$$

Then, if  $n = 1$ , we have  $M = M'$ . For  $n \geq 2$ , we have some controlling  $c_n M' f \leq M f \leq C_n M' f$ . We can also define

$$M'' f(x) = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y)| dy, \quad Q \text{ is a box.}$$

Then  $M' f \leq M'' f \leq 2^n M' f$ .

#### Lemma 2.4.1

Let  $\mathcal{F} = \{B_j = B(x_j, r_j)\}_{j=1}^N$  be open balls in metric space  $(X, d)$ . Let  $mB_j = B(x_j, mr_j)$ . Then there exists  $\{B'_i\}_{i=1}^l \subseteq \mathcal{F}$  such that  $B'_i \cap B'_j = \emptyset$  for every  $i \neq j$ , and  $\bigcup_{j=1}^N B_j \subseteq \bigcup_{i=1}^l 3B'_i$ .

#### Theorem 2.4.2

$$\|Mf\|_{1,\infty} \leq 3^n \|f\|_1.$$

**Remark 2.4.3** — This estimate also holds for  $M', M'', \widetilde{M}$ .

Define the space of non-negative radial decreasing function

$$\nu_0(\mathbb{R}^n) = \{\phi(x) = \phi_0(|x|) : \phi_0 : (0, \infty) \rightarrow [0, \infty) \text{ decreasing}, \phi \in L^1(\mathbb{R}^n)\}.$$

#### Proposition 2.4.4

If  $\phi \in \nu_0(\mathbb{R}^n)$ , then  $\sup_{t>0} |\phi_t * f(x)| \leq \|\phi\|_1 Mf(x), \forall f \in L^1_{loc}(\mathbb{R}^n)$ .

#### Corollary 2.4.5

If  $|\phi(x)| \leq \psi(x) \in \nu_0(\mathbb{R}^n)$ , then  $f \mapsto \sup_{t>0} \phi_t * f$  is weak  $(1, 1)$ .

Define  $\nu_1(\mathbb{R}^n) := \{\phi \in L^1(\mathbb{R}^n) : \exists \psi \in \nu_0(\mathbb{R}^n), |\phi| \leq \psi \text{ a.e.}\}.$

#### Corollary 2.4.6

If  $1 \leq p \leq \infty$ ,  $f \in L^p(\mathbb{R}^n)$ ,  $\phi \in \nu_1(\mathbb{R}^n)$ , then  $\lim_{t \rightarrow 0+} \phi_t * f(x) = \left(\int_{\mathbb{R}^n} \phi\right) f(x)$  a.e. .

Recall Poisson kernel  $P_1$ , Gauss kernel  $W_1$ , Fejér kernel  $F_1$ . Then

$$P_1, W_1 \in \nu_0(\mathbb{R}^n), \quad F_1 \in \nu_1(\mathbb{R}).$$

#### Corollary 2.4.7

If  $f \in L^1_{loc}(\mathbb{R}^n)$ , then  $\lim_{r \rightarrow 0+} \frac{1}{|B_r|} \int_{B_r} f(x-y) dy = f(x)$  a.e. . Moreover,

$$\lim_{r \rightarrow 0+} \frac{1}{|B_r|} \int_{B_r} |f(x-y) - f(x)| dy = 0 \text{ a.e. .}$$

**Lemma 2.4.8**

$f \in L^1_{loc}(\mathbb{R}^n)$ ,  $f \neq 0$ , then  $Mf \notin L^1(\mathbb{R}^n)$ .

If  $Mf(x_0) = 0$  for some  $x_0 \in \mathbb{R}^n$ , then  $f(x) = 0$  a.e. .

**Theorem 2.4.9**

$B \subset \mathbb{R}^n$  bounded, then  $\exists C > 0$  such that

$$\int_B Mf \leq 2|B| + C \int_{\mathbb{R}^n} |f| \ln^+ |f| dx.$$

*Proof.* Note that

$$\int_B Mf = 2 \int_0^\infty |\{x \in B : Mf(x) \geq 2\lambda\}| d\lambda \leq 2|B| + \int_1^\infty |\{x \in B : Mf(x) \geq 2\lambda\}| d\lambda.$$

Let  $f = f_1 + f_2$ , where  $f_1 = f \mathbb{1}_{\{x: |f(x)| > \lambda\}}$ ,  $f_2 = f - f_1$ , then  $|Mf_2| \leq \lambda$ . Hence

$$|\{x \in B : Mf(x) > 2\lambda\}| \leq |\{x \in B : Mf_1(x) > \lambda\}| \leq \frac{C}{\lambda} \int_{\{x: |f(x)| > \lambda\}} |f|.$$

Bring back to the integral and apply Fubini theorem.  $\square$

**Theorem 2.4.10**

Let  $w \geq 0$ ,  $w \in L^1_{loc}(\mathbb{R}^n)$ , then

$$\begin{aligned} \int_{\mathbb{R}^n} Mf(x)^p w(x) dx &\leq C_p \int_{\mathbb{R}^n} |f(x)|^p dx. \\ \int_{\{x: Mf(x) > \lambda\}} w(x) dx &\leq \frac{C}{\lambda} \int_{\mathbb{R}^n} |f(x)| Mw(x) dx. \end{aligned}$$

*Proof.* WLOG,  $w \in L^1(\mathbb{R}^n)$ , consider  $(X, \mu) = (\mathbb{R}^n, w(x)dx)$ ,  $(Y, \nu) = (\mathbb{R}^n, Mw(x)dx)$ . For  $w \neq 0$ , we have estimate  $Mw(x) \geq \frac{c}{1+|x|^n}$ , hence  $M : X \rightarrow Y$  is  $(\infty, \infty)$ . It suffices to show  $M$  is weak  $(1, 1)$ . Let  $E_\lambda = \{x : Mf(x) \geq \lambda\}$ , for all compact  $K \subset E_\lambda$ , we have  $K \subseteq \bigcup_{j=1}^N 3B_j$  for some  $B_j$ . We claim that

$$\int_{3B_j} w(x) dx \leq \frac{4^n}{\lambda} \int_{B_j} |f(x)| Mw(x) dx.$$

It suffices to show that  $4^n |B_j| \inf_{B_j} Mw(x) \geq \int_{3B_j} w(x) dx$ . Assume the radius of  $B_j$  is  $r_j$ , then for every  $y \in B_j$ , we have  $B(y, 4r_j) \supseteq 3B_j$ . Hence

$$Mw(y) \geq \frac{1}{4^n |B_j|} \int_{B(y, 4r_j)} w(z) dz \geq \frac{1}{4^n |B_j|} \int_{3B_j} w(x) dx.$$

$\square$

Let  $\widetilde{F}_N = F_N(t) \mathbb{1}_{\{-\frac{1}{2}, \frac{1}{2}\}}$ ,  $\widetilde{P}_r(t) = P_r(t) \mathbb{1}_{\{-\frac{1}{2}, \frac{1}{2}\}}$ .

**Lemma 2.4.11**

If  $f \in L^p(\mathbb{T})$ ,  $1 \leq p \leq \infty$ , then

$$\lim_{n \rightarrow \infty} \sigma_n f(x) = f(x) \text{ a.e. }, \quad \lim_{r \rightarrow 1^-} P_r * f(x) = f(x) \text{ a.e. }.$$

*Proof.* Let  $\Omega_1 f = \limsup_{n \rightarrow \infty} |\sigma_n f(x) - f(x)|$ ,  $\Omega_2 f(x) = \limsup_{r \rightarrow 1^-} |P_r * f(x) - f(x)|$ , then  $\Omega_1 f = \Omega_2 f = 0$  for every  $f \in C(\mathbb{T})$ . Note that  $\Omega_1, \Omega_2$  are both weak  $(1, 1)$  by the estimate of convolution. Hence  $\Omega_1 f, \Omega_2 f = 0$  a.e.  $\square$

**§2.5 The dyadic maximal function**

Let

$$\mathcal{Q}_k := \left\{ \prod_{i=1}^n \left[ \frac{a_i}{2^k}, \frac{a_i+1}{2^k} \right) : a_1, a_2, \dots, a_n \in \mathbb{Z} \right\}, \quad \mathcal{Q}_* = \bigcup_k \mathcal{Q}_k.$$

Then

1.  $\forall x \in \mathbb{R}, k \in \mathbb{Z}$ , there exists unique  $Q \in \mathcal{Q}_k$  such that  $x \in Q$ .
2.  $\forall A, B \in \mathcal{Q}_*$ , then  $A \cap B = \emptyset$  or  $A \subseteq B$  or  $B \subseteq A$ .
3.  $\forall A \in \mathcal{Q}_k, j < k$ , there exists unique  $B \in \mathcal{Q}_j$  such that  $A \subseteq B$ .

For all  $f \in L^1_{loc}(\mathbb{R}^n)$ , define

$$E_k f = \sum_{Q \in \mathcal{Q}_k} \left( \frac{1}{|Q|} \int_Q f \right) \mathbb{1}_Q = \mathbb{E}(f | \sigma(\mathcal{Q}_k)).$$

**Definition 2.5.1.** Define the **dyadic maximal function**  $M_d f(x) = \sup_k |E_k f(x)|$ .

**Remark 2.5.2** — Note that in the definition of dyadic maximal function, it does not take the absolute value of  $f$  directly. Hence, we do **not** have  $M_d f = M_d |f|$  in general.

**Theorem 2.5.3**

The following holds:

1.  $M_d$  is weak  $(1, 1)$ . Moreover,  $\|M_d f\|_{1, \infty} \leq \|f\|_1$ .
2. If  $f \in L^1_{loc}$ , then  $E_k f \rightarrow f$  a.e. ( $k \rightarrow +\infty$ ).

*Proof.* For  $f \geq 0$ , for every  $\lambda > 0$ , let

$$E_\lambda := \{x \in \mathbb{R}^n : M_d f(x) > \lambda\},$$

then  $E_\lambda = \bigcup_{k \in \mathbb{Z}} \Omega_k$ , where

$$\Omega_k := \{x \in \mathbb{R}^n : E_k f(x) > \lambda, \quad E_j f \leq \lambda, \forall j < k\}.$$

Then  $\Omega_k$  forms a disjoint union of  $E_\lambda$ . The following proof is easy.  $\square$

**Theorem 2.5.4** (Calderón-Zygmund Decomposition)

$\forall f \in L^1(\mathbb{R}^n)$ ,  $f \geq 0$ ,  $\lambda > 0$ .  $\exists \{A_j\} \subset \mathcal{Q}_*$ ,  $A_i \cap A_j = \emptyset$  for every  $i \neq j$  such that

- (i)  $f(x) \leq \lambda$  for almost every  $x \notin \bigcup_j A_j$ .
- (ii)  $\left| \bigcup_j A_j \right| \leq \frac{1}{\lambda} \|f\|_1$ .
- (iii)  $\lambda < \frac{1}{|A_j|} \int_{A_j} f \leq 2^n \lambda$  for every  $A_j$ .

**Remark 2.5.5** — This theorem also gives somehow “reverse” weak  $(1,1)$ , that is

$$|E_\lambda| \geq \frac{1}{2^n \lambda} \int_{E_\lambda} f.$$

For  $f$  with  $\text{supp } f \subseteq Q \in \mathcal{Q}_*$ ,  $f \geq 0$ , we have

$$\int_Q M_d f = \int_0^\infty |E_\lambda \cap Q| d\lambda \leq |Q| + \int_Q f \ln^+ M_d f.$$

Note that  $B \ln^+ A \leq B \ln^+ B + \frac{A}{e}$ . Take  $A = M_d f$  and  $B = f$  we have

$$(1 - e^{-1}) \int_Q M_d f \leq |Q| + \int_Q f \ln^+ f.$$

Conversely, assume  $|Q| = 1$  and  $\|f\|_1 = 1$ , then  $E_k f(x) = 0$  for every  $x \notin Q$  and  $k \geq 0$ . Moreover,  $E_k f(x) \leq 2^{nk} \leq 2^{-n}$  for  $x \notin Q$  and  $k \leq -1$ . Let  $\lambda_0 = 2^{-n}$ , then  $E_\lambda \subseteq Q$  for every  $\lambda > \lambda_0$ . We have

$$\int_Q M_d f(x) dx \geq \int_{\lambda_0}^\infty |E_\lambda| d\lambda \geq \frac{1}{2^n} \int_Q f(x) \ln^+ \frac{f(x)}{\lambda_0}.$$

**Proposition 2.5.6**

$M_d f \in L^1(Q)$  if and only if  $f \in L \ln L(Q)$ .

**§2.6 Some other covering lemmas****Theorem 2.6.1** (Vitali)

$\{B_j\}_{j \in \mathcal{J}}$  be a family of open balls in  $\mathbb{R}^n$ , then there exists countable disjoint  $\{B_k\} \subseteq \{B_j\}_{j \in \mathcal{J}}$  such that  $\bigcup_{j \in \mathcal{J}} B_j \subseteq_k 5B_k$ .

**Theorem 2.6.2** (Besicovitch)

$A \subseteq \mathbb{R}^n$  bounded,  $\mathcal{F} = \{B_x\}_{x \in A}$  where  $B_x = B(x, r_x)$ . Then there exists countable  $\{B_j\} \subseteq \mathcal{F}$  such that

$$A \subseteq \mathcal{F}, \quad \sum_j \mathbb{1}_{B_j}(x) \leq C_n.$$

Let  $\mu$  be a Radon measure on  $\mathbb{R}^n$ , define

$$M_\mu f(x) = \sup_{r>0} \frac{1}{\mu(B(x,r))} \int_{B(x,r)} |f(y)| d\mu(y).$$

**Corollary 2.6.3**

$$\|M_\mu f\|_{1,\infty} \leq C_n \|f\|_{L^1(\mu)}.$$

**Remark 2.6.4** — Note that the definition of  $M_\mu$  is **different** with

$$M'_\mu f(x) := \sup_{r>0} \frac{1}{\mu(B_r)} \int_{B_r} |f(x-y)| d\mu(y).$$

Because  $\mu$  may not be invariant under translation.  $M'_\mu$  is not weak  $(1,1)$  in measure  $\mu$ . Consider  $X = \mathbb{R}$ ,  $d\mu = e^{|x|} dx$ ,  $f = \mathbb{1}_{\{-1,1\}}$ . Then  $M'_\mu f(x) = \frac{1-e^{-2}}{2} \notin L^{1,\infty}(\mu)$ .