

1 Combinatorial

2 Permutations

2.1 Cycles

Let $g_S(n)$ be the number of n -permutations whose cycle lengths all belong to the set S . Then

$$\sum_{n=0}^{\infty} g_S(n) \frac{x^n}{n!} = \exp \left(\sum_{n \in S} \frac{x^n}{n} \right)$$

2.2 Derangements

Permutations of a set such that none of the elements appear in their original position.

$$D(n) = (n-1)(D(n-1) + D(n-2)) = nD(n-1) + (-1)^n = \left\lfloor \frac{n!}{e} \right\rfloor$$

2.3 Burnside's lemma

Given a group G of symmetries and a set X , the number of elements of X up to symmetry equals

$$\frac{1}{|G|} \sum_{g \in G} |X^g|,$$

where X^g are the elements fixed by g ($g.x = x$).

If $f(n)$ counts “configurations” (of some sort) of length n , we can ignore rotational symmetry using $G = \mathbb{Z}_n$ to get

$$g(n) = \frac{1}{n} \sum_{k=0}^{n-1} f(\gcd(n, k)) = \frac{1}{n} \sum_{k|n} f(k) \phi(n/k).$$

3 Partitions and subsets

3.1 Partition function

Number of ways of writing n as a sum of positive integers, disregarding the order of the summands.

$$p(0) = 1, \quad p(n) = \sum_{k \in \mathbb{Z} \setminus \{0\}} (-1)^{k+1} p(n - k(3k-1)/2)$$

$$p(n) \sim 0.145/n \cdot \exp(2.56\sqrt{n})$$

$$\begin{array}{c|ccccccccccccc} n & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 20 & 50 & 100 \\ \hline p(n) & 1 & 1 & 2 & 3 & 5 & 7 & 11 & 15 & 22 & 30 & 627 & \sim 2e5 & \sim 2e8 \end{array}$$

3.2 Lucas' Theorem

Let n, m be non-negative integers and p a prime. Write $n = n_k p^k + \dots + n_1 p + n_0$ and $m = m_k p^k + \dots + m_1 p + m_0$. Then $\binom{n}{m} \equiv \prod_{i=0}^k \binom{n_i}{m_i} \pmod{p}$.

4 General purpose numbers

4.1 Bernoulli numbers

EGF of Bernoulli numbers is $B(t) = \frac{t}{e^t - 1}$ (FFT-able). $B[0, \dots] = [1, -\frac{1}{2}, \frac{1}{6}, 0, -\frac{1}{30}, 0, \frac{1}{42}, \dots]$
Sums of powers:

$$\sum_{i=1}^n i^m = \frac{1}{m+1} \sum_{k=0}^m \binom{m+1}{k} B_k \cdot (n+1)^{m+1-k}$$

Euler-Maclaurin formula for infinite sums:

$$\begin{aligned} \sum_{i=m}^{\infty} f(i) &= \int_m^{\infty} f(x) dx - \sum_{k=1}^{\infty} \frac{B_k}{k!} f^{(k-1)}(m) \\ &\approx \int_m^{\infty} f(x) dx + \frac{f(m)}{2} - \frac{f'(m)}{12} + \frac{f''(m)}{720} + O(f^{(5)}(m)) \end{aligned}$$

4.2 Stirling numbers of the first kind

Number of permutations on n items with k cycles.

$$c(n, k) = c(n - 1, k - 1) + (n - 1)c(n - 1, k), \quad c(0, 0) = 1$$

$$\sum_{k=0}^n c(n, k)x^k = x(x + 1) \dots (x + n - 1)$$

$$c(8, k) = 8, 0, 5040, 13068, 13132, 6769, 1960, 322, 28, 1$$

$$c(n, 2) = 0, 0, 1, 3, 11, 50, 274, 1764, 13068, 109584, \dots$$

4.3 Eulerian numbers

Number of permutations $\pi \in S_n$ in which exactly k elements are greater than the previous element. k j :s s.t. $\pi(j) > \pi(j + 1)$, $k + 1$ j :s s.t. $\pi(j) \geq j$, k j :s s.t. $\pi(j) > j$.

$$E(n, k) = (n - k)E(n - 1, k - 1) + (k + 1)E(n - 1, k)$$

$$E(n, 0) = E(n, n - 1) = 1$$

$$E(n, k) = \sum_{j=0}^k (-1)^j \binom{n+1}{j} (k+1-j)^n$$

4.4 Stirling numbers of the second kind

Partitions of n distinct elements into exactly k groups.

$$S(n, k) = S(n - 1, k - 1) + kS(n - 1, k)$$

$$S(n, 1) = S(n, n) = 1$$

$$S(n, k) = \frac{1}{k!} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} j^n$$

4.5 Bell numbers

Total number of partitions of n distinct elements. $B(n) = 1, 1, 2, 5, 15, 52, 203, 877, 4140, 21147, \dots$. For p prime,

$$B(p^m + n) \equiv mB(n) + B(n + 1) \pmod{p}$$

4.6 Labeled unrooted trees

on n vertices: n^{n-2}

on k existing trees of size n_i : $n_1 n_2 \dots n_k n^{k-2}$

with degrees d_i : $(n - 2)! / ((d_1 - 1)! \dots (d_n - 1)!)$

4.7 Catalan numbers

$$C_n = \frac{1}{n+1} \binom{2n}{n} = \binom{2n}{n} - \binom{2n}{n+1} = \frac{(2n)!}{(n+1)!n!}$$

$$C_0 = 1, \quad C_{n+1} = \frac{2(2n+1)}{n+2} C_n, \quad C_{n+1} = \sum C_i C_{n-i}$$

$$C_n = 1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, \dots$$

- sub-diagonal monotone paths in an $n \times n$ grid.
- strings with n pairs of parenthesis, correctly nested.
- binary trees with $n + 1$ leaves (0 or 2 children).
- ordered trees with $n + 1$ vertices.
- ways a convex polygon with $n + 2$ sides can be cut into triangles by connecting vertices with straight lines.
- permutations of $[n]$ with no 3-term increasing subseq.