

An independent four-generator counterexample; this is not the cubic model.

DGLA $\mathfrak{g}$

Degree one: α, ξ, E ; degree two: β

$$d\alpha = d\xi = 0, \quad dE = \beta$$

$$[\xi, \xi] = 2\beta, \quad [\alpha, E] = 2\beta$$

p

Abelian DGLA $\mathfrak{h}$

Degree one: α, ξ

Zero differential and zero brackets

$$p(E) = p(\beta) = 0$$

p fixes α, ξ and has acyclic kernel $\mathbb{C}E \xrightarrow{d} \mathbb{C}\beta$.

Hence p is a strict DGLA quasi-isomorphism.

$$\gamma = u\alpha + v\xi + wE$$

$$w + v^2 + 2uw = 0$$

$$w = -v^2/(1 + 2u)$$

$$\text{MC}(\mathfrak{h}) : u\alpha + v\xi$$

Same smooth formal germ
in two parameters

$$\mathbb{C}[[u, v]]$$

Over local Artin algebras, $1 + 2u$ is a unit; there are no degree-zero gauge directions.

$$\text{In } \mathfrak{g}: \mathcal{L}_{\text{res}} = \mathbb{C}\beta, T_{\text{resp}} = \mathbb{C}\alpha, \rho_{\text{res}}(\alpha) = 2.$$

$$\text{In } \mathfrak{h}: \mathcal{L}'_{\text{res}} = 0.$$

The Maurer–Cartan functor is preserved, but the marked pair is erased.