

No assumption that $K = \ker H$ is invariant

$$U = \bigcap_{j=0}^{n-1} \ker(HA^j) \subseteq K \subseteq R, \quad n = \dim R > 0$$

U is the largest A -invariant subspace contained in K .



The full orbit is unchanged: $A_\tau = A + \tau H$

Every A_τ has the same U and the same action $A|_U$.

Choose $R = U \oplus X$. On R/U , the blocks range over $B + LC$;
the induced pair (B, C) is observable.



First quotient witness

$$\begin{aligned} \chi_{A_{\tau_1}} &= \chi_{A|_U} p_1 \\ \gcd(p_1, \chi_{A|_U}) &= 1 \end{aligned}$$



Second quotient witness

$$\begin{aligned} \chi_{A_{\tau_2}} &= \chi_{A|_U} p_2 \\ \gcd(p_2, \chi_{A|_U} p_1) &= 1 \end{aligned}$$



Stable-core characteristic and minimal floors

Coprime quotient spectra split the extension for these two witnesses:

$$\begin{aligned} \mu_{A_{\tau_i}} &= \mu_{A|_U} \mu_{B+L_i C} \quad (i = 1, 2). \\ \gcd_{\tau} \chi_{A_{\tau}} &= \chi_{A|_U}, \quad \gcd_{\tau} \mu_{A_{\tau}} = \mu_{A|_U} \end{aligned}$$

Over an infinite field, spectral avoidance supplies the two quotient blocks.

The perturbations still vanish on K , not merely on U .

If K is invariant, $U = K$; zero-dimensional polynomials are 1.