

Static extraction from finite landing data

$$\begin{aligned} M &\xrightarrow{\Lambda} R_\Lambda \xrightarrow{D} F \xrightarrow{q} F/N, & \Gamma &= qD\Lambda \\ K_\Lambda &= R_\Lambda \cap D^{-1}(N) &= \Lambda(\ker \Gamma) &\simeq \ker \Gamma / \ker \Lambda \end{aligned}$$

Global landing descent

$$\begin{aligned} T \ker \Lambda &\subseteq \ker \Lambda \\ \iff \text{a unique } A_\Lambda \text{ satisfies} \\ A_\Lambda \Lambda &= \Lambda T \end{aligned}$$

Additional normalized closure

$$\begin{aligned} T \ker \Gamma &\subseteq \ker \Gamma \\ \text{Under global descent, this is equivalent} \\ &\text{to invariance of } K_\Lambda. \end{aligned}$$

Stable source kernel, always defined for $m = \dim M > 0$

$$\mathcal{O} = \begin{bmatrix} \Gamma \\ \Gamma T \\ \vdots \\ \Gamma T^{m-1} \end{bmatrix}, \quad W = M_\infty = \ker \mathcal{O}$$

W is the largest T -invariant subspace of $\ker \Gamma$.

Three distinct dynamical outcomes

Global descent $K_\infty = \Lambda W \simeq W / \ker \Lambda$
 Local descent: $TJ \subseteq J$ $\Lambda W \simeq W/J$, $J = W \cap \ker \Lambda$
 No local descent W/J_{fut} retains all future landing outputs

$$\begin{aligned} J_{\text{fut}} &= \bigcap_{j=0}^{m-1} \ker(\Lambda T^j) \subseteq W \\ W/J_{\text{fut}} &\simeq \text{im}[w \mapsto (\Lambda w, \Lambda T w, \dots, \Lambda T^{m-1} w)] \end{aligned}$$

The future-output quotient may be larger than ΛW .
 It does not define static landing dynamics when local descent fails.