

Finite invariant-source closure

$$N^{(0)} = \text{im } R_0, \quad N^{(r+1)} = N^{(r)} + \sum_i C_i N^{(r)}$$

$$N^{(0)} \subseteq N^{(1)} \subseteq \dots \subseteq N_{\text{eq}} = \mathbb{k}\langle C_1, \dots, C_s \rangle \text{im } R_0$$

The sequence stabilizes in finite dimension.



Minimality

N_{eq} is the unique smallest subspace containing the seed image and invariant under every C_i .



Intertwining data imply functoriality

$$\Phi_F R_0 = R'_0 \Phi_W, \quad \Phi_F C_i = C'_i \Phi_F$$

$$\implies \Phi_F(N_{\text{eq}}) \subseteq N'_{\text{eq}}$$



Generated coverage when Φ_F is invertible

$$\Phi_F(N_{\text{eq}}) = N'_{\text{eq}} \iff \text{im } R'_0 \subseteq \Phi_F(N_{\text{eq}})$$

Mapping onto the target seed image is sufficient, but not necessary.



Insertion into the landing problem

$$\text{For } N_{\text{eq}} \subseteq H_0(P_\bullet), \text{ use } N = \iota(N_{\text{eq}}) \subseteq F,$$

$$q_{\text{eq}} : F \rightarrow F/N, \quad \Gamma_{\text{eq}} = q_{\text{eq}} D\Lambda.$$