

Finite marked input and supplied observation data

$$\begin{aligned} \mathcal{D} &= (P_\bullet, \mathfrak{G}_{\text{mark}}, R_0; \iota, F, M, T, \Lambda, D) \\ \iota : H_0(P_\bullet) &\hookrightarrow F, \quad \Lambda : M \rightarrow R, \quad D : R \rightarrow F, \quad T \in \text{End}(M) \end{aligned}$$

Descend the marked chain maps

$$\begin{aligned} dG = Gd &\implies \bar{G} \in \text{End}(H_0) \\ \mathcal{A}_{\text{eq}}^{\text{eff}} &= \mathbb{k}\langle \bar{G} : G \in \mathfrak{G}_{\text{mark}} \rangle \end{aligned}$$

Generate the normalized source

$$\begin{aligned} N_{\text{eq}} &= \mathcal{A}_{\text{eq}}^{\text{eff}} \text{im } R_0 \subseteq H_0 \\ q_{\text{eq}} : F &\rightarrow F/\iota(N_{\text{eq}}) \end{aligned}$$

Normalize the output and compute the stable kernel

$$\Gamma_{\text{eq}} = q_{\text{eq}} D \Lambda, \quad W = \bigcap_{j=0}^{m-1} \ker(\Gamma_{\text{eq}} T^j), \quad m = \dim M > 0$$

Landing descent: $T \ker \Lambda \subseteq \ker \Lambda$

$$K_\infty = \Lambda W \simeq W / \ker \Lambda \text{ carries the induced } T\text{-action.}$$

Its characteristic and minimal polynomials are the two forced divisors of the *original full* normalized tilt orbit on $R_\Lambda = \text{im } \Lambda$.

The marking and the maps ι, D, Λ, T are supplied data.
A bare complex does not reconstruct them. For $M = 0$, both divisors are 1.