

Relation presentation and zeroth homology

$$P_1 \xrightarrow{d_1} P_0 \xrightarrow{\varepsilon} F_{\text{rel}} \longrightarrow 0, \quad H_0 = P_0 / \text{im } d_1 \xrightarrow[\sim]{\bar{\varepsilon}} F_{\text{rel}}$$

Temporary equation algebra

$\mathcal{A}_{\text{eq}}$ acts by chain maps.

$$\bar{\rho} : \mathcal{A}_{\text{eq}} \rightarrow \text{End}(H_0)$$

$$C(a) = \bar{\varepsilon} \bar{\rho}(a) \bar{\varepsilon}^{-1}$$

$$\mathcal{C}_{\text{syz}} = \text{im } C$$

$$\simeq \mathcal{A}_{\text{eq}} / \text{Ann}(F_{\text{rel}})$$

Supplied marked chain maps

$$\mathfrak{G}_{\text{mark}} = \{G_1, \dots, G_t\}$$

$$dG_j = G_j d \implies \bar{G}_j \in \text{End}(H_0)$$

$$\mathcal{A}_{\text{eq}}^{\text{eff}} = \mathbb{k}\langle \bar{G} : G \in \mathfrak{G}_{\text{mark}} \rangle$$

The smallest unital subalgebra containing the descended marks.

Identify the represented actions, then generate the source

$$\mathcal{C}_{\text{syz}} \simeq \mathcal{A}_{\text{eq}}^{\text{eff}}, \quad N_{\text{eq}} = \mathcal{A}_{\text{eq}}^{\text{eff}} \text{ im } R_0$$

Transport uses $\bar{\varepsilon}$; this is not literal matrix equality in unrelated bases.

Homotopy-compatible comparison of corresponding marks

$$\Phi G - G' \Phi = d' H_G + H_G d \implies \bar{\Phi} \bar{G} = \bar{G}' \bar{\Phi}$$

If $\bar{\Phi}$ is invertible, the effective algebras are conjugate.

No extension of the source action outside the relation image is needed.