

1. Operator closure on H_0 , with $h = \dim H_0$

$$V_0 = \mathbb{k}I, \quad V_{j+1} = V_j + \sum_{G \in \mathfrak{G}_{\text{mark}}} \bar{G}V_j$$

$$V_{j+1} = V_j \implies V_j = \mathcal{A}_{\text{eq}}^{\text{eff}}$$

At most h^2 strict increases; at most $h^2 - 1$ when $h > 0$.



2. Source closure and normalization output

$$N_0 = \text{im } R_0, \quad N_{j+1} = N_j + \sum_{G \in \mathfrak{G}_{\text{mark}}} \bar{G}N_j$$

$$N_{\text{eq}} = \mathcal{A}_{\text{eq}}^{\text{eff}} \text{im } R_0, \quad q_{\text{eq}} : F \rightarrow F/\iota(N_{\text{eq}}), \quad \Gamma_{\text{eq}} = q_{\text{eq}}D\Lambda$$



3. Stable assembly under landing descent

$$T \ker \Lambda \subseteq \ker \Lambda, \quad W = \bigcap_{j=0}^{m-1} \ker(\Gamma_{\text{eq}} T^j), \quad m = \dim M > 0$$

$$\ker \Lambda \subseteq W, \quad \tilde{T} : W/\ker \Lambda \rightarrow W/\ker \Lambda$$

$$\chi_{\text{forced}} = \chi_{\tilde{T}}, \quad \mu_{\text{forced}} = \mu_{\tilde{T}}$$

$$\text{If } T \ker \Gamma_{\text{eq}} \subseteq \ker \Gamma_{\text{eq}}, \text{ then } W = \ker \Gamma_{\text{eq}}.$$

No filtration is needed for termination. If the whole effective algebra preserves a length- L filtration, its shift ideal J satisfies $J^L = 0$.

Both output polynomials are 1 when $M = 0$.