

Exact rational data and the unchanged full orbit

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 2 & 0 \\ 1 & 0 & 3 \end{pmatrix}, \quad H = \begin{pmatrix} 0 & 0 & 1 \end{pmatrix}, \quad A_\tau = A + \tau H$$

Only the third column varies.

Noninvariant kernel and its invariant core

$$U = \langle e_2 \rangle \subset K = \langle e_1, e_2 \rangle \subset R = \mathbb{Q}^3$$
$$Ae_1 = e_3 \notin K, \quad U = \bigcap_{j=0}^2 \ker(HA^j), \quad A|_U = 2I_U$$

First cyclic witness

$$\tau_0 = (0, 0, -3)^\top$$
$$\chi_{A+\tau_0 H} = (S - 2)S^2$$
$$\mu_{A+\tau_0 H} = (S - 2)S^2$$

Second cyclic witness

$$\tau_1 = (-1, 0, -3)^\top$$
$$\chi_{A+\tau_1 H} = (S - 2)(S^2 + 1)$$
$$\mu_{A+\tau_1 H} = (S - 2)(S^2 + 1)$$

Exact stable-core floor

$$\gcd_\tau \chi_{A_\tau} = S - 2 = \chi_{A|_U}, \quad \gcd_\tau \mu_{A_\tau} = S - 2 = \mu_{A|_U}$$

The two witnesses realize both gcds inside the original tilt family.