



# Recommended Vaccinations for Children

FEB. 11, 2026 •



**PURPOSE**


Guide for parents and caregivers to ensure their child is up-to-date on recommended childhood vaccines.

## Childhood Immunization Schedule by Recommendation Group

### Immunizations Recommended for All Children

| Birth                                                   | 1 mo | 2 mos | 4 mos    | 6 mos    | 7 mos    | 8 mos | 12 m |
|---------------------------------------------------------|------|-------|----------|----------|----------|-------|------|
| Diphtheria, tetanus, acellular pertussis (DTaP < 7 yrs) |      |       | 1st dose | 2nd dose | 3rd dose |       |      |
| Tetanus, diphtheria, acellular pertussis (Tdap ≥ 7 yrs) |      |       |          |          |          |       |      |
| Haemophilus influenzae type b (Hib)                     |      |       | 1st dose | 2nd dose | 3rd dose |       |      |
| Pneumococcal conjugate (PCV15, PCV20)                   |      |       | 1st dose | 2nd dose | 3rd dose |       |      |
| Inactivated poliovirus (IPV < 18 yrs)                   |      |       | 1st dose | 2nd dose | 3rd dose |       |      |
| Measles, mumps, rubella (MMR)                           |      |       |          |          |          |       |      |
| Varicella (VAR)                                         |      |       |          |          |          |       |      |
| Human papillomavirus (HPV)                              |      |       |          |          |          |       |      |

Childhood immunization schedule



 Some children should get this dose of vaccine at this age depending on their dose series

### Immunizations recommended for Certain High-Risk Groups or Populations

| Birth                                              | 1 mo   | 2 mos    | 4 mos | 6 mos | 7 mos    | 8 mos | 12 mos   | 15 mos | 18 mos | 19 m |
|----------------------------------------------------|--------|----------|-------|-------|----------|-------|----------|--------|--------|------|
| Respiratory syncytial virus (RSV-mAb) <sup>1</sup> | 1 dose |          |       |       |          |       |          |        |        |      |
| Respiratory syncytial virus (RSV-mAb) <sup>2</sup> | 1 dose |          |       |       |          |       | 2nd dose |        |        |      |
| Hepatitis B (HepB) <sup>3</sup>                    | dose   | 2nd dose |       |       | 3rd dose |       |          |        |        |      |

| Birth                           | 1 mo | 2 mos | 4 mos    | 6 mos | 7 mos | 8 mos | 12 mos | 15 mos | 18 mos | 19 m |
|---------------------------------|------|-------|----------|-------|-------|-------|--------|--------|--------|------|
| Dengue <sup>4</sup>             |      |       |          |       |       |       |        |        |        |      |
| Meningococcal ACWY <sup>5</sup> |      |       | 2, 3, or |       |       |       |        |        |        |      |
| Meningococcal B <sup>6</sup>    |      |       |          |       |       |       |        |        |        |      |
| Hepatitis A (HepA) <sup>7</sup> |      |       |          |       |       |       |        |        |        |      |

<sup>1</sup> All children whose mother did not have the vaccine should get one dose.

<sup>2</sup> High-risk children, such as those with chronic lung disease, should receive a second dose at ages 8 to 19 months.

<sup>3</sup> Vaccination recommended for infants born to women who tested positive for the hepatitis B virus or whose status is unknown.

<sup>4</sup> Recommended ONLY if living in areas with endemic dengue AND with a laboratory confirmation of previous dengue infection.


<sup>5</sup> For high-risk groups (e.g., those with anatomic or functional asplenia or HIV infection), those traveling to countries with hyperendemic or epidemic meningococcal disease, and first-year college students living in residential housing, vaccination is recommended.

<sup>6</sup> Recommended for high-risk groups, e.g., with anatomic or functional asplenia, and during outbreaks.

<sup>7</sup> Vaccination recommended for international travel to areas with high or intermediate hepatitis A endemicity.

## Immunizations Based on Shared Clinical Decision-Making

| Birth                                                    | 1 mo | 2 mos | 4 mos    | 6 mos    | 7 mos    | 8 mos | 12 mos |
|----------------------------------------------------------|------|-------|----------|----------|----------|-------|--------|
| Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series) |      |       | 1st dose | 2nd dose | 3rd dose |       |        |
| COVID-19 (1vCOV-mRNA, 1vCOV-aPS)                         |      |       |          |          | 2nd dose |       |        |
| Influenza (IIV3, cclIV3)                                 |      |       |          |          |          |       |        |
| OR                                                       |      |       |          |          |          |       |        |
| Influenza (LAIV3)                                        |      |       |          |          |          |       |        |
| Hepatitis A (HepA)                                       |      |       |          |          |          |       |        |
| Hepatitis B (HepB)*                                      |      |       | 1st dose | 2nd dose |          |       |        |
| Meningococcal ACWY                                       |      |       |          |          |          |       |        |
| Meningococcal B                                          |      |       |          |          |          |       |        |



 Some children should get this dose of vaccine at this age depending on their dose series

\* For parents deciding whether to vaccinate for HepB in infants born to women who tested negative for the hepatitis B virus, it is suggested that the initial dose is administered no earlier than 2 months of age.

## What diseases do these vaccines protect against?

| Vaccine-Preventable Disease                                                                                                                                                 | Disease Complications                                                      |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|
| <b>Diphtheria</b><br>Illness caused by a toxin produced by bacteria that infects the nose, throat, and sometimes skin.                                                      | Swelling of the heart muscle, heart failure, coma, paralysis, death        |
| <b>Tetanus (Lockjaw)</b><br>Bacterial infection of brain and nerves caused by spores found in soil and dust everywhere; spores enter the body through wounds or broken skin | Seizures, broken bones, difficulty breathing, death                        |
| <b>Pertussis (Whooping Cough)</b><br>Contagious bacterial infection of the lungs and airway; spread through air and direct contact                                          | Infection of the lungs (pneumonia), death; especially dangerous for babies |

| Vaccine-Preventable Disease                                                                                                                                                                                       | Disease Complications                                                                                                                                                      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Hib (<i>Haemophilus influenzae</i> type b)</b><br>Contagious bacterial infection of the lungs, brain and spinal cord, or bloodstream; spread through air and direct contact                                    | Depends on the part of the body infected, but can include brain damage, hearing loss, loss of arm or leg, death                                                            |
| <b>Pneumococcal</b><br>Bacterial infections of ears, sinuses, lungs, or bloodstream; spread through direct contact with respiratory droplets like saliva or mucus                                                 | Depends on the part of the body infected, but can include infection of the lungs (pneumonia), blood poisoning, infection of the lining of the brain and spinal cord, death |
| <b>Polio</b><br>Contagious viral infection of nerves and brain; spread through the mouth from stool on contaminated hands, food or liquid, and by air and direct contact                                          | Paralysis, death                                                                                                                                                           |
| <b>Measles (Rubeola)</b><br>Contagious viral infection that causes high fever, cough, red eyes, runny nose, and rash; spread through air and direct contact                                                       | Brain swelling, infection of the lungs (pneumonia), death                                                                                                                  |
| <b>Mumps</b><br>Contagious viral infection that causes fever, tiredness, swollen cheeks, and tender swollen jaw; spread through air and direct contact                                                            | Brain swelling, painful and swollen testicles or ovaries, deafness, death                                                                                                  |
| <b>Rubella (German Measles)</b><br>Contagious viral infection that causes low-grade fever, sore throat, and rash; spread through air and direct contact                                                           | Very dangerous in pregnant women; can cause miscarriage or stillbirth, premature delivery, severe birth defects                                                            |
| <b>Chickenpox (Varicella)</b><br>Contagious viral infection that causes fever, headache, and an itchy, blistering rash; spread through air and direct contact                                                     | Infected sores, brain swelling, infection of the lungs (pneumonia), death                                                                                                  |
| <b>HPV (Human papillomavirus)</b><br>Contagious viral infection spread by close skin-to-skin touching, including during sex                                                                                       | Genital warts and many types of cancers later in life, including cancers of the cervix, vagina, penis, anus, and throat                                                    |
| <b>RSV (Respiratory syncytial virus)</b><br>Contagious viral infection of the nose, throat, and sometimes lungs; spread through air and direct contact                                                            | Infection of the lungs (pneumonia) and small airways of the lungs; especially dangerous for infants and young children                                                     |
| <b>Hepatitis B</b><br>Contagious viral infection of the liver; spread through contact with infected body fluids such as blood or semen                                                                            | Chronic liver infection, liver failure, liver cancer, death                                                                                                                |
| <b>Dengue</b><br>Viral infection spread by bite from infected mosquito; causes, fever, headache, pain behind the eyes, rash, joint pain, body ache, nausea, loss of appetite, feeling tired, abdominal pain       | Severe bleeding, seizures, shock, damage to the liver, heart, and lungs, death                                                                                             |
| <b>Mpox</b><br>Contagious viral infection spread through close, often skin-to-skin contact, including sex; causes a painful rash, fever, headache, tiredness, cough, runny nose, sore throat, swollen lymph nodes | Infected sores, brain swelling, infection of the lungs (pneumonia), eye infection, blindness, death                                                                        |
| <b>Meningococcal</b><br>Contagious bacterial infection of the lining of the brain and spinal cord or the bloodstream; spread through air and direct contact                                                       | Loss of arm or leg, deafness, seizures, death                                                                                                                              |
| <b>Rotavirus</b><br>Contagious viral infection of the gut; spread through the mouth from hands and food contaminated with stool                                                                                   | Severe diarrhea, dehydration, death                                                                                                                                        |
| <b>COVID-19</b><br>Contagious viral infection of the nose, throat, or lungs; may feel like a cold or flu. Spread through air and direct contact                                                                   | Infection of the lungs (pneumonia); blood clots; liver, heart or kidney damage; long COVID; death                                                                          |
| <b>Influenza (Flu)</b><br>Contagious viral infection of the nose, throat, and sometimes lungs; spread through air and direct contact                                                                              | Infection of the lungs (pneumonia), sinus and ear infections, worsening of underlying heart or lung conditions, death                                                      |
| <b>Hepatitis A</b><br>Contagious viral infection of the liver; spread by contaminated food or drink or close contact with an infected person                                                                      | Liver failure, death                                                                                                                                                       |

 [Get email updates](#)

More Schedule Resources

- [Learn about vaccine-preventable diseases](#)
- [Syndicate the schedules on your website](#)
- [Compliant version of the schedule](#)

For Other Groups

- [Aged 19 Years and Older](#)
- [Pregnant Women](#)

SOURCES

CONTENT SOURCE:

National Center for Immunization and Respiratory Diseases; Immunization Services Division