

JOINT DISTRIBUTION IN RESIDUE CLASSES OF FAMILIES OF MULTIPLICATIVE FUNCTIONS I

AKASH SINGHA ROY

ABSTRACT. We study the joint distribution of families of multiplicative functions in residue classes, allowing the moduli to vary within a wide range and assuming some natural control on the average behavior of the functions at (some fixed power of) the primes. As applications, we obtain essentially best possible analogues of the Siegel–Walfisz theorem for families of multiplicative functions that can be controlled by polynomials at some powers of primes. (This class includes several interesting arithmetic functions such as Euler’s totient $\varphi(n)$, the sum-of-divisors $\sigma(n)$, and more generally, the sum-of-divisor-powers $\sigma_r(n) := \sum_{d|n} d^r$, and so on.) Our results (optimally) extend works of Narkiewicz, Rayner, Śliwa, Dobrowolski, Fomenko and others. One of the main ideas behind our arguments is the detection of a certain “mixing” (or “quantitative ergodicity”) phenomenon via methods from sieve theory and the anatomy of integers. Additionally, we combine ideas and machinery from classical analytic number theory, character sums, linear algebra over rings, as well as arithmetic and algebraic geometry.

1. INTRODUCTION

We say that a function $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ is **equidistributed mod $q \in \mathbb{Z}^+$** if $\#\{n \leq x : f(n) \equiv a \pmod{q}\} \sim x/q$ as $x \rightarrow \infty$. This notion has been studied for *additive* functions: See results of Delange [6, 7], which have been partially extended in [26, 27, 1, 32]. However, for *multiplicative* functions, this is not the correct notion to work with: For instance, using classical results (such as those implicit in [16]), it can be shown that Euler’s totient function $\varphi(n)$ is almost always divisible by any fixed integer q , and hence is **not** equidistributed modulo any fixed $q > 1$. Motivated by this, Narkiewicz defines a function $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ to be **weakly equidistributed (WUD) mod q** if there are infinitely many n for which $(f(n), q) = 1$, and if

$$\#\{n \leq x : f(n) \equiv a \pmod{q}\} \sim \frac{1}{\varphi(q)} \#\{n \leq x : (f(n), q) = 1\} \quad \text{as } x \rightarrow \infty,$$

for every coprime residue $a \pmod{q}$.¹ Hence, our sample space is $\{n : (f(n), q) = 1\}$, and every coprime residue mod q gets its fair share of the sample space. For example, if f is WUD mod 6, then the residues 1 and 5 mod 6 each asymptotically receive 50% of the sample space $\{n : (f(n), 6) = 1\}$. Likewise, given a family of functions $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$, we say that $(f_1, \dots, f_K)$ are **WUD mod q** if there are infinitely many n for which $(f_1(n) \dots f_K(n), q) = 1$, and if for all coprime residues $a_1, \dots, a_K \pmod{q}$, we have

$$(1.1) \quad \#\{n \leq x : (\forall i) f_i(n) \equiv a_i \pmod{q}\} \sim \frac{1}{\varphi(q)^K} \#\{n \leq x : (f_1(n) \dots f_K(n), q) = 1\} \quad \text{as } x \rightarrow \infty.$$

The notion of weak equidistribution has been studied by several authors. Narkiewicz gave a criterion [19, Theorem 1] deciding weak equidistribution of any “polynomially-defined” multiplicative function; here, $f : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ is **polynomially-defined** if there exist $V \in \mathbb{Z}^+$ and polynomials $\{F_v\}_{v=1}^V \subset \mathbb{Z}[T]$ satisfying $f(p^v) = F_v(p)$ for all primes p and all $v \in [V]$. (Hence, F_v controls the behavior of f at the v -th powers of primes.) As a consequence, he showed that $\varphi(n)$ is WUD mod q iff $(q, 6) = 1$. (Related are works of Dence–Pomerance [8] and Banks–Shparlinski [2, Theorem 3.1].) Śliwa [34] used Narkiewicz’s criterion to show that the sum-of-divisors $\sigma(n) = \sum_{d|n} d$ is WUD mod q iff $6 \nmid q$. This was extended to the functions

2020 *Mathematics Subject Classification.* Primary 11A25; Secondary 11N36, 11N37, 11N64, 11N69.

Key words and phrases. multiplicative function, uniform distribution, equidistribution, weak uniform distribution, joint distribution, distribution in residue classes.

¹Here, (a, b) denotes the gcd of a and b , unless stated otherwise.

$\sigma_r(n) = \sum_{d|n} d^r$ (which also appear as Fourier coefficients of Eisenstein series) by Narkiewicz, Rayner, Śliwa, Dobrowolski and Fomenko [34, 11, 24, 22, 23, 28, 29]. Finally, the most general results were also obtained by Narkiewicz in [21, 20], where he gave a complete characterization deciding when an arbitrary *family* of polynomially-defined multiplicative functions is weakly equidistributed to a fixed modulus.

In all these works, q is assumed to be *fixed*. It is then natural to ask what happens when q is allowed to *grow* with x , in analogy with the Siegel–Walfisz theorem from prime number theory. Analogues of Siegel–Walfisz are always sought both due to their independent interest and applications to other problems, and such analogues have been studied in other contexts, such as for smooth numbers and mean values of multiplicative functions. In our context, the explicit question is: Can we find analogues of the Siegel–Walfisz theorem for the joint distribution of families of multiplicative functions? To formalize this, given $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ and a set $\mathcal{Q} \subset \mathbb{Z}^+$, we say that $(f_1, \dots, f_K)$ are WUD (uniformly) mod $q \in \mathcal{Q}$, if:

- For every $q \in \mathcal{Q}$, we have $(\prod_{i=1}^K f_i(n), q) = 1$ for infinitely many n , and
- (1.1) holds as $x \rightarrow \infty$, uniformly in $q \in \mathcal{Q}$ and in coprime residues $a_1, \dots, a_K \bmod q$.

The only results in this direction seem to appear in [17, 25, 27], which make partial progress towards extending *highly special cases* of Narkiewicz’s criteria [19, 21] to varying q . In [17], Lebowitz-Lockard, Pollack and the author show that $\varphi(n)$ is WUD uniformly mod *prime* $q \in [5, (\log x)^{K_0}]$ for any fixed K_0 . In [25], Pollack and the author consider multiplicative $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ for which there exist nonconstant $F_1, \dots, F_K \in \mathbb{Z}[T]$ with the product $F_1 \dots F_K$ separable², such that $f_i(p) = F_i(p)$ for all i and all primes p : We show that such $(f_1, \dots, f_K)$ are WUD uniformly mod $q \leq (\log x)^{1/K-\epsilon}$ that are “almost prime” in a certain sense. Finally, [27] partially improves this only for $K = 1$, optimizing the range of q and replacing “almost prime” by the condition that q is supported on sufficiently large primes.

The above restrictions are unnecessary in the fixed- q results, but they cannot be overcome with the methods available in [17, 25, 27]. For instance, most arguments in [27] do not generalize to families (i.e. to $K > 1$), and the separability and the (roughness) condition on q are both strictly necessary for the arguments. Moreover, Narkiewicz’s criteria [19, 21] allow a certain flexibility with prime powers and the possibility of sparse input sets (see § 7.2), both of which are crucial in most of the other fixed- q results cited above; however [17, 25, 27] cannot allow any such flexibility. Hence, these works cannot give satisfactory uniform extensions of most of the other works of Narkiewicz, Rayner, Śliwa and others.

In this manuscript, we remove *all* these limitations and give essentially optimal uniform extensions of Narkiewicz’s most general criteria [21, 20] (for a single growing modulus), hence also completely extending all the cited works. Our q will vary either within optimal ranges *or* up to any fixed power of $\log x$, hence we also obtain essentially best possible qualitative analogues of the *Siegel–Walfisz theorem* for a general family of polynomially-defined multiplicative functions. To do this, we not only need to refine the methods in the previous works (which were restricted to the anatomy of integers), but also use several ideas not considered before, such as from linear algebra over rings, and from algebraic and arithmetic geometry. As mentioned in § 1.2, some of these ideas have potential applications to other related problems as well.

1.1. The Main Results. Since the general version of Narkiewicz’s criterion in [21] requires some technical set-up, we first give uniform analogues of a special case to illustrate the most important ideas. In section 7, we state our most general results and the additional ingredients they require.

Consider multiplicative functions $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ for which there exist polynomials $F_1, \dots, F_K \in \mathbb{Z}[T]$ satisfying $f_i(p) = F_i(p)$ for all primes p and all $i \in [K]$. For any $q \in \mathbb{Z}^+$, we define

$$\alpha(q) := \varphi(q)^{-1} \# \{u \in U_q : F_1(u) \dots F_K(u) \in U_q\}, \text{ where } U_q := (\mathbb{Z}/q\mathbb{Z})^\times.$$

Let $\mathcal{Q}(f_1, \dots, f_K)$ denote the set of all $q \in \mathbb{Z}^+$ with $\alpha(q) \neq 0$ that satisfy the following property:

²i.e. having no repeated roots in $\mathbb{C}$

For all Dirichlet characters $(\chi_1, \dots, \chi_K) \neq (\chi_0, \dots, \chi_0) \bmod q$ satisfying $\prod_{i=1}^K \chi_i(F_i(u)) = 1$ on its “unit support” $\{u \in U_q : \prod_{i=1}^K F_i(u) \in U_q\}$, we have $\prod_{i=1}^K \chi_i(f_i(2^j)) = -1$ for all $j \geq 1$.

Here χ_0 is the trivial character mod q . Narkiewicz’s criterion in this special setting is as follows.

Theorem 1.1. [21] *Assume $\alpha(q) \neq 0$. Then $(f_1, \dots, f_K)$ are WUD mod q iff $q \in \mathcal{Q}(f_1, \dots, f_K)$.*

Now assume that $F_1, \dots, F_K$ are multiplicatively independent, i.e. $\prod_{i=1}^K F_i^{c_i}$ is nonconstant in $\mathbb{Q}(T)$ for all integers $(c_1, \dots, c_K) \neq (0, \dots, 0)$.³ Factor each F_i as $r_i \prod_{j=1}^M G_j^{\mu_{i,j}}$ with $r_i \in \mathbb{Z}$, $\mu_{i,j} \in \mathbb{Z}_{\geq 0}$, and with $\{G_j\}_{j=1}^M \subset \mathbb{Z}[T]$ being pairwise coprime primitive⁴ irreducibles, such that each G_j appears with a positive exponent $\mu_{i,j}$ in some F_i . Let E_0 be the $M \times K$ matrix $\left((\mu_{i,j})_{\substack{1 \leq i \leq K \\ 1 \leq j \leq M}} \right)^\top$, and $\beta \in \mathbb{Z}^+$ be the largest invariant factor of E_0 . Since $\{F_i\}_i$ are multiplicatively independent, the columns of E_0 are $\mathbb{Q}$ -linearly independent (i.e. $\text{rank}(E_0) = K \leq M$), so β is the last diagonal entry in the Smith Normal Form of E_0 .

Fix any $B_0 > 0$. We say that q satisfies Hypothesis $IFH(F_1, \dots, F_K; B_0)$ if all its prime divisors ℓ exceeding B_0 satisfy $\gcd(\ell - 1, \beta) = 1$. Note that if $\prod_{i=1}^K F_i$ is separable (as was assumed in previous works [25, 27]) then $\beta = 1$, so that any q satisfies $IFH(F_1, \dots, F_K; B_0)$ for any $B_0 > 0$.

Theorem 1.2. *Fix any $K, B_0 > 0$ and $\epsilon \in (0, 1)$, and assume that $F_1, \dots, F_K$ are multiplicatively independent. Then $(f_1, \dots, f_K)$ are WUD mod $q \leq (\log x)^{K_0}$ lying in $\mathcal{Q}(f_1, \dots, f_K)$, such that q satisfies $IFH(F_1, \dots, F_K; B_0)$ along with at least one of the following two conditions:*

- (i) $q \leq (\log x)^{(1-\epsilon)\alpha(q)} (K-1/D_{\min})^{-1}$, where $D_{\min} := \min\{\deg F_i : 1 \leq i \leq K\}$, **or**
- (ii) q is squarefree and $q^{K-1} D_{\min}^{\omega(q)} \leq (\log x)^{(1-\epsilon)\alpha(q)}$.

For $K = D_{\min} = 1$ (single multiplicative function controlled by a linear polynomial at the primes), (i) and (ii) are tautological, so we have uniformity in $q \leq (\log x)^{K_0}$. Some applications of Theorem 1.2: $\varphi(n)$ is WUD mod $q \leq (\log x)^{K_0}$ coprime to 6 (extending Narkiewicz [19, Cor. 2]); $\sigma(n)$ is WUD mod *odd* $q \leq (\log x)^{K_0}$ (partially extending Śliwa [34]); (φ, σ) are WUD mod $q \leq (\log x)^{(1-\epsilon)\alpha_2(q)}$ coprime to 6, where $\alpha_2(q) = \prod_{\ell|q} (\ell - 3)/(\ell - 1)$ (extending Narkiewicz [20, Theorem 1] for a single modulus). A simple consequence of (ii) and (4.1) below is that if $F_1, \dots, F_K$ are multiplicatively independent, then $(f_1, \dots, f_K)$ are WUD mod prime $q \leq (\log x)^{(1/(K-1)-\epsilon)\alpha(q)}$ satisfying $q \in \mathcal{Q}(f_1, \dots, f_K)$ and $\gcd(q - 1, \beta) = 1$.

In section 6, we show that for any K and $D_{\min}$ (except $K = D_{\min} = 1$), both ranges in (i) and (ii) are essentially optimal, in that “ $1 - \epsilon$ ” cannot be replaced by “ $1 + \epsilon$ ” in either. In fact, if $K > 1$ and $D_{\min} = 1$, then this optimality holds in a much stronger sense: The ranges in (i) and (ii) are optimal no matter how many of the F_i we assume to be linear and even if we take all the F_i to be essentially arbitrary linear polynomials. In particular, the range $q \leq (\log x)^{(1-\epsilon)\alpha_2(q)}$ above is optimal even for the *particular example* of (φ, σ) . Moreover, the multiplicative independence hypothesis and hypothesis $IFH(F_1, \dots, F_K; B_0)$ are also both necessary in Theorem 1.2 and in Theorem 1.3 below: We shall establish this in the sequel [33].

As our constructions in section 6 show, obstructions to uniformity come from prime inputs. We can modify those constructions to produce more obstructions coming from inputs n that have *too few* large prime factors. Hence to restore uniformity in the full Siegel-Walfisz range, we need to restrict our n to those with *sufficiently many* large prime factors. To this end, let $\mathbf{P}_R(\mathbf{n})$ denote the R -th largest prime factor of n counted with multiplicity (with the convention $\mathbf{P}_R(\mathbf{n}) := 1$ if $\Omega(\mathbf{n}) < R$).

³This hypothesis is easily satisfied in almost all applications: For example, it is satisfied if $\prod_{i=1}^K F_i$ is separable, or (more generally) if each F_i has an irreducible factor that is not present in the other F_j (for $j \neq i$).

⁴A polynomial in $\mathbb{Z}[T]$ is *primitive* when the greatest common divisor of its coefficients is 1.

Theorem 1.3. Fix $K_0, B_0 > 0$ and assume that $F_1, \dots, F_K$ are multiplicatively independent. Then

$$(1.2) \quad \sum_{\substack{n \leq x: P_R(n) > q \\ (\forall i) f_i(n) \equiv a_i \pmod{q}}} 1 \sim \frac{1}{\varphi(q)^K} \sum_{\substack{n \leq x: P_R(n) > q \\ (\prod_i f_i(n), q) = 1}} 1 \sim \frac{1}{\varphi(q)^K} \sum_{\substack{n \leq x \\ (\prod_i f_i(n), q) = 1}} 1 \quad \text{as } x \rightarrow \infty,$$

uniformly in $q \leq (\log x)^{K_0}$ lying in $\mathcal{Q}(f_1, \dots, f_K)$ and satisfying $IFH(F_1, \dots, F_K; B_0)$, as well as uniformly in coprime residues $a_1, \dots, a_K$ modulo q . Here we can take

$$(1.3) \quad R = \begin{cases} KD + 1, & \text{in general, where } D := \sum_{i=1}^K \deg F_i, \\ 2K + 1, & \text{if } q \text{ is squarefree,} \\ 2, & \text{if } q \text{ squarefree, } K = 1, F_1 \text{ is not squarefull (i.e. } F_1 \text{ has a simple root in } \mathbb{C}). \end{cases}$$

For instance, (φ, σ) are WUD uniformly modulo $q \leq (\log x)^{K_0}$ coprime to 6 if we restrict to inputs n with $P_5(n) > q$. (Without this restriction, we only have the aforementioned optimal range $q \leq (\log x)^{(1-\epsilon)\alpha_2(q)}$.) Another application: By Narkiewicz [20, 21], $(\varphi, \sigma, \sigma_2)$ are WUD modulo any fixed q having $P^-(q) > 23$. By Theorem 1.2, this weak equidistribution holds uniformly modulo any such $q \leq (\log x)^{(1/2-\epsilon)c_q}$ where $c_q := \prod_{\substack{\ell|q \text{ prime} \\ \ell \equiv -1 \pmod{4}}} \frac{\ell-3}{\ell-1} \cdot \prod_{\substack{\ell|q \text{ prime} \\ \ell \equiv 1 \pmod{4}}} \frac{\ell-5}{\ell-1}$. Uniformity is restored modulo all $q \leq (\log x)^{K_0}$ with $P^-(q) > 23$, if we restrict to n with $P_{13}(n) > q$. This can be weakened to $P_7(n) > q$, if q is also assumed squarefree.

The smaller the value of R in (1.2), the larger our set wherein equidistribution occurs, and the better our result. In section 6, we show that the third value of R in (1.3) is optimal, while for any $K \in \mathbb{Z}^+$, the second value is at most “one step away” from being optimal (in that it cannot be lowered to $2K - 1$). Note that even the $K = 1$ special case of Theorem 1.3 improves over [27, Theorem 1.4]. It is not entirely clear whether the first value of R in (1.3) can be improved in the relatively special setting of Theorem 1.3 above; however, we remark that in the more general setting of Theorem 7.6 below, the corresponding value of R in (7.11) does turn to be optimal (as discussed after Theorem 7.6).

One important theme in our arguments is a “mixing” phenomenon originally observed in [27]: Intuitively, given $G \in \mathbb{Z}[T]$ and q supported on large primes, if we choose uniformly at random, units $u_1, u_2, \dots \pmod{q}$ such that each $G(u_j)$ is a unit mod q , then in the sequence of partial products $\{G(u_1) \dots G(u_j)\}_{j=1}^\infty$, every unit mod q appears equally often. Here we detect and generalize this “mixing” via more careful arguments from sieve theory, the anatomy of integers and character sums (extensions of Weil bounds).

Moreover, we need several ideas and machinery not considered in previous works, such as from linear algebra over rings, and algebraic/arithmetical geometry (via regular sequences and counting rational points on varieties over finite fields). We also need a theorem of Halász (which requires us to carefully estimate certain “pretentious distances”⁵), and we need to *modify* the Landau–Selberg–Delange “LSD” method. (Known forms of LSD or other mean value results are *not* sufficient to get the desired uniformity in q .)

1.2. Beyond the polynomially-defined regime. The aforementioned “mixing”, extended suitably, can be used to study the joint weak equidistribution of multiplicative functions $h_1, \dots, h_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ assuming only a natural control on the *average* behavior of the h_i at the primes. Owing to this generality, we expect the following results to have applications beyond the “polynomially-defined” regime, such as for Ramanujan’s τ -function and Fourier coefficients of modular forms.

We assume that for some (large) *threshold parameter* y and for all Dirichlet characters $(\chi_1, \dots, \chi_K) \pmod{q}$, we have an estimate of the form $\sum_{y < p \leq Y} \prod_i \chi_i(h_i(p)) \approx \alpha_{\widehat{\chi}} \sum_{y < p \leq Y} \prod_i \chi_0(h_i(p))$ for all $Y \geq y$, where $\alpha_{\widehat{\chi}} \in \mathbb{C}$ depends only on $\widehat{\chi} := (\chi_1, \dots, \chi_K)$. Assuming this with an essentially arbitrary error term, we bound the *extent* to which $(h_1, \dots, h_K)$ fails to be weakly equidistributed mod q . Our first result shows

⁵in the terminology of Granville and Soundararajan [12]

that for any $J \in \mathbb{Z}^+$ for which the J -th moment $\sum_{\chi_1, \dots, \chi_K} |\alpha_{(\chi_1, \dots, \chi_K)}|^J$ is small enough, $(h_1, \dots, h_K)$ are WUD mod q , upon restricting to inputs n with J many large prime factors (exceeding the *threshold* y).

In what follows, we set $\hat{\chi} := (\chi_1, \dots, \chi_K)$ and let $\mathfrak{f}(\hat{\chi})$ denote the lcm of the conductors of χ_i . We write $\hat{h} := (h_1, \dots, h_K)$, $h := h_1 \cdots h_K$, and $\hat{\chi}(\hat{h}(n)) := \chi_1(h_1(n)) \cdots \chi_K(h_K(n))$.

Theorem 1.4. *Let $h_1, \dots, h_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ be multiplicative functions. Consider any $q \in \mathbb{Z}^+$, $M \geq 1$, $y \geq q$, and a decreasing function $\mathcal{E} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, such that for all $\hat{\chi} \bmod q$ and all $Y \geq y$,*

$$(1.4) \quad \sum_{y < p \leq Y} \hat{\chi}(\hat{h}(p)) = \alpha_{\hat{\chi}} \sum_{y < p \leq Y} \chi_0(h(p)) + O(MY\mathcal{E}(y)),$$

with $\alpha_{\hat{\chi}} \in \mathbb{C}$ in the unit disk. Then for all d dividing q , all $a_i \in U_q$, all $x \geq y$ and all $J \in \mathbb{Z}^+$, we have

$$(1.5) \quad \sum_{\substack{n \leq x: P_J(n) > y \\ (\forall i) h_i(n) \equiv a_i \pmod{q}}} 1 - \left(\frac{\varphi(d)}{\varphi(q)} \right)^K \sum_{\substack{n \leq x: P_J(n) > y, (h(n), q) = 1 \\ (\forall i) h_i(n) \equiv a_i \pmod{d}}} 1 \\ \ll \frac{1}{\varphi(q)^K} \sum_{\substack{\hat{\chi} \bmod q \\ \mathfrak{f}(\hat{\chi}) \nmid d}} |\alpha_{\hat{\chi}}|^J \sum_{\substack{n \leq x \\ (h(n), q) = 1}} 1 + \frac{Jx}{y} + Mx\mathcal{E}(y) \log x.$$

The implied constant here depends only on the implied constant in (1.4).

Note that (1.4) is in general weaker than a hypothesis like $\sum_{y < p \leq Y} \hat{\chi}(\hat{h}(p)) = \beta_{\hat{\chi}} \sum_{y < p \leq Y} 1 + O(MY\mathcal{E}(y))$. Further, (1.5) just needs (1.4) for all $\hat{\chi}$ with $\mathfrak{f}(\hat{\chi}) \nmid d$. Taking $J = d = 1$ in (1.5) directly bounds the failure of weak equidistribution; however, to make the right side of (1.5) negligible in applications, we often need to select a suitable parameter $J \rightarrow \infty$. We thus see what happens for the n with $P_J(n) \leq y$.

Our estimates will be in terms of a parameter that controls the (suitably normalized) count of large prime solutions to the congruences $h_i(p) \equiv a_i \pmod{q}$. Given any $h_i : \mathbb{Z}^+ \rightarrow \mathbb{Z}$, any $q, r \in \mathbb{Z}^+$ and any $z > 0$, let $\xi_r(q; z) := \xi_r(q; z; \{h_i\}_{i=1}^K) > 0$ be *any* parameter satisfying the bound

$$(K_r) \quad \#\{P_1 \cdots P_r \leq x : P_i \text{ primes}, P_1 > z, q < P_r < \cdots < P_1, (\forall i) h_i(P_1) \cdots h_i(P_r) \equiv a_i \pmod{q}\} \\ \leq \xi_r(q; z) \cdot x(\log \log x)^r / \log(z/q)$$

for all $x \geq z$ and all $a_1, \dots, a_K \in U_q$. Thus, $\xi_r(q; z)$ controls the number of squarefree integers $A \leq x$ with exactly r prime factors (all suitably large), that satisfy $h_i(A) \equiv a_i \pmod{q}$ for all i . In particular, $\xi_1(q; z)$ controls the number of primes $P \in (z, x)$ satisfying $h_i(P) \equiv a_i \pmod{q}$ for all i . Note that this is genuinely a *definition*, not a *hypothesis*, for the choice $\xi_r(q; z) = 1$ always satisfies (K_r) by Chebyshev's estimates, for all possible $\{h_i\}_i, q, r$ and z . Let $\Psi(x, z)$ be the number of z -smooth numbers up to x .

Theorem 1.5. *Fix $K \in \mathbb{Z}^+$. Uniformly in **all** multiplicative functions $h_1, \dots, h_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$, all $q, J \in \mathbb{Z}^+$, all $a_i \in U_q$, and in all real numbers $x \geq z \geq y \geq q$, we have*

$$(1.6) \quad \sum_{\substack{n \leq x: P_J(n) \leq y \\ (\forall i) h_i(n) \equiv a_i \pmod{q}}} 1 \ll \xi_1(q; z) \frac{x(2 \log \log x)^{J+1}}{\log(z/q)} \exp \left(\sum_{p \leq y} \frac{\mathbb{1}_{(h(p), q) = 1}}{p} \right) + \Psi(x, z) + \frac{x}{y}.$$

$$(1.7) \quad \sum_{\substack{n \leq x: P_J(n) \leq y \\ (h(n), q) = 1}} 1 \ll \frac{x(2 \log \log x)^J}{\log z} \exp \left(\sum_{p \leq y} \frac{\mathbb{1}_{(h(p), q) = 1}}{p} \right) + \Psi(x, z) + \frac{x}{y}.$$

The implied constants depend only on K . For any $R \leq J$, if we also impose the condition $P_R(n) > q$ on the left of (1.6), then the same bound holds, only with $\xi_1(q; z)$ and $(2 \log \log x)^{J+1}$ replaced by

$$\xi_R(q; z) + R^K \sum_{\substack{1 \leq r < R \\ 1 \leq s < K}} \frac{\xi_r(q; z)}{q^{\max\{s, R-r-s\}}} \quad \text{and} \quad (2 \log \log x)^{J+R}, \quad \text{respectively.}$$

Combining Theorem 1.4 with Theorem 1.5, we get a bound the difference

$$\sum_{\substack{n \leq x: P_R(n) > q \\ (\forall i) h_i(n) \equiv a_i \pmod{q}}} 1 - \left(\frac{\varphi(d)}{\varphi(q)} \right)^K \sum_{\substack{n \leq x: (h(n), q) = 1 \\ (\forall i) h_i(n) \equiv a_i \pmod{d}}} 1$$

for all $d \mid q$, all $x, y, z > 0$, and all $J, R \in \mathbb{Z}^+$ satisfying $q \leq y \leq z \leq x$ and $J \geq R$. The flexibility in choosing d, y, z, J, R in this bound (especially the fact that z and J now appear only on the right of this bound) are very useful in applications: For instance, by taking $d = R = 1$ and a large J , we can make the right side negligible while including *all* inputs n on the left. The strength of this bound basically depends on the bounds available on $\sum_{f(\widehat{\chi}) \nmid d} |\alpha_{\widehat{\chi}}|^J$, and on $\xi_r(q; z)$ for $r \in [R]$. As discussed in Remark 2, all the “ $f(\widehat{\chi}) \nmid d$ ” conditions can be replaced by the more general “ $\widehat{\chi} \notin \mathcal{S}$ ”, with $\mathcal{S}$ an arbitrary subset of the set of K -tuples of characters mod q ; the corresponding variants then would be uniform in $\mathcal{S}$.

We also mention the following estimate, which forms a key input in the proofs of Theorems 1.2 and 1.3, and may also be of independent interest or have applications to other problems.

Proposition 1.6. *Under the hypotheses of Theorem 1.4, we have for any $\widehat{\chi} \bmod q$,*

$$\sum_{n \leq x} \widehat{\chi}(\widehat{h}(n)) \ll \frac{x}{\log x} \exp \left(\sum_{\substack{p \leq y \\ (h(p), q) = 1}} \frac{1}{p} + |\alpha_{\widehat{\chi}}| \sum_{\substack{y < p \leq x \\ (h(p), q) = 1}} \frac{1}{p} + O(M\mathcal{E}(y)(\log x)^3) \right).$$

The implied constant depends only on the implied constant in (1.4).

Finally, we remark that Narkiewicz’s weak equidistribution criteria [19, 21] are *much* more general than Theorem 1.1: They allow the possibility that the behavior of our f_i at the ν -th powers of primes is *most significant* (in a certain sense), for some fixed but arbitrary $\nu \in \mathbb{Z}^+$; this also allows the possibility of the input sets $\{n : \gcd(\prod_i f_i(n), q) = 1\}$ being sparse. However, all the results and methods from all previous work on growing moduli are unable to handle any $\nu > 1$. Adapting our methods for Theorems 1.2 and 1.3 with additional arguments, we obtain (in section 7) optimal uniform results in this greater generality, along with several new applications that could not be addressed with previous work on growing moduli. We also give extensions of the results in subsection § 1.2 that apply for certain sparse input sets.

In sections 2 to 5, we will first prove the results in subsection § 1.2, and then deduce those in § 1.1 with additional arguments. In a future paper, we shall study the joint weak equidistribution of a family of multiplicative functions $(f_1, \dots, f_K)$ to growing moduli $(q_1, \dots, q_K)$, defined appropriately.

Conventions. We do not consider the zero function as multiplicative (thus, $f(1) = 1$ for any multiplicative function f). By $P(n)$ and $P^-(n)$, we mean the largest and least prime factors of n respectively (with $P(1) := 1$, $P^-(1) := \infty$). By the z -smooth part (resp. z -rough part) of n , we mean the largest z -smooth (resp. z -rough) integer dividing n . When there is no danger of confusion, we write $(a_1, \dots, a_K)$ in place of $\gcd(a_1, \dots, a_K)$. Throughout, we use the letters p and ℓ to denote primes. Given a tuple of characters $\widehat{\psi} := (\psi_1, \dots, \psi_K) \bmod q$, we use $f(\psi_i)$ for the conductor of ψ_i , and we let $f(\widehat{\psi}) := \text{lcm}[f(\psi_1), \dots, f(\psi_K)]$.

Implied constants in $\ll$ and O -notation, and implicit constants in qualifiers like “sufficiently large”, may depend on any parameters declared as “fixed”. Other dependence will be noted explicitly (with, say, parentheses or subscripts). $\log_k$ denotes the k -th iterate of the natural logarithm.

2. THE GENERAL MIXING PHENOMENON: PROOF OF THEOREMS 1.4 AND 1.5

Proof of Theorem 1.4. We say that $n \leq x$ is “convenient” (or “conv”) if the J largest prime divisors of n exceed y and are distinct, i.e. if n can be uniquely written as $mP_J \cdots P_1$, where

$$(2.1) \quad L_m := \max\{y, P(m)\} < P_J < \cdots < P_1.$$

Now, the number of $n \leq x$ having a repeated prime factor exceeding y is at most $\sum_{p>y} \sum_{m \leq x/p^2} 1 \ll x/y$, which is absorbed in the right of (1.5). Hence, to complete the proof of Theorem 1.4, it suffices to show (1.5) with the conditions “ $P_J(n) > y$ ” replaced by the conditions that n is convenient.

Using the orthogonality of Dirichlet characters, we detect the congruences $h_i(n) \equiv a_i \pmod{q}$ to get

$$(2.2) \quad \sum_{\substack{n \leq x \text{ conv} \\ (\forall i) \ h_i(n) \equiv a_i \pmod{q}}} 1 = \frac{1}{\varphi(q)^K} \sum_{\widehat{\chi} \bmod q} \overline{\chi}_1(a_1) \cdots \overline{\chi}_K(a_K) \sum_{n \leq x \text{ conv}} \widehat{\chi}(\widehat{h}(n)).$$

For any $\widehat{\chi} \bmod q$ for which $\mathfrak{f}(\widehat{\chi}) \mid d$, there exists a unique tuple of characters $\widehat{\psi} := (\psi_1, \dots, \psi_K) \bmod d$ such that $\chi_i = \chi_0 \psi_i$ for all i . Thus the total of contribution of all such $\widehat{\chi}$ to the right of (2.2) equals

$$(2.3) \quad \frac{1}{\varphi(q)^K} \sum_{\widehat{\psi} \bmod d} \overline{\psi}_1(a_1) \cdots \overline{\psi}_K(a_K) \sum_{\substack{n \leq x \text{ conv} \\ (h(n), q) = 1}} \widehat{\psi}(\widehat{h}(n)) = \left(\frac{\varphi(d)}{\varphi(q)} \right)^K \sum_{\substack{n \leq x \text{ conv}: (h(n), q) = 1 \\ (\forall i) \ h_i(n) \equiv a_i \pmod{d}}} 1.$$

Now consider any $\widehat{\chi} \bmod q$. Splitting any convenient n uniquely as $mP_J \cdots P_1$ above, we obtain

$$(2.4) \quad \sum_{n \leq x \text{ conv}} \widehat{\chi}(\widehat{h}(n)) = \sum_{m \leq x} \widehat{\chi}(\widehat{h}(m)) \cdot \frac{1}{J!} \sum_{\substack{P_1, \dots, P_J > L_m \text{ distinct} \\ P_1 \cdots P_J \leq x/m}} \widehat{\chi}(\widehat{h}(P_1)) \cdots \widehat{\chi}(\widehat{h}(P_J)),$$

where we have replaced the ordering on $P_1, \dots, P_J$ in (2.1) by the weaker condition “ $P_1, \dots, P_J$ distinct” at the cost of the $J!$ factor. For each $i \in [J]$, we see by (1.4) that

$$(2.5) \quad \sum_{\substack{P_i \leq x/mP_1 \cdots P_{i-1}P_{i+1} \cdots P_J \\ P_i > L_m, r \neq i \implies P_r \neq P_i}} \widehat{\chi}(\widehat{h}(P_i)) = \sum_{L_m < P_i \leq x/mP_1 \cdots P_{i-1}P_{i+1} \cdots P_J} \widehat{\chi}(\widehat{h}(P_i)) + O(J) \\ = \alpha_{\widehat{\chi}} \sum_{\substack{P_i \leq x/mP_1 \cdots P_{i-1}P_{i+1} \cdots P_J \\ P_i > L_m, r \neq i \implies P_r \neq P_i}} \chi_0(h(P_i)) + O\left(J + \frac{Mx\mathcal{E}(y)}{mP_1 \cdots P_{i-1}P_{i+1} \cdots P_J}\right),$$

The sum of the O -term over all m and all the other P_r in (2.4) is $\ll x/y + J^{-1}Mx\mathcal{E}(y) \log x$. (To see this, we use the $(J-1)!$ from (2.4) to restore the ordering on $P_1, \dots, P_{i-1}, P_{i+1}, \dots, P_J$, and then note that $mP_1 \cdots P_{i-1}P_{i+1} \cdots P_J \leq x/y$.) Hence using (2.5) for $i = 1, \dots, J$ successively, we get from (2.4),

$$(2.6) \quad \sum_{n \leq x \text{ conv}} \widehat{\chi}(\widehat{h}(n)) = (\alpha_{\widehat{\chi}})^J \sum_{m \leq x} \widehat{\chi}(\widehat{h}(m)) \sum_{\substack{P_1, \dots, P_J \\ P_1 \cdots P_J \leq x/m \\ L_m < P_J < \cdots < P_1}} \chi_0(h(P_1 \cdots P_J)) + O\left(\frac{Jx}{y} + Mx\mathcal{E}(y) \log x\right).$$

The first sum on the right side is $\ll |\alpha_{\widehat{\chi}}|^J \#\{n \leq x \text{ conv} : (h(n), q) = 1\}$. Using this bound for all character tuples $\widehat{\chi} \bmod q$ for which $\mathfrak{f}(\widehat{\chi}) \nmid d$, and combining this with (2.3), we obtain Theorem 1.4. $\square$

Proof of Theorem 1.5. We first show the analogue of (1.6) for a general R . By the argument in the sentence after (2.1), we see that the n that are either z -smooth or have a repeated prime factor exceeding y contribute $O(\Psi(x, z) + x/y)$. Let $\sum_{n \leq x}^*$ denote a sum over $n \leq x$ with no repeated prime factor

exceeding y , that satisfy $P_R(n) > q$, $P_J(n) \leq y$, $P(n) > z$, and $h_i(n) \equiv a_i \pmod{q}$ for all i . Let $E_{r,s}$ be the number of n counted in $\sum_{n \leq x}^* 1$ with $\#\{p > q : p \parallel n\} = r$ and $\#\{p > q : p^2 \mid n\} = s$. Then

$$(2.7) \quad \sum_{n \leq x}^* 1 \leq \sum_{\substack{n \leq x \\ \#\{p > q : p \parallel n\} \geq R}}^* 1 + \sum_{\substack{n \leq x \\ \#\{p > q : p^2 \mid n\} \geq K}}^* 1 + \sum_{\substack{1 \leq r < R \\ 1 \leq s < K}} E_{r,s}.$$

Any n counted in $E_{r,s}$ is of the form $mp_1^{c_1} \cdots p_s^{c_s} A$, where $P(m) \leq q$, all $p_j > q$ are primes, all $c_j \geq 2$, where A is squarefree with $\Omega(A) = r$, $P^-(A) > q$, $P(A) > z$, and where all the factors are pairwise coprime. Thus $(h(m), q) = 1$, $h_i(A) \equiv a_i h_i(mp_1^{c_1} \cdots p_s^{c_s})^{-1} \pmod{q}$, and $\sum_{j=1}^s c_j \geq R - r$ (as $P_R(n) > q$). By (K_r) , the number of possible A is $\ll \xi_r(q; z) x (\log_2 x)^r / (mp_1^{c_1} \cdots p_s^{c_s} \log(z/q))$. Next,

$$(2.8) \quad \sum_{\substack{m: P(m) \leq q \\ (h(m), q) = 1}} \frac{1}{m} \leq \prod_{p \leq q} \left(1 + \sum_{r \geq 1} \frac{\mathbb{1}_{(h(p^r), q) = 1}}{p^r} \right) \ll \exp \left(\sum_{p \leq q} \frac{\mathbb{1}_{(h(p), q) = 1}}{p} \right),$$

and $\sum_{p_1, \dots, p_s > q} p_1^{-c_1} \cdots p_s^{-c_s} \ll q^{-(c_1 + \cdots + c_s - s)}$. The sum of $q^{-(c_1 + \cdots + c_s - s)}$ over all possible c_i is both $\ll q^s \prod_{j=1}^s (\sum_{c_j \geq 2} q^{-c_j}) \ll q^{-s}$, and $\ll \sum_{t \geq R-r} t^s / q^{t-s} \ll R^K / q^{R-r-s}$ (taking $t = \sum_{j=1}^s c_j$). Hence

$$E_{r,s} \ll R^K \frac{\xi_r(q; z)}{q^{\max\{s, R-r-s\}}} \cdot \frac{x (\log_2 x)^R}{\log(z/q)} \exp \left(\sum_{p \leq q} \frac{\mathbb{1}_{(h(p), q) = 1}}{p} \right).$$

The other two sums on the right of (2.7) are bounded analogously: For example, to bound the second sum, we write $n = mp_1^{c_1} \cdots p_K^{c_K} P$ with $P_J(m) \leq y$, $P > z$, and all $p_j > q$, $c_j \geq 2$. By Chebyshev's estimates, the count of P is $\ll x / (mp_1^{c_1} \cdots p_K^{c_K} \log z)$. This time in place of (2.8), we use

$$(2.9) \quad \sum_{\substack{m \leq x: P_J(m) \leq y \\ (h(m), q) = 1}} \frac{1}{m} \leq \sum_{\substack{A \leq x \\ \Omega(A) \leq J}} \frac{1}{A} \sum_{\substack{B: P(B) \leq y \\ (h(B), q) = 1}} \frac{1}{B} \ll (2 \log_2 x)^J \exp \left(\sum_{p \leq y} \frac{\mathbb{1}_{(h(p), q) = 1}}{p} \right),$$

where we have written $m = AB$ with $P(B) \leq y < P^-(A)$ (so that $P_J(n) \leq y$ forces $\Omega(A) \leq J$), bounded $\sum 1/B$ analogously to (2.8), and noted that $\sum 1/A \leq (1 + \sum_{p \leq x} 1/p)^J$. This establishes (1.6).

To show (1.7), we again impose the conditions $P(n) > z$ and that n has no repeated prime factor exceeding y . Write any such $n = mP$ with $\max\{z, P(m)\} < P \leq x/m$, so that $P_J(m) \leq y$ and $(h(m), q) = 1$. By Chebyshev's estimates, the number of P is $\ll x / (m \log z)$. Bounding $\sum 1/m$ by (2.9) yields (1.7). $\square$

Remark. Given any set $\mathcal{S}$ of K -tuples of characters mod q , applying (2.6) for $\hat{\chi} \notin \mathcal{S}$, we obtain a variant of (1.5) with the condition $\mathfrak{f}(\hat{\chi}) \nmid d$ replaced by $\hat{\chi} \notin \mathcal{S}$, and with the second term on the left replaced by $\varphi(q)^{-K} \sum_{\hat{\chi} \in \mathcal{S}} \bar{\chi}_1(a_1) \cdots \bar{\chi}_K(a_K) \sum_{n \leq x: P_J(n) > y} \hat{\chi}(\hat{h}(n))$. This estimate is uniform in all $\mathcal{S}$.

3. A "HALASZ-TYPE" CHARACTER SUM BOUND: PROOF OF PROPOSITION 1.6

Consider any $\hat{\chi} \pmod{q}$. By Halász's Theorem (as stated in [35, Corollary III.4.12]), we have

$$(3.1) \quad \sum_{n \leq x} \hat{\chi}(\hat{h}(n)) \ll \frac{x}{L} + x \exp \left(- \min_{-L \leq t \leq L} \mathcal{D}(x, t) \right),$$

uniformly in $x, L \geq 2$, where by Mertens' Theorem,

$$(3.2) \quad \mathcal{D}(x, t) := \sum_{p \leq x} \frac{1 - \operatorname{Re}(\hat{\chi}(\hat{h}(p)) p^{-it})}{p} \geq \log_2 x - \sum_{\substack{p \leq y \\ (h(p), q) = 1}} \frac{1}{p} - \left| \sum_{y < p \leq x} \frac{\hat{\chi}(\hat{h}(p))}{p^{1+it}} \right| + O(1).$$

Set $L := \log x$. Cover $(y, x]$ with disjoint subintervals $I_\eta := (\eta, \eta(1 + 1/\log^2 x)]$, such that the rightmost of these may jut out past x but not past $x(1 + 1/\log^2 x)$. Uniformly in $p \in I_\eta$ and $t \in [-L, L]$, we have $p = \eta(1 + O(1/\log^2 x))$, so that $p^{-(1+it)} = \eta^{-(1+it)}(1 + O(1/\log x))$. Combining these observations with the estimates $\sum_\eta \sum_{p \in I_\eta} 1/\eta \ll \sum_{p \leq 2x} 1/p \ll \log_2 x$ and $\sum_{x < p \leq x(1+1/\log^2 x)} 1/p \ll 1/\log^3 x$, we get

$$(3.3) \quad \sum_{y < p \leq x} \frac{\widehat{\chi}(\widehat{h}(p))}{p^{1+it}} = \sum_\eta \frac{1}{\eta^{1+it}} \sum_{p \in I_\eta} \widehat{\chi}(\widehat{h}(p)) + O\left(\frac{\log_2 x}{\log x}\right).$$

Using (1.4) to estimate $\sum_{p \in I_\eta} \widehat{\chi}(\widehat{h}(p))$ and noting that there are $O(\log^3 x)$ many I_η 's, we obtain

$$\sum_{y < p \leq x} \frac{\widehat{\chi}(\widehat{h}(p))}{p^{1+it}} = \alpha_{\widehat{\chi}} \sum_\eta \frac{1}{\eta^{1+it}} \sum_{p \in I_\eta} \chi_0(h(p)) + O\left(\frac{\log_2 x}{\log x} + M\mathcal{E}(y)(\log x)^3\right).$$

Using the observations before (3.3), we can replace the double sum above by $\alpha_{\widehat{\chi}} \sum_{y < p \leq x} \chi_0(h(p))/p^{1+it}$ upto an error $\ll \log_2 x / \log x$. Inserting the resulting estimate into (3.2), we obtain

$$(3.4) \quad \mathcal{D}(x, t) \geq \log_2 x - \sum_{\substack{p \leq y \\ (h(p), q)=1}} \frac{1}{p} - |\alpha_{\widehat{\chi}}| \sum_{\substack{y < p \leq x \\ (h(p), q)=1}} \frac{1}{p} + O(1 + M\mathcal{E}(y)(\log x)^3).$$

Finally, inserting this bound into (3.1) completes the proof of Proposition 1.6. $\square$

4. PREP FOR THEOREMS 1.2-1.3: CHARACTER SUMS, MODULE THEORY, ARITHMETIC GEOMETRY

Until section 6, we assume the set up of Theorems 1.2 and 1.3, and let $\mathbf{f} = \prod_{i=1}^K \mathbf{f}_i$, $\mathbf{F} = \prod_{i=1}^K \mathbf{F}_i$. Since $\alpha(q) > 0$, and $\alpha(q) = \prod_{\ell|q} \alpha(\ell)$ by the Chinese Remainder Theorem, we have $\alpha(\ell) > 0$ for each $\ell \mid q$. Hence $\alpha(\ell) = 1 - \frac{1}{\ell-1} \#\{u \in U_\ell : F(u) \equiv 0 \pmod{\ell}\} \geq 1 - \frac{\min\{\ell-2, D\}}{\ell-1}$. By Mertens' Theorem,

$$(4.1) \quad \alpha(q) \gg (\log_2(3q))^{-D} \text{ uniformly in } q \geq 1 \text{ s.t. } \alpha(q) \neq 0, \text{ where } D = \deg F = \sum_{i=1}^K \deg F_i.$$

For an odd prime power ℓ^e , let ψ_{ℓ^e} denote a character mod ℓ^e which generates the character group of U_{ℓ^e} . We will need the following version of the Weil bounds, as in [36, Corollary 2.3].

Proposition 4.1. *Let ℓ be a prime. Consider any $H \in \mathbb{Z}[T]$ which is not of the form $c \cdot G^{\ell-1}$ in $\mathbb{F}_\ell[T]$. Then $|\sum_{u \bmod \ell} \psi_\ell(H(u))| \leq (r-1)\sqrt{\ell}$ where r is the number of distinct roots of H (in $\mathbb{C}$).*

We will also need Cochrane's [5, Theorems 1.2, 7.1; eq. (1.15)] extension of this to higher prime powers.

Proposition 4.2. *Let ℓ be a prime, $H \in \mathbb{Z}[T]$ be nonconstant with $\text{ord}_\ell(H) = 0$. Define $t = \text{ord}_\ell(H')$ and $M = \max\{\text{mult}_\theta(\ell^{-t}H') : \theta \in \mathbb{F}_\ell, H(\theta) \neq 0\}$. Let $e \geq t+2$ be an integer.*

- (i) *If $\ell > 2$, then $\left| \sum_{u \bmod \ell^e} \psi_{\ell^e}(H(u)) \right| \leq \left(\sum_{\substack{\theta \in \mathbb{F}_\ell \\ H(\theta) \neq 0}} \text{mult}_\theta(\ell^{-t}H') \right) \ell^{t/(M+1)} \ell^{e(1-1/(M+1))}$.*
- (ii) *Assume that $e \geq t+3$ and let ψ be the character mod 2^e defined by $\psi(5) = \exp(2\pi i/2^{e-2})$ and $\psi(-1) = 1$. Then $\left| \sum_{u \bmod 2^e} \psi(H(u)) \right| \leq (12.5) 2^{t/(M+1)} 2^{e(1-1/(M+1))}$.*

The non-constantness of $H \in \mathbb{Z}[T]$ guarantees that $t = \text{ord}_\ell(H')$ is finite, so that $\ell^{-t}H'$ can be viewed as a nonzero element of $\mathbb{F}_\ell[T]$. Hence, $\text{mult}_\theta(\ell^{-t}H') \in \mathbb{Z}_{\geq 0}$ is the multiplicity of $\theta \in \mathbb{F}_\ell$ as a zero of $\ell^{-t}H'$.

The following result controls the quantities appearing in the above bounds.

Lemma 4.3. *There exists a constant $C := C(F_1, \dots, F_K) > 0$ satisfying all the following:*

- (1) *For any prime $\ell > C$ satisfying $\gcd(\ell - 1, \beta) = 1$, the only tuple $(A_1, \dots, A_K) \in [\ell - 1]^K$ for which $\prod_{i=1}^K F_i^{A_i}$ is of the form $c \cdot G^{\ell-1}$ in $\mathbb{F}_\ell[T]$ is $(A_1, \dots, A_K) = (\ell - 1, \dots, \ell - 1)$.*
- (2) *$\text{ord}_\ell(\sum_{i=1}^K A_i F'_i \prod_{j \neq i} F_j) \leq \mathbb{1}_{\ell \leq C} C$ for any prime ℓ and $A_1, \dots, A_K \in \mathbb{Z}^+$ with $\ell \nmid \gcd(A_1, \dots, A_K)$.*
- (3) *Consider any integer $r \geq 2$, any prime ℓ , and any $A_1, \dots, A_K \in \mathbb{Z}^+$ with $\ell \nmid \gcd(A_1, \dots, A_K)$. Let $H(T) := T^{\varphi(\ell^r)} \prod_{i=1}^K F_i(T)^{A_i}$, $\tau_\ell := \text{ord}_\ell(H')$, and $\tilde{F}(T) := \sum_{i=1}^K A_i F'_i(T) \prod_{j \neq i} F_j(T)$. Then*

$$(4.2) \quad \tau_\ell = \begin{cases} \text{ord}_\ell(\tilde{F}) = 0, & \text{if } \ell > C, \text{ and } r \geq 2 \\ \text{ord}_\ell(\tilde{F}) \leq C, & \text{if } \ell \leq C, \text{ ord}_\ell(F) = 0, \text{ and } r \geq C + 2. \end{cases}$$

In both cases above, we have $\text{mult}_\theta(\ell^{-\tau_\ell} H') = \text{mult}_\theta(\ell^{-\tau_\ell} \tilde{F})$ for any $\theta \in \mathbb{F}_\ell$ for which $\theta F(\theta) \neq 0$, so that $\sum_{\theta \in \mathbb{F}_\ell: H(\theta) \neq 0} \text{mult}_\theta(\ell^{-\tau_\ell} H') = \sum_{\theta \in \mathbb{F}_\ell: \theta F(\theta) \neq 0} \text{mult}_\theta(\ell^{-\tau_\ell} \tilde{F}) \leq \deg(\tilde{F}) \leq D - 1$.

Proof. Write each $F'_i \prod_{j \neq i} F_j =: \sum_{r=0}^{D-1} c_{i,r} T^r$ in $\mathbb{Z}[T]$. Let $\tilde{\beta} \in \mathbb{Z}^+$ be the largest invariant factor of the $D \times K$ matrix $M_0 := \left((c_{i,r})_{\substack{1 \leq i \leq K \\ 0 \leq r < D}} \right)^\top$. Fix $C > \max \left\{ 2, \tilde{\beta}, \prod_{j=1}^M |\text{disc}(G_j)| \cdot \prod_{1 \leq j \neq j' \leq M} |\text{res}(G_j, G_{j'})| \right\}$, such that C exceeds the sizes of the leading coefficients of all F_i and G_j . (Here $\{G_j\}_{j=1}^M \subset \mathbb{Z}[T]$ are as before Theorem 1.2; disc and res are the discriminant and resultant.) We show that any such C satisfies all the assertions. Note that $\prod_{j=1}^M G_j$ is separable over $\mathbb{F}_\ell$ (i.e. squarefree in $\overline{\mathbb{F}_\ell}[T]$) for all $\ell > C$.

Proof of (1). Consider any prime $\ell > C$ and any $(A_1, \dots, A_K) \in [\ell - 1]^K$ satisfying $\gcd(\ell - 1, \beta) = 1$ and $\prod_{i=1}^K F_i^{A_i} = c \cdot G^{\ell-1}$ for some $c \in U_\ell$ and $G \in \mathbb{F}_\ell[T]$. Since each $F_i = r_i \prod_{j=1}^M G_j^{\mu_{i,j}}$, we have $c \cdot G^{\ell-1} = \rho \prod_{j=1}^M G_j^{\sum_{i=1}^K \mu_{i,j} A_i}$ for some $\rho \in U_\ell$. Factoring out all leading coefficients and comparing exponents, we get $\sum_{i=1}^K \mu_{i,j} A_i \equiv 0 \pmod{\ell - 1}$ for all $j \in [M]$ (as $\prod_{j=1}^M G_j$ is squarefree in $\overline{\mathbb{F}_\ell}[T]$).

This gives the matrix congruence $E_0(A_1 \dots A_K)^\top \equiv 0 \pmod{\ell - 1}$. Let $\text{diag}(\beta_1, \dots, \beta_K)$ be the Smith normal form of E_0 , so each $\beta_i \in \mathbb{Z}^+$, $\beta_i \mid \beta_{i+1}$, and $\beta = \beta_K$. Consider invertible integer matrices P_0, R_0 s.t. $P_0 E_0 R_0 = \text{diag}(\beta_1, \dots, \beta_K)$. The last matrix congruence yields $\text{diag}(\beta_1, \dots, \beta_K) R_0^{-1} (A_1 \dots A_K)^\top \equiv 0 \pmod{\ell - 1}$. Since $\gcd(\ell - 1, \beta_K) = 1$, we have $\gcd(\ell - 1, \beta_1 \dots \beta_K) = 1$, so that writing out the last congruence elementwise, we get $R_0^{-1} (A_1 \dots A_K)^\top \equiv 0 \pmod{\ell - 1}$. Hence $(A_1 \dots A_K)^\top \equiv 0 \pmod{\ell - 1}$.

Proof of (2, 3). Since $(\prod_{i=1}^K F_i^{b_i})' = \sum_{i=1}^K b_i F'_i \prod_{j \neq i} F_j$ for any $\{b_i\}_{i=1}^K \subset \mathbb{Q}$, the multiplicative independence of $\{F_i\}_{i=1}^K$ forces the $\mathbb{Q}$ -linear independence of $\left\{ F'_i \prod_{j \neq i} F_j \right\}_{i=1}^K$. Hence the matrix M_0 above has rank K ($\leq D$), its Smith Normal Form has K positive diagonal entries, and $\tilde{\beta}$ is the last diagonal entry.

Let $\tilde{F} := \sum_{i=1}^K A_i F'_i \prod_{j \neq i} F_j$ and $d := \text{ord}_\ell(\tilde{F})$. Since $\tilde{F}(T) = \sum_{r=0}^{D-1} (\sum_{i=1}^K c_{i,r} A_i) T^r$, we have the matrix congruence $M_0(A_1 \dots A_K)^\top \equiv 0 \pmod{\ell^d}$. An argument analogous to that given in subpart (1) shows that if $\ell^d \nmid \tilde{\beta}$, then $\ell \mid (A_1, \dots, A_K)$, a contradiction. Thus $\ell^d \mid \tilde{\beta}$, i.e. $d \leq v_\ell(\tilde{\beta}) \leq \mathbb{1}_{\ell \leq C} C$, proving (2).

Now (4.2) follows from the identity $H'(T) = \varphi(\ell^r) T^{\varphi(\ell^r)-1} \prod_{i=1}^K F_i(T)^{A_i} + T^{\varphi(\ell^r)} \tilde{F}(T) \prod_{i=1}^K F_i(T)^{A_i-1}$. This identity also shows that in both cases of (4.2), we have $\ell^{-\tau_\ell} H'(T) \equiv T^{\varphi(\ell^r)} \cdot \ell^{-\tau_\ell} \tilde{F}(T) \cdot \prod_{i=1}^K F_i(T)^{A_i-1}$ in the ring $\mathbb{F}_\ell[T]$, which establishes the remaining assertion of subpart (3) of the lemma. $\square$

For any tuple of characters $\hat{\chi} = (\chi_1, \dots, \chi_K) \pmod{q}$, we define

$$(4.3) \quad \alpha_{\hat{\chi}} = (\alpha(q) \varphi(q))^{-1} \sum_{u \in U_q} \hat{\chi}(\hat{F}(u)) = (\alpha(q) \varphi(q))^{-1} \sum_{u \pmod{q}} \chi_0(u) \prod_{i=1}^K \chi_i(F_i(u)).$$

With B_0 and D as in Theorems 1.2 and 1.3, and with C as in the proof of Lemma 4.3,

$$(4.4) \quad \text{Fix } C_0 > \max\{B_0, C, (32D)^{4D+2}\}, \text{ and then fix an integer } \kappa > 100D(DC_0^{2C_0})^{4C_0}.$$

Let $\mathbf{Q}_0$ be the largest C_0 -smooth $(\kappa + 1)$ -free divisor of q , i.e. $\mathbf{Q}_0 = \prod_{\ell \leq C_0} \ell^{\min\{v_\ell(q), \kappa\}} \ll 1$.

Proposition 4.4. (1) We have $\sum_{\substack{\widehat{\chi} \bmod q \\ \mathfrak{f}(\widehat{\chi}) \nmid \mathbf{Q}_0}} |\alpha_{\widehat{\chi}}|^N = o(1)$ as $N \rightarrow \infty$, uniformly in all q .

(2) Bound (K_r) holds with $h_i = f_i$, where uniformly in all $q \in \mathbb{Z}^+$ and $z \in (q, x]$, we have

$$(4.5) \quad \xi_r(q; z) \ll \begin{cases} \varphi(q)^{-K} \exp(O((\log q)^{1-1/D})), & \text{for each fixed } r \geq KD + 1. \\ q^{-r/D+\epsilon}, & \text{for each fixed } r \leq KD \text{ and any fixed } \epsilon > 0. \\ q^{-1/D_{\min}} \cdot \log_2(3q), & \text{if } r = 1, \text{ where } D_{\min} = \min_i \deg(F_i). \end{cases}$$

(3) Bound (K_r) holds with $h_i = f_i$, where uniformly in all **squarefree** q and $z \in (q, x]$, we have

$$(4.6) \quad \xi_r(q; z) \ll \begin{cases} \varphi(q)^{-K} \exp(O(\sqrt{\log q})), & \text{for each fixed } r \geq 2K + 1. \\ q^{-r/2+\epsilon}, & \text{for each fixed } r \leq 2K \text{ and any fixed } \epsilon > 0. \\ D_{\min}^{\omega(q)} \cdot \varphi(q)^{-1}, & \text{if } r = 1. \\ \varphi(q)^{-1} \exp(O(\sqrt{\log q})), & \text{if } r = 2, K = 1, \text{ and } F_1 \text{ is not squarefull.} \end{cases}$$

Proof. As seen before (4.1), $\alpha(\ell) \geq 1 - \frac{\min\{\ell-2, D\}}{\ell-1} > 0$ for each $\ell \mid q$, which gives both $\text{ord}_\ell(F) = 0$, and

$$(4.7) \quad \alpha(\ell) \geq \frac{1}{2} \text{ if } \ell > C_0; \quad \alpha(\ell) \geq \frac{1}{C_0} \text{ if } \ell \leq C_0; \quad \alpha(2) = 1 \text{ if } 2 \mid q.$$

We assume in the entire proof that $D > 1$; the case $D = 1$ (i.e. $K = 1$ and F_1 linear) is much simpler. We first claim that for any tuple $\widehat{\chi} = (\chi_1, \dots, \chi_K)$ of Dirichlet characters mod q , we have

$$(4.8) \quad |\alpha_{\widehat{\chi}}| \leq \prod_{\substack{\ell > C_0 \\ \ell \mid \mathfrak{f}(\widehat{\chi})}} \left(4D \cdot \ell^{-v_\ell(\mathfrak{f}(\widehat{\chi}))/D} \right) \prod_{\substack{\ell \leq C_0 \\ v_\ell(\mathfrak{f}(\widehat{\chi})) \geq \kappa+1}} \left(2DC_0^{C_0+1} \cdot \ell^{-v_\ell(\mathfrak{f}(\widehat{\chi}))/D} \right).$$

To show this, we start by writing each $\chi_i = \prod_{\ell^e \parallel q} \chi_{i,\ell}$ for some character $\chi_{i,\ell} \bmod \ell^e$. Then with $\alpha_{\widehat{\chi},\ell} := (\alpha(\ell)\varphi(\ell^e))^{-1} \sum_{u \bmod \ell^e} \chi_{0,\ell}(u) \prod_{i=1}^K \chi_{i,\ell}(F_i(u))$, we have $\alpha_{\widehat{\chi}} = \prod_{\ell^e \parallel q} \alpha_{\widehat{\chi},\ell}$ by the Chinese Remainder Theorem. Now let $r_\ell := v_\ell(\mathfrak{f}(\widehat{\chi}))$, so $\ell^{r_\ell} = \text{lcm}[\mathfrak{f}(\chi_{1,\ell}), \dots, \mathfrak{f}(\chi_{K,\ell})]$. Let $\psi_{i,\ell}$ denote the character mod ℓ^{r_ℓ} inducing $\chi_{i,\ell}$, so that **at least one of $(\psi_{1,\ell}, \dots, \psi_{K,\ell})$ is primitive mod ℓ^{r_ℓ}** . Note that

$$(4.9) \quad \alpha_{\widehat{\chi},\ell} = (\alpha(\ell)\varphi(\ell^{r_\ell}))^{-1} \sum_{u \bmod \ell^{r_\ell}} \chi_{0,\ell}(u) \prod_{i=1}^K \psi_{i,\ell}(F_i(u)).$$

Since $|\alpha_{\widehat{\chi}}| = \prod_{\ell^e \parallel q} |\alpha_{\widehat{\chi},\ell}|$, our claim (4.8) will follow once we show that for each prime power $\ell^e \parallel q$,

$$(4.10) \quad |\alpha_{\widehat{\chi},\ell}| \leq \begin{cases} (4D) \cdot \ell^{-r_\ell/D} = (4D) \cdot \ell^{-v_\ell(\mathfrak{f}(\widehat{\chi}))/D}, & \text{if } \ell > C_0, r_\ell \geq 1. \\ (2DC_0^{C_0+1}) \cdot \ell^{-r_\ell/D} = (2DC_0^{C_0+1}) \cdot \ell^{-v_\ell(\mathfrak{f}(\widehat{\chi}))/D}, & \text{if } 2 < \ell \leq C_0, r_\ell \geq \kappa + 1. \\ (DC_0^{C_0+1}) \cdot 2^{-r_2/D} = (DC_0^{C_0+1}) \cdot 2^{-v_2(\mathfrak{f}(\widehat{\chi}))/D} & \text{if } \ell = 2 \text{ divides } q, \text{ and } r_2 \geq \kappa + 1. \end{cases}$$

First consider $\ell > 2$. Let ψ_ℓ be a character generating the character group mod ℓ^{r_ℓ} . Then each $\psi_{i,\ell} = \psi_\ell^{A_i}$ for some $A_i \in [\varphi(\ell^{r_\ell})]$. Since at least one of $(\psi_{i,\ell})_{i=1}^K$ is primitive mod ℓ^{r_ℓ} , we must have

$$(4.11) \quad (A_1, \dots, A_K) \not\equiv \begin{cases} 0 \pmod{\ell-1}, & \text{if } r_\ell = 1 \\ 0 \pmod{\ell}, & \text{if } r_\ell \geq 2. \end{cases}$$

By (4.9), $\alpha_{\widehat{\chi}, \ell} = (\alpha(\ell)\varphi(\ell^{r_\ell}))^{-1} \sum_{u \bmod \ell^{r_\ell}} \psi_\ell(H(u))$ with $H(T) := T^{\varphi(\ell^{r_\ell})} \prod_{i=1}^K F_i(T)^{A_i}$. We bound the sum as follows: If $\ell > C_0$ and $r_\ell = 1$, we use Proposition 4.1, Lemma 4.3(1) and (4.11), noting that the prime divisors $\ell > C_0 > \max\{C, B_0\}$ satisfy $\gcd(\ell - 1, \beta) = 1$ by hypothesis $IFH(F_1, \dots, F_K; B_0)$; hence $|\sum_{u \bmod \ell} \psi_\ell(H(u))| \leq D\ell^{1/2} \leq D\ell^{1-1/D}$ for all $\ell > C_0$. If either $\ell > C_0, r_\ell \geq 2$ or $2 < \ell \leq C_0, r_\ell \geq \kappa + 1$, then Lemma 4.3(3) and (4.11) give both $\tau_\ell := \text{ord}_\ell(H') \leq \mathbb{1}_{\ell \leq C_0} C_0 \leq r_\ell - 2$ (as $\kappa > C_0 + 2$ by (4.4)) and $\sum_{\theta \in \mathbb{F}_\ell: H(\theta) \neq 0} \text{mult}_\theta(\ell^{-\tau_\ell} H') \leq D - 1$; hence $|\sum_{u \bmod \ell^{r_\ell}} \psi_\ell(H(u))| \leq D(C_0^{C_0})^{\mathbb{1}_{\ell \leq C_0}} \ell^{r_\ell(1-1/D)}$ by Proposition 4.2(i). Combining all this with (4.7) and $\varphi(\ell^{r_\ell}) \geq \ell^{r_\ell}/2$, we get the first two bounds in (4.10).

Consider the third case of (4.10), i.e. $2 \mid q$, $\alpha(2) = 1$, $\text{ord}_2(F) = 0$, $r_2 \geq \kappa + 1 > C_0 + 2D + 4$ by (4.7) and (4.4). Recall that the character group mod 2^{r_2} is generated by the characters ψ, η with $\psi(5) = \exp(2\pi i/2^{r_2-2})$, $\psi(-1) = 1$, $\eta(5) = 1$, $\eta(-1) = -1$. Thus each $\psi_{i,2} = \psi^{A_i} \eta^{B_i}$ with $A_i \in [2^{r_2-2}]$, $B_i \in [2]$; here $\mathbf{2} \nmid (\mathbf{A}_1, \dots, \mathbf{A}_K)$ as some $(\psi_{i,2})_{i=1}^K$ is primitive mod 2^{r_2} . By (4.9), $\alpha_{\widehat{\chi}, 2} = \varphi(2^{r_2})^{-1} \sum_{u \bmod 2^{r_2}} \psi \left(\prod_{i=1}^K F_i(u)^{A_i} \right) \eta \left(u^2 \prod_{i=1}^K F_i(u)^{B_i} \right)$. Writing $u = 4v + \lambda$ and using $\mathfrak{f}(\eta) = 4$ gives (4.12)

$$\alpha_{\widehat{\chi}, 2} = \frac{1}{2^{r_2-1}} \sum_{\substack{\lambda = \pm 1 \\ \prod_{i=1}^K F_i(\lambda) \equiv 1 \pmod{2}}} \eta \left(\prod_{i=1}^K F_i(\lambda)^{B_i} \right) \cdot \frac{1}{4} \sum_{v \bmod 2^{r_2}} \psi(g_\lambda(v)), \text{ with } g_\lambda(T) := \prod_{i=1}^K F_i(4T + \lambda)^{A_i}.$$

Now $g'_\lambda(T) = 4\widetilde{F}(4T + \lambda) \prod_{i=1}^K F_i(4T + \lambda)^{A_i-1}$ where $\widetilde{F} := \sum_{i=1}^K A_i F'_i \prod_{j \neq i} F_j$. For each λ in (4.12), $\prod_{i=1}^K F_i(4T + \lambda) \equiv \prod_{i=1}^K F_i(\lambda) \equiv 1 \pmod{2}$, so $t_\lambda := \text{ord}_2(g'_\lambda) = 2 + \text{ord}_2(\widetilde{F}(4T + \lambda))$. From the last two sentences, we get both $\text{mult}_\theta(2^{-t_\lambda} g'_\lambda) = \text{mult}_\theta(2^{-(t_\lambda-2)} \widetilde{F}(4T + \lambda)) \leq \deg(\widetilde{F}) \leq D - 1$ for any $\theta \in \mathbb{F}_2$, and

$$t_\lambda = 2 + \text{ord}_2(\widetilde{F}(4T + \lambda)) \leq 2D + \text{ord}_2(\widetilde{F}) \leq 2D + C_0 \leq \kappa - 3 \leq r_2 - 3;$$

here the first inequality uses the fact that $\text{ord}_2(W(4T + \lambda)) \leq \text{ord}_2(W) + 2\deg(W)$ for any $W \in \mathbb{Z}[T]$ (which can be seen by a straightforward divisibility argument), and the second inequality uses Lemma 4.3(2). Hence, using Proposition 4.2(ii) and the observations $t_\lambda \leq 2D + C_0$ and $\max_{\theta \in \mathbb{F}_2} \text{mult}_\theta(2^{-t_\lambda} g'_\lambda) \leq D - 1$ above, we can bound each inner sum in (4.12), thus obtaining the last bound in (4.10).

Proof of Proposition 4.4 (1). Since $\#\{\widehat{\chi} \bmod q : \mathfrak{f}(\widehat{\chi}) \mid m\} \leq m^K$, (4.8) yields for any $N \in \mathbb{Z}^+$,

$$(4.13) \quad \sum_{\substack{\widehat{\chi} \bmod q \\ \mathfrak{f}(\widehat{\chi}) \nmid Q_0}} |\alpha_{\widehat{\chi}}|^N \leq \sum_{\substack{m \mid q \\ m \nmid Q_0}} m^K \prod_{\substack{\ell \mid m \\ \ell > C_0}} \left(4D \cdot \ell^{-v_\ell(m)/D} \right)^N \prod_{\substack{\ell \leq C_0 \\ v_\ell(m) \geq \kappa+1}} \left(2DC_0^{C_0+1} \cdot \ell^{-v_\ell(m)/D} \right)^N.$$

Since Q_0 is the largest C_0 -smooth $(\kappa + 1)$ -free divisor of q , any m in (4.13) either: (i) has $P(m) > C_0$ or (ii) has $P(m) \leq C_0$ but is not $(\kappa + 1)$ -free. Let E_1 denote the contribution of all m of type (i) to the rightmost sum in (4.13), and E_2 be that of type (ii). Writing each m in E_1 as $m = pM$ for some $p > C_0$,

$$(4.14) \quad E_1 \leq \sum_{p > C_0} \frac{(4D)^N}{p^{N/D-K}} \sum_{M \mid q} M^K \prod_{\substack{\ell \mid M \\ \ell > C_0}} \left(4D \cdot \ell^{-v_\ell(M)/D} \right)^N \prod_{\substack{\ell \leq C_0 \\ v_\ell(M) \geq \kappa+1}} \left(2DC_0^{C_0+1} \cdot \ell^{-v_\ell(M)/D} \right)^N.$$

As $N \rightarrow \infty$, the first sum above is at most $(4D)^N \sum_{p > C_0} p^{-N/2D} \leq (4DC_0^{-1/4D})^N \sum_{p > C_0} p^{-5} \leq (4DC_0^{-1/4D})^N = o(1)$ by (4.4). Since $\prod_{\ell \leq C_0} (1 + \sum_{1 \leq v \leq \kappa} \ell^{vK}) \ll 1$, the second sum in (4.14) is

$$\begin{aligned} &\ll \prod_{\substack{\ell \mid q \\ \ell > C_0}} \left(1 + \sum_{1 \leq v \leq v_\ell(q)} \frac{(4D)^N}{\ell^{v(N/D-K)}} \right) \cdot \prod_{\substack{\ell \mid q \\ \ell \leq C_0}} \left(1 + \sum_{\kappa+1 \leq v \leq v_\ell(q)} \frac{(2DC_0^{C_0+1})^N}{\ell^{v(N/D-K)}} \right) \\ &\ll \exp \left((4DC_0^{-1/4D})^N \sum_{\ell > C_0} \sum_{v \geq 1} \ell^{-5v} \right) \cdot \exp \left((2DC_0^{C_0+1} 2^{-\kappa/4D})^N \sum_{\ell \leq C_0} \sum_{v \geq 1} \ell^{-5v} \right) \ll 1 \end{aligned}$$

by (4.4) and the fact that $N/D - K \geq N/(2D) \geq 10$. Hence overall $E_1 = o(1)$ as $N \rightarrow \infty$.

Any m counted in E_2 is C_0 -smooth but not $(\kappa + 1)$ -free, and hence $m = AB$ for some $(\kappa + 1)$ -full integer $A > 1$ and some $(\kappa + 1)$ -free integer B . Since $B \leq \prod_{\ell \leq C_0} \ell^\kappa \ll_{C_0, \kappa} 1$, and $N \rightarrow \infty$, we have

$$\begin{aligned} E_2 &\ll \sum_{\substack{A > 1 : P(A) \leq C_0 \\ A \text{ is } (\kappa+1)\text{-full}}} \prod_{\ell \mid A} \frac{(2DC_0^{C_0+1})^N}{\ell^{v_\ell(A)(N/D-K)}} \leq \prod_{\ell \leq C_0} \left(1 + \sum_{v \geq \kappa+1} \frac{(2DC_0^{C_0+1})^N}{\ell^{Nv/2D}} \right) - 1 \\ &\leq \left| \exp \left(O \left(\sum_{\ell \leq C_0} \frac{(2DC_0^{C_0+1})^N}{\ell^{\kappa N/2D}} \right) \right) - 1 \right| \leq \left| \exp \left(O \left(\left(\frac{2DC_0^{C_0+1}}{2^{\kappa/2D}} \right)^N \right) \right) - 1 \right| \ll \left(\frac{2DC_0^{C_0+1}}{\kappa/2D} \right)^N, \end{aligned}$$

so that (4.4) yields $E_2 = o(1)$ as $N \rightarrow \infty$. Collecting all the estimates, we obtain subpart (1).

Proof of Proposition 4.4 (2). Fix any $r \in \mathbb{Z}^+$ and consider any $z \in (q, x]$. We may write

$$\sum_{\substack{P_1, \dots, P_r : P_1 \cdots P_r \leq x \\ P_1 > z, q < P_r < \dots < P_1 \\ (\forall i) f_i(P_1) \cdots f_i(P_r) \equiv a_i \pmod{q}}} 1 \leq \sum_{(v_1, \dots, v_r) \in \mathcal{V}_{r,K}(q; \hat{a})} \sum_{\substack{q < P_r < \dots < P_2 \leq x \\ (\forall j \geq 2) P_j \equiv v_j \pmod{q}}} \sum_{\substack{z < P_1 \leq x/P_2 \cdots P_r \\ P_1 \equiv v_1 \pmod{q}}} 1,$$

with $\mathcal{V}_{r,K}(q; \hat{a}) := \{(v_1, \dots, v_r) \in U_q^r : (\forall i) \prod_{j=1}^r F_i(v_j) \equiv a_i \pmod{q}\}$. By Brun–Titchmarsh, the innermost sum above is $\ll x/(\varphi(q)P_2 \cdots P_r \log(z/q))$; moreover, by Brun–Titchmarsh and partial summation, we have $\sum_{\substack{q < P_j \leq x \\ P_j \equiv v_j \pmod{q}}} 1/P_j \ll \log_2 x/\varphi(q)$ for each $j \in \{2, \dots, r\}$. Combining these two estimates,

$$(4.15) \quad \sum_{\substack{P_1, \dots, P_r : P_1 \cdots P_r \leq x \\ P_1 > z, q < P_r < \dots < P_1 \\ (\forall i) f_i(P_1) \cdots f_i(P_r) \equiv a_i \pmod{q}}} 1 \ll \frac{\#\mathcal{V}_{r,K}(q; \hat{a})}{\varphi(q)^r} \cdot \frac{x(\log_2 x)^{r-1}}{\log(z/q)}.$$

The bound $\#\mathcal{V}_{1,K}(q; \hat{a}) \ll q^{1-1/D_{\min}} \ll q^{-1/D_{\min}} \varphi(q) \log_2(3q)$, coming from Konyagin [14, 15], now yields the last bound in (4.5). Next, detecting the congruences $\prod_{j=1}^r F_i(v_j) \equiv a_i \pmod{q}$ by orthogonality,

$$(4.16) \quad \varphi(q)^K \cdot \frac{\#\mathcal{V}_{r,K}(q; \hat{a})}{\varphi(q)^r} = \alpha(q)^r \left| \sum_{\hat{\chi} \bmod q} \bar{\chi}_1(a_1) \cdots \bar{\chi}_K(a_K) (\alpha_{\hat{\chi}})^r \right| \leq \sum_{\hat{\chi} \bmod q} |\alpha_{\hat{\chi}}|^r$$

$$(4.17) \quad \leq \sum_{M \mid q} M^K \prod_{\substack{\ell \mid M \\ \ell > C_0}} \left(4D \cdot \ell^{-v_\ell(M)/D} \right)^r \prod_{\substack{\ell \mid M : \ell \leq C_0 \\ v_\ell(M) \geq \kappa+1}} \left(2DC_0^{C_0+1} \cdot \ell^{-v_\ell(M)/D} \right)^r,$$

where we used (4.8). The last sum above is precisely the second sum in (4.14), and hence again is

$$(4.18) \quad \ll \prod_{\substack{\ell \mid q \\ \ell \leq C_0}} \left(1 + \sum_{\kappa+1 \leq v \leq v_\ell(q)} \frac{(2DC_0^{C_0+1})^r}{\ell^{v(r/D-K)}} \right) \cdot \prod_{\substack{\ell \mid q \\ \ell > C_0}} \left(1 + \sum_{1 \leq v \leq v_\ell(q)} \frac{(4D)^r}{\ell^{v(r/D-K)}} \right).$$

If $r \geq KD + 1$, this is $\ll \exp(O(\sum_{\ell \mid q} \sum_{v \geq 1} \ell^{-v/D})) \leq \exp(O(\sum_{\ell \leq \omega(q)} \ell^{-1/D})) \leq \exp(O((\log q)^{1-1/D}))$.

If $r \leq KD$, then (4.18) is $\ll \prod_{\ell \mid q} O(v_\ell(q) \ell^{v_\ell(q)(K-r/D)}) \leq q^{K-r/D} e^{O(\omega(q))} \prod_{\ell \mid q} v_\ell(q) \ll_\varepsilon q^{K-r/D+\varepsilon}$. Inserting these two bounds into (4.17), and then into (4.15), we get the first two bounds in (4.5).

Proof of Proposition 4.4 (3). A simpler version of the argument for (4.8) gives $|\alpha_{\hat{\chi}}| \leq \prod_{\substack{\ell \nmid \hat{\chi} \\ \ell > C_0}} (4D \ell^{-1/2})$ for any $\hat{\chi} \bmod \text{squarefree } q$. With this, proceeding as above gives the first two bounds in (4.6). The third

bound in (4.6) follows from (4.15) and $\#\mathcal{V}_{1,K}(q;\widehat{a}) = \prod_{\ell|q} \#\mathcal{V}_{1,K}(\ell;\widehat{a}) \ll (D_{\min})^{\omega(q)}$ (by the Chinese Remainder Theorem and the fact that $\#\{u \in \mathbb{F}_\ell : W(u) = 0\} \leq \deg(W)$ for all $W \in \mathbb{Z}[T]$ and $\ell \gg_W 1$).

It only remains to show the fourth bound in (4.6), i.e. for $r = 2$, $K = 1$, and $F_1 \in \mathbb{Z}[T]$ not squarefull. Proceeding as in the first paragraph of the proof of Lemma 4.3, we can find a constant C_1 exceeding the size of the leading coefficient of F_1 , such that F_1 has at least one simple root in $\overline{\mathbb{F}}_\ell$ for all $\ell > C_1$. We claim that $F_1(X)F_1(Y) - w$ is irreducible in $\overline{\mathbb{F}}_\ell[X, Y]$ for all $\ell > C_1$ and all $w \in U_\ell$. Indeed, suppose

$$(4.19) \quad F_1(X)F_1(Y) - w = U(X, Y)V(X, Y) = \left(\sum_{0 \leq j \leq r} u_j(Y)X^j \right) \left(\sum_{0 \leq j \leq s} v_j(Y)X^j \right) \text{ in } \overline{\mathbb{F}}_\ell[X, Y],$$

where $U(X, Y) = \sum_{j=0}^r u_j(Y)X^j$, $V(X, Y) = \sum_{j=0}^s v_j(Y)X^j \in \overline{\mathbb{F}}_\ell[X, Y]$. For any root $\theta \in \overline{\mathbb{F}}_\ell$ of F_1 , taking $Y = \theta$ in (4.19) shows that $U(X, \theta)$ and $V(X, \theta)$ are constant. Hence if both $r, s > 0$, then $u_r(\theta) = v_s(\theta) = 0$, giving $\prod_{\theta: F_1(\theta)=0} (Y - \theta) \mid \gcd(u_r(Y), v_s(Y))$; but then comparing the highest degree terms in

X on the two extremes of (4.19) yields $\prod_{\theta: F_1(\theta)=0} (Y - \theta)^2 \mid F_1(Y)$, contradicting that F_1 has a simple root

in $\overline{\mathbb{F}}_\ell$. Thus $rs = 0$. Assuming $s = 0$, we have $V(X, Y) = v_0(Y)$ with $F_1(X)F_1(Y) - w = U(X, Y)v_0(Y)$. Substituting a root of F_1 for X shows that $V(X, Y) = v_0(Y)$ is constant, proving the claimed irreducibility.

The Hasse–Weil bound in [18, Corollary 2b] now gives $\#\mathcal{V}_{2,1}(\ell; w) \leq \varphi(\ell)(1 + O(\ell^{-1/2}))$ uniformly in $\ell > C_1$, $w \in U_\ell$. Hence from $\#\mathcal{V}_{2,1}(q; a) = \prod_{\ell|q} \#\mathcal{V}_{2,1}(\ell; a)$ and (4.15), we get the last bound in (4.6). $\square$

4.1. Setting up for Theorems 1.2 and 1.3. The following estimates will be used repeatedly.

Lemma 4.5. [27, Proposition 2.1, Lemma 2.4] *Fix any multiplicative function $w : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ for which there exists a nonconstant $W \in \mathbb{Z}[T]$ satisfying $w(p) = W(p)$ for all primes p . Fix any $K_0 > 0$. Then for all $X \gg 1$ and uniformly in $q \leq (\log x)^{K_0}$ with $\alpha_W(q) := \varphi(q)^{-1} \#\{u \in U_q : W(u) \in U_q\} > 0$, we have*

$$\sum_{\substack{n \leq X \\ (w(n), q)=1}} 1 = \frac{X}{(\log X)^{1-\alpha_W(q)}} \exp(O((\log_2(3q))^{O(1)})), \quad \sum_{\substack{p \leq X \\ (W(p), q)=1}} \frac{1}{p} = \alpha_W(q) \log_2 X + O((\log_2(3q))^{O(1)}).$$

To show Theorems 1.2 and 1.3, we invoke Theorems 1.4 and 1.5 with $h_i = f_i$, $q \leq (\log x)^{K_0}$, $d = Q_0$, $y = \exp((\log x)^{\epsilon/2})$, $J = \lfloor \log_3 x \rfloor$, $z = x^{1/\log_2 x}$ (with Q_0 as after (4.4), K_0 and ϵ as in Theorems 1.2-1.3, and $\epsilon := 1$ for Theorem 1.3). Writing $\sum_{y < p \leq Y} \widehat{\chi}(\widehat{f}(p)) = \sum_{u \in U_q} \widehat{\chi}(\widehat{F}(u)) \sum_{\substack{y < p \leq Y \\ p \equiv u \pmod{q}}} 1$, Siegel–Walfisz

gives (1.4) with $\alpha_{\widehat{\chi}}$ as in (4.3), $M = \varphi(q)$ and $\mathcal{E}(y) = \exp(-c\sqrt{\log y})$, with $c > 0$ an absolute constant.

Any quantity below is “negligible” if it is $o(\varphi(q)^{-K} \#\{n \leq x : (f(n), q) = 1\})$, with the number inside $o()$ estimated by Lemma 4.5. The right of (1.5) is negligible by Proposition 4.4(1). For $q \leq (\log x)^{K_0}$, the condition $P_J(n) > y$ in the *second* sum in (1.5) is ignorable upto negligible error, by (1.7) and the facts

$$(4.20) \quad \sum_{\substack{p \leq y \\ (f(p), q)=1}} \frac{1}{p} = \frac{\alpha(q)\epsilon}{2} \log_2 x + O((\log_2(3q))^{O(1)}); \quad \alpha(q) \gg \frac{1}{(\log_3 x)^D}; \quad \Psi(x, z) \leq \frac{x}{(\log x)^{(1+o(1)) \log_3 x}},$$

which use Lemma 4.5, (4.1), and known estimates on smooth numbers [4, p. 15], respectively. This gives

$$(4.21) \quad \sum_{\substack{n \leq x: P_J(n) > y \\ (\forall i) \ f_i(n) \equiv a_i \pmod{q}}} 1 - \left(\frac{\varphi(Q_0)}{\varphi(q)} \right)^K \sum_{\substack{n \leq x: (f(n), q)=1 \\ (\forall i) \ f_i(n) \equiv a_i \pmod{Q_0}}} 1 = o\left(\frac{1}{\varphi(q)^K} \sum_{\substack{n \leq x \\ (f(n), q)=1}} 1 \right).$$

For q in Theorem 1.2(i)-(ii), the condition $P_J(n) > y$ in (4.21) is ignorable upto negligible error by (1.6), (4.20), and the third bounds in (4.5), (4.6). For q, R in Theorem 1.3, the difference between the left of (1.2) and the first sum in (4.21) is the number $\#\{n \leq x : P_R(n) > q, P_J(n) \leq y, (\forall i) f_i(n) \equiv a_i \pmod{q}\}$ which is negligible by the last assertion of Theorem 1.5, (4.20), (4.5) and (4.6).⁶ The second asymptotic in (1.2) is clear, as the condition $P_R(n) \leq q$ forces $P_J(n) \leq y$ which can be dealt with by (1.7) and (4.20).

By (4.21) and the above observations, Theorems 1.2 and 1.3 would follow once we show that

$$(4.22) \quad \#\{n \leq x : (f(n), q) = 1, (\forall i) f_i(n) \equiv a_i \pmod{Q_0}\} \sim \varphi(Q_0)^{-K} \#\{n \leq x : (f(n), q) = 1\}.$$

This is clear if $Q_0 = 1$, or if $q^* := \prod_{\ell|q} \ell \ll 1$. (If $q^* \ll 1$, apply Theorem 1.1 to the integer $\text{lcm}[Q_0, q^*] \ll 1$). By orthogonality, (4.1) and Lemma 4.5, estimate (4.22) would follow once we show that: There exists a constant $\delta_0 := \delta_0(C_0, \kappa) < 1$ such that for all $q \leq (\log x)^{K_0}$ with $q \in \mathcal{Q}(f_1, \dots, f_K)$, $Q_0 > 1$ and $q^* \gg 1$,

$$(4.23) \quad \sum_{n \leq x} \mathbb{1}_{(f(n), q) = 1} \widehat{\psi}(\widehat{f}(n)) \ll \frac{x}{(\log x)^{1 - \delta_0 \alpha(q)}} \text{ for all nontrivial character tuples } \widehat{\psi} \pmod{Q_0}.$$

If $\widehat{\psi}(\widehat{F}(u))$ is *not* constant on $\{u \in U_{Q_0} : F(u) \in U_{Q_0}\}$, then (4.23) follows by applying Proposition 1.6 to the character tuple $\widehat{\chi} \pmod{q}$ induced by $\widehat{\psi}$, and noting that $|\alpha_{\widehat{\chi}}| = (\alpha(q)\varphi(q))^{-1} |\sum_{u \in U_q} \widehat{\chi}(\widehat{F}(u))| = (\alpha(Q_0)\varphi(Q_0))^{-1} |\sum_{u \in U_{Q_0}} \widehat{\psi}(\widehat{F}(u))|$ is bounded away from 1 by a *constant* (depending on C_0 and κ), since $Q_0 \ll_{C_0, \kappa} 1$. It thus only remains to show (4.23) when $\widehat{\psi}(\widehat{F}(u))$ is constant on $\{u \in U_{Q_0} : F(u) \in U_{Q_0}\}$.

5. MODIFYING THE LANDAU–SELBERG–DELANGE METHOD

We show (4.23) in the case when $\widehat{\psi}(\widehat{F}(u))$ takes a constant value $c_{\widehat{\psi}}$ on $\{u \in U_{Q_0} : F(u) \in U_{Q_0}\}$. Recall that $q \leq (\log x)^{K_0}$ lies $\mathcal{Q}(f_1, \dots, f_K)$, and has $q^* = \prod_{\ell|q} \ell \gg 1$ and $Q_0 > 1$. Note also that coprimality with q and q^* are equivalent, so that $\alpha(q^*) = \alpha(q) > 0$. We will modify the Landau–Selberg–Delange method, borrowing ideas from work of Scourfield [30]. We may assume throughout that $x \in \mathbb{Z}^+$.

5.1. Perron's formula and analytic continuations. We start by considering the Dirichlet series $\mathcal{F}(s) := \sum_{n \geq 1} \mathbb{1}_{(f(n), q) = 1} \widehat{\psi}(\widehat{f}(n)) / n^s$. By Perron's Formula [35, Theorem II.2.3], we have

$$(5.1) \quad \sum_{n \leq x} \mathbb{1}_{(f(n), q) = 1} \widehat{\psi}(\widehat{f}(n)) = \frac{1}{2\pi i} \int_{1 + \frac{1}{\log x} - iT}^{1 + \frac{1}{\log x} + iT} \frac{\mathcal{F}(s)x^s}{s} ds + O\left(x \sum_{n \geq 1} \frac{\mathbb{1}_{(f(n), q) = 1}}{n^{1 + \frac{1}{\log x}} (1 + T|\log(x/n)|)}\right)$$

uniformly in $T \geq 1$. The total contribution of all $n \notin (3x/4, 5x/4)$ to the O -term above is

$$(5.2) \quad \ll \frac{x}{T} \sum_{n \geq 1} \frac{1}{n^{1 + 1/\log x}} \ll \frac{x}{T} \exp\left(\sum_p \frac{1}{p^{1 + 1/\log x}}\right) \ll \frac{x}{T} \exp\left(\sum_{p \leq x} \frac{1}{p} + \sum_{j \geq 0} \exp(-2^j) \sum_{x^{2^j} < p \leq x^{2^{j+1}}} \frac{1}{p}\right),$$

which is $\ll x(\log x)/T$. Next, for any $n \in (3x/4, x - 1]$, we have $n = x - v$ for some $v \in [1, x/4] \cap \mathbb{Z}$, so that $|\log(x/n)| = -\log(1 - v/x) \gg v/x$. Thus, the contribution of all such n to the O -term in (5.1) is $\ll T^{-1} \sum_{1 \leq v < x/4} x/v \ll x(\log x)/T$; likewise, so is the contribution of all $n \in [x + 1, 5x/4)$. Hence

$$(5.3) \quad \sum_{n \leq x} \mathbb{1}_{(f(n), q) = 1} \widehat{\psi}(\widehat{f}(n)) = \frac{1}{2\pi i} \int_{1 + \frac{1}{\log x} - iT}^{1 + \frac{1}{\log x} + iT} \frac{\mathcal{F}(s)x^s}{s} ds + O\left(\frac{x \log x}{T}\right).$$

For $s \in \mathbb{C}$, write $\sigma := \text{Re}(s)$ and $t := \text{Im}(s)$. Define $\mathcal{L}_m(t) := \log(m(|t| + 1))$. Let $c_1 > 0$ be an absolute constant such that any $m \in \mathbb{Z}^+$, the product $\prod_{\eta \bmod m} L(s, \eta)$ has at most one zero ρ_e (the ‘‘Siegel zero’’) in the region $\sigma > 1 - c_1/\mathcal{L}_m(t)$; here $\rho_e \in \mathbb{R}$ is simple and is associated to a real character $\eta_e \pmod{m}$. In

⁶Note that $\max\{s + r/D, (KD + 1) - s - r + r/D\} \geq K + 1/2$ for $r \in [KD]$, and $\max\{s + r/2, (2K + 1) - s - r/2\} \geq K + 1/2$.

⁷i.e., for all tuples of characters $\psi = (\psi_1, \dots, \psi_K) \pmod{Q_0}$, except for the tuple of trivial characters $\pmod{Q_0}$

what follows, any term(s) involving ρ_e or η_e can be ignored if ρ_e does not exist. Now if $(\alpha(q^*), c_{\hat{\psi}}) \neq (1, 1)$, then we cut the complex plane along the ray $(-\infty, 1]$. If $(\alpha(q^*), c_{\hat{\psi}}) = (1, 1)$ and ρ_e exists, then we cut the complex plane along the ray $(-\infty, \rho_e]$. In the remaining case, we make no branch cut.

Lemma 5.1. *For $\sigma > 1$, we have $\mathcal{F}(s) = F_{\hat{\psi}}(s)G_{\hat{\psi}}(s)$, where $F_{\hat{\psi}}(s) := \prod_{\eta \bmod q^*} L(s, \eta)^{\alpha(q^*)c_{\hat{\psi}}\gamma(\eta)}$, $\gamma(\eta) := (\alpha(q^*)\varphi(q^*))^{-1} \sum_{u: uF(u) \in U_{q^*}} \bar{\eta}(u)$, and $G_{\hat{\psi}}$ is analytic on $\sigma > 1 - 1/\log q^*$ with $G_{\hat{\psi}}(s) \ll (\log_2 q^*)^3$ uniformly therein. Hence $\mathcal{F}(s)$ continues analytically into the region $\{s : \sigma > 1 - c_1/\mathcal{L}_{q^*}(t)\} - [1 - c_1, 1]$.*

Proof. By orthogonality mod q^* and the identity $\log L(s, \eta) = \sum_{p, v \geq 1} \eta(p^v)/vp^{vs}$ for $\sigma > 1$, we have

$$(5.4) \quad \sum_{p \equiv b \pmod{q^*}} \frac{1}{p^s} = \frac{1}{\varphi(q^*)} \sum_{\eta \bmod q^*} \bar{\eta}(b) \log L(s, \eta) - \sum_{\substack{p, v \geq 2 \\ p^v \equiv b \pmod{q^*}}} \frac{1}{vp^{vs}}.$$

Moreover, by the Euler product of the Dirichlet series $\mathcal{F}(s)$, we have for $\sigma > 1$,

$$(5.5) \quad \mathcal{F}(s) = \prod_{p: pF(p) \in U_{q^*}} \left(1 - \frac{c_{\hat{\psi}}}{p^s}\right)^{-1} \cdot \prod_p \left(1 + \sum_{v \geq 1} \frac{\mathbb{1}_{(f(p^v), q^*)=1} \hat{\psi}(\hat{f}(p^v))}{p^{vs}}\right) \left(1 - \frac{c_{\hat{\psi}} \mathbb{1}_{pF(p) \in U_{q^*}}}{p^s}\right).$$

Expanding the logarithm of the first product into Taylor series, and splitting the sum $\sum_{p: pF(p) \in U_{q^*}} p^{-s}$ as $\sum_{b \in U_{q^*}: F(b) \in U_{q^*}} \sum_{p \equiv b \pmod{q^*}} p^{-s}$, we see by (5.4) that this logarithm equals

$$(5.6) \quad \alpha(q^*)c_{\hat{\psi}} \sum_{\eta \bmod q^*} \gamma(\eta) \log L(s, \eta) + \left(\sum_{\substack{p, v \geq 2 \\ pF(p) \in U_{q^*}}} \frac{(c_{\hat{\psi}})^v}{vp^{vs}} - \sum_{\substack{p, v \geq 2 \\ pF(p^v) \in U_{q^*}}} \frac{c_{\hat{\psi}}}{vp^{vs}} \right).$$

Hence $\mathcal{F}(s) = F_{\hat{\psi}}(s)G_{\hat{\psi}}(s)$, where $G_{\hat{\psi}}(s)$ is the second product in (5.5) times the exponential of the parenthesized expression in (5.6). Now since $\sum_{p, v \geq 2} p^{-2v/3} \ll 1$, the last two sums in (5.6) are analytic functions of size $O(1)$ on $\sigma > 2/3$. The contribution of all $p \nmid q^*$ to the second product in (5.5) is exactly $\prod_{p \nmid q^*} (1 + \sum_{v \geq 2} p^{-vs} \{ \mathbb{1}_{(f(p^v), q^*)=1} \hat{\psi}(\hat{f}(p^v)) - c_{\hat{\psi}} \mathbb{1}_{(F(p)f(p^{v-1}), q^*)=1} \cdot \hat{\psi}(\hat{f}(p^{v-1})) \})$), which is analytic of size $O(1)$ on $\sigma > 2/3$. Finally for $\sigma > 1 - 1/\log q^*$, the total contribution of all $p \mid q^*$ to the second product in (5.5) has size $\leq \exp(\sum_{p \mid q^*} \sum_{v \geq 1} p^{-v\sigma}) \ll \exp(\sum_{p \mid q^*} p^{-1} \exp(\log p / \log q^*)) \ll \exp(e \sum_{p \leq p_{\omega(q^*)}} p^{-1}) \ll (\log_2 q^*)^3$ with $p_{\omega(q^*)}$ being the $\omega(q^*)$ -th prime. Collecting all these observations completes the proof. $\square$

Lemma 5.2. *Fix $W \in \mathbb{Z}[T]$ nonconstant. Consider squarefree $Q > 1$ with $\alpha_W(Q) := \varphi(Q)^{-1} \#\{u : uW(u) \in U_Q\} > 0$. Let $\gamma(\eta) = (\alpha_W(Q)\varphi(Q))^{-1} \sum_{u: uW(u) \in U_Q} \bar{\eta}(u)$. For s with $\sigma > 1 - c_1/2\mathcal{L}_Q(t)$, we have $|\sum_{\eta \bmod Q} \alpha_W(Q)\gamma(\eta)L'(s, \eta)/L(s, \eta) + \alpha_W(Q)/(s-1) - \alpha_W(Q)\gamma(\eta_e)/(s-\rho_e)| \ll \mathcal{L}_Q(t)$.*

This follows from Scourfield [30, Lemmas 15(i), 9(ii)], taking “ ξ ” there to be $e^{6\mathcal{L}_Q(t)}$, and using $L(z, \eta) = L(z, \eta') \prod_{\ell \mid Q} (1 - \eta'(\ell)/\ell^z)$ and $\gamma(\eta) = \gamma(\eta')$ when $\eta \bmod Q$ is induced by the primitive character η' . Note that if $\alpha_W(Q) = 1$ (which happens if, say, $\alpha_W(Q) > 0$ and $W(T) = c \cdot T^k$ for some constants c, k), then $\{u : uW(u) \in U_Q\} = U_Q$, so that $\gamma(\eta) = 0$ for all nontrivial $\eta \bmod Q$, and Lemma 5.2 is a classical result.

In our setting, we consider the function $H_{\hat{\psi}}(s) = F_{\hat{\psi}}(s) (s-1)^{\alpha(q^*)c_{\hat{\psi}}} (s-\rho_e)^{-\alpha(q^*)\gamma(\eta_e)c_{\hat{\psi}}}$ which analytically continues into $\sigma > 1 - c_1/\mathcal{L}_{q^*}(t)$. Lemma 5.2 yields $|H'_{\hat{\psi}}(z)/H_{\hat{\psi}}(z)| \ll \mathcal{L}_{q^*}(\text{Im}(z))$ uniformly in $z \in \mathbb{C}$ with $\text{Re}(z) > 1 - c_1/2\mathcal{L}_{q^*}(\text{Im}(z))$. Since $\log |H_{\hat{\psi}}(w)/H_{\hat{\psi}}(s)| \leq \left| \int_{\text{Re}(s)}^{\text{Re}(w)} |H'_{\hat{\psi}}(u+it)/H_{\hat{\psi}}(u+it)| du \right|$, we get

$$(5.7) \quad H_{\hat{\psi}}(s) \asymp H_{\hat{\psi}}(w), \text{ uniformly in } s, w \in \mathbb{C} \text{ having } \text{Im}(s) = \text{Im}(w) =: t$$

$$\text{and } 1 - c_1/2\mathcal{L}_{q^*}(t) \leq \text{Re}(s) \leq \text{Re}(w) \leq 1 + 100/\mathcal{L}_{q^*}(t).$$

5.2. The contour shift. We assume that $\rho_e \bmod q^*$ exists and $(\alpha(q^*), c_{\hat{\psi}}) \neq (1, 1)$, otherwise much simpler versions of our argument suffice. (We mention the modifications for the other possibilities at the end of this section.) By scaling c_1 if necessary, we may then *also* assume that $\rho_e > 1 - 5c_1/(24 \log q^*)$, and that $f(\eta_e)$ (which is squarefree) is large enough that it is divisible by a prime exceeding $D + 2$.

Define $T = \exp(\sqrt{\log x})$, $\tilde{\rho}_e = (2 + \rho_e)/3$, and $\vartheta(t) = 1 - c_1/(4\mathcal{L}_{q^*}(t))$. Let $\delta_e, \delta > 0$ be any parameters satisfying $\vartheta(0) < \rho_e - 2\delta_e$ and $\rho_e + 2\delta_e < \tilde{\rho}_e < 1 - 2\delta$. (We will eventually let $\delta, \delta_e \rightarrow 0+$.) Consider the contour Γ_1 consisting of the segment $\Gamma_2 = \{\sigma + iT : \sigma \in [\vartheta(T), 1 + 1/\log x]\}$, the curve $\Gamma_3 = \{\vartheta(t) + it : t \in [0, T]\}$, the segments $\Gamma_4 = [\vartheta(0), \rho_e - \delta_e]$, $\Gamma_6 = [\rho_e + \delta_e, \tilde{\rho}_e]$ and $\Gamma_7 = [\tilde{\rho}_e, 1 - \delta]$ all *above* the branch cut, the semicircle $\Gamma_5 = \{\rho_e + \delta_e \cdot e^{i\theta} : 0 < \theta < \pi\}$, the circle $\Gamma_8 = \{1 + \delta \cdot e^{i\theta} : -\pi < \theta < \pi\}$, and the reflections $\bar{\Gamma}_j$ of Γ_j about the real line (for $2 \leq j \leq 7$), with all components directed as in Figure 1.




FIGURE 1. The Contour Γ_1 in the case when the Siegel-zero ρ_e exists and $(\alpha(q^*), c_{\hat{\psi}}) \neq (1, 1)$

By Cauchy's integral theorem and the final assertion of Lemma 5.1, we now obtain from (5.3),

$$(5.8) \quad \sum_{n \leq x} \mathbb{1}_{(f(n), q)=1} \hat{\psi}(\hat{f}(n)) = \frac{1}{2\pi i} \int_{\Gamma_1} \frac{\mathcal{F}(s)x^s}{s} ds + O\left(\frac{x \log x}{T}\right).$$

We will show that all parts of Γ_1 except Γ_7 and $\bar{\Gamma}_7$ are negligible. The following bounds will be useful.

Proposition 5.3. (1) We have $|\mathcal{F}(s)| \ll \log x$ uniformly in $s \in \Gamma_2 \cup \Gamma_3 \cup \bar{\Gamma}_2 \cup \bar{\Gamma}_3$.
(2) For any fixed $\epsilon > 0$, we have $|H_{\hat{\psi}}(s)| \ll_{\epsilon} (\log x)^{\epsilon \alpha(q^*)/5}$ uniformly in real $s \in [1 - c_1/4 \log q^*, 1]$.

Proof. (1) Let $\mu(t) := 1 + c_1/(7\mathcal{L}_{q^*}(t))$. Consider any $s \in \Gamma_2 \cup \Gamma_3 \cup \bar{\Gamma}_2 \cup \bar{\Gamma}_3$. By Lemma 5.1, we have $\mathcal{F}(s) \ll (\log_2 q^*)^3 |F_{\hat{\psi}}(s)| \ll (\log_2 q^*)^3 |H_{\hat{\psi}}(s)| \cdot |s - 1|^{-\alpha(q^*)\text{Re}(c_{\hat{\psi}})} |s - \rho_e|^{\alpha(q^*)\text{Re}(\gamma(\eta_e)c_{\hat{\psi}})}$. Hence by (5.7),

$$(5.9) \quad \mathcal{F}(s) \ll (\log_2 q^*)^3 \cdot |H_{\hat{\psi}}(\mu(t) + it)| \cdot |s - 1|^{-\alpha(q^*)\text{Re}(c_{\hat{\psi}})} |s - \rho_e|^{\alpha(q^*)\text{Re}(\gamma(\eta_e)c_{\hat{\psi}})}.$$

Using the Dirichlet series of $\log L(\mu(t) + it, \eta)$ and interchanging sums, we see that $|F_{\hat{\psi}}(\mu(t) + it)| = |\exp(\sum_{\eta \bmod q^*} \gamma(\eta) \alpha(q^*) c_{\hat{\psi}} \log L(\mu(t) + it, \eta))| \leq \exp(\sum_{p, r \geq 1} r^{-1} p^{-r\mu(t)} |\sum_{\eta \bmod q^*} \alpha(q^*) \gamma(\eta) \eta(p^r)|)$. Now

$$\sum_{\eta \bmod q^*} \alpha(q^*) \gamma(\eta) \eta(p^r) = \frac{1}{\varphi(q^*)} \sum_{u: uF(u) \in U_{q^*}} \sum_{\eta \bmod q^*} \bar{\eta}(u) \eta(p^r) = \sum_{u: uF(u) \in U_{q^*}} \mathbb{1}_{p^r \equiv u \bmod q^*} = \mathbb{1}_{p^r F(p^r) \in U_{q^*}}.$$

Hence $|F_{\hat{\psi}}(\mu(t) + it)| \leq \exp(\sum_{p, r \geq 1} r^{-1} p^{-r\mu(t)}) = \zeta(\mu(t)) = 1/(\mu(t) - 1) + O(1) \ll \mathcal{L}_{q^*}(t)$. By the definition of $H_{\hat{\psi}}(\mu(t) + it)$, we thus obtain the following bound, which holds uniformly in all $t \in \mathbb{R}$.

$$(5.10) \quad |H_{\hat{\psi}}(\mu(t) + it)| \ll \mathcal{L}_{q^*}(t) |\mu(t) + it - 1|^{\alpha(q^*) \operatorname{Re}(c_{\hat{\psi}})} |\mu(t) + it - \rho_e|^{-\alpha(q^*) \operatorname{Re}(\gamma(\eta_e) c_{\hat{\psi}})}.$$

Finally, we observe that for both $\theta \in \{1, \rho_e\}$, we have $|s - \theta| \geq c_1/48 \mathcal{L}_{q^*}(t)$ for all $s = \sigma + it$ lying on $\Gamma_2 \cup \Gamma_3 \cup \bar{\Gamma}_2 \cup \bar{\Gamma}_3$. (This is clear if $|t| > 1/100$, or if $\theta = 1$, $|t| \leq 1/100$; if $\theta = \rho_e$ and $|t| \leq 1/100$, this follows from $\rho_e > 1 - 5c_1/(24 \log q^*)$.) The last observation easily yields $|s - \theta| \asymp |\mu(t) + it - \theta|$. Inserting this estimate and (5.10) into (5.9), we get $\mathcal{F}(s) \ll (\log_2 q^*)^3 \mathcal{L}_{q^*}(t) \ll (\log_3 x)^3 \log(q^* T) \ll \log x$.

(2) By (5.7) and (5.10), $|H_{\hat{\psi}}(s)| \ll |H_{\hat{\psi}}(\mu(0))| \ll (\log q^*)^3 (1 - \rho_e)^{-\alpha(q^*)}$ for $s \in [1 - c_1/4 \log q^*, 1]$. The desired bound now follows from $1 - \rho_e \gg (q^*)^{-\epsilon/20K_0} \gg (\log x)^{-\epsilon/20}$ (by Siegel's Theorem) and (4.1). $\square$

Proposition 5.4. *Set $I_j := \int_{\Gamma_j} \mathcal{F}(s) x^s / s \, ds$ and $\bar{I}_j := \int_{\bar{\Gamma}_j} \mathcal{F}(s) x^s / s \, ds$. Then*

$$(5.11) \quad |I_2| + |I_3| + |\bar{I}_2| + |\bar{I}_3| + |I_4 + \bar{I}_4| + |I_6 + \bar{I}_6| \ll x \exp(-(\log x)^{1/3}).$$

$$\lim_{\delta_e \rightarrow 0+} I_5 = \lim_{\delta_e \rightarrow 0+} \bar{I}_5 = \lim_{\delta \rightarrow 0+} I_8 = 0.$$

Proof. The bounds on $I_2, I_3, \bar{I}_2, \bar{I}_3$ are immediate by Proposition 5.3(1) and $|s| \gg t + 1$ for $s \in \Gamma_3 \cup \bar{\Gamma}_3$. To prove (5.11), we only show the bound for $I_4 + \bar{I}_4$; the argument for $I_6 + \bar{I}_6$ is analogous. For $s \in \Gamma_4$, we have $(s - 1)^{-\alpha(q^*) c_{\hat{\psi}}} \cdot (s - \rho_e)^{\alpha(q^*) \gamma(\eta_e) c_{\hat{\psi}}} = |1 - s|^{-\alpha(q^*) c_{\hat{\psi}}} \cdot |\rho_e - s|^{\alpha(q^*) \gamma(\eta_e) c_{\hat{\psi}}} \cdot e^{-i\pi \alpha(q^*) c_{\hat{\psi}}} \cdot e^{i\pi \alpha(q^*) \gamma(\eta_e) c_{\hat{\psi}}}$; the same holds for $s \in \bar{\Gamma}_4$ with the opposite sign of the power of e . Hence by the definition of $H_{\hat{\psi}}(s)$,

$$(5.12) \quad |I_4 + \bar{I}_4| \ll \int_{\vartheta(0)}^{\rho_e - \delta_e} \frac{|H_{\hat{\psi}}(u) G_{\hat{\psi}}(u)| x^u}{u} (1 - u)^{-\alpha(q^*) \operatorname{Re}(c_{\hat{\psi}})} \cdot (\rho_e - u)^{\alpha(q^*) \operatorname{Re}(\gamma(\eta_e) c_{\hat{\psi}})} \, du$$

$$\ll x^{\rho_e} (\log x)^{\alpha(q^*)/200} \cdot (\log_2 q^*)^3 \cdot (1 - \rho_e)^{-\alpha(q^*)} \int_{\vartheta(0)}^{\rho_e - \delta_e} (\rho_e - u)^{\alpha(q^*) \operatorname{Re}(\gamma(\eta_e) c_{\hat{\psi}})} \, du$$

by Lemma 5.1 and Proposition 5.3(2). By our hypothesis in the first paragraph of § 5.2, the conductor $f(\eta_e)$ has a prime factor $\ell_0 > D + 2$. As such, factoring $\eta_e = \prod_{\ell | q^*} \eta_{e, \ell}$ with $\eta_{e, \ell}$ a character mod ℓ ,

$$(5.13) \quad |\alpha(q^*) \gamma(\eta_e)| = \prod_{\ell | q^*} \frac{1}{\varphi(\ell)} \left| \sum_{\substack{u \in U_{\ell} \\ F(u) \in U_{\ell}}} \eta_{e, \ell}(u) \right| \leq \frac{1}{\ell_0 - 1} \left| - \sum_{\substack{u \in U_{\ell_0} \\ F(u) \equiv 0 \pmod{\ell_0}}} \eta_{e, \ell_0}(u) \right| \leq \frac{D}{D + 1}.$$

Hence the integral in (5.12) is at most $(\rho_e - \vartheta(0))^{1 + \alpha(q^*) \operatorname{Re}(\gamma(\eta_e) c_{\hat{\psi}})} / (1 + \alpha(q^*) \operatorname{Re}(\gamma(\eta_e) c_{\hat{\psi}})) \ll 1$. The bound on $|I_4 + \bar{I}_4|$ in (5.11) now follows from $1 - \rho_e \gg (\log x)^{-1/200}$ (Siegel's Theorem) and (4.1).

To show $\lim_{\delta_e \rightarrow 0+} I_5 = 0$, let $M_{\hat{\psi}}$ be the maximum of $|H_{\hat{\psi}}(s)|$ on a fixed small closed disk centered at ρ_e that is contained in the region $\sigma > 1 - c_1/4 \mathcal{L}_{q^*}(t)$. Then $M_{\hat{\psi}}$ is finite as $H_{\hat{\psi}}(s)$ is holomorphic on the region $\sigma > 1 - c_1/\mathcal{L}_{q^*}(t)$. Parametrizing Γ_5 as $s = \rho_e + \delta_e \cdot e^{i\theta}$ for $0 \leq \theta \leq \pi$, invoking Lemma 5.1 and arguing as above, we find that $|I_5| \ll M_{\hat{\psi}} (\log_2 q^*)^3 x^{\rho_e + \delta_e} \cdot (1 - \rho_e - \delta_e)^{-\alpha(q^*)} \cdot (\delta_e)^{1 + \alpha(q^*) \operatorname{Re}(\gamma(\eta_e) c_{\hat{\psi}})}$. Now

(5.13) yields $\lim_{\delta_e \rightarrow 0+} I_5 = 0$. The arguments for $\bar{I}_5$ and I_8 are analogous: For I_8 , we only need the fact that $\lim_{\delta \rightarrow 0+} \delta^{1-\alpha(q^*)\operatorname{Re}(c_{\hat{\psi}})} = 0$, since $(\alpha(q^*), c_{\hat{\psi}}) \neq (1, 1)$ as assumed in the first paragraph of § 5.2. $\square$

Letting $\delta, \delta_e \rightarrow 0+$ and invoking Proposition 5.4, we thus obtain from (5.8),

$$(5.14) \quad \sum_{n \leq x} \mathbb{1}_{(f(n), q)=1} \hat{\psi}(\hat{f}(n)) = \frac{1}{2\pi i} \int_{\Gamma_0} \frac{\mathcal{F}(s)x^s}{s} ds + O\left(x \exp(-(\log x)^{1/3})\right),$$

with Γ_0 being the contour being the union of the two horizontal segments joining the points 1 and $\tilde{\rho}_e$ above and below the branch cut, directed like Γ_7 . (It is easily seen that the integral in (5.14) converges.)

5.3. Bounding the remaining integral in (5.14). As before, we have for any fixed $\epsilon > 0$,

$$(5.15) \quad \left| \int_{\Gamma_0} \frac{\mathcal{F}(s)x^s}{s} ds \right| \ll \int_{\tilde{\rho}_e}^1 \frac{|H_{\hat{\psi}}(u)G_{\hat{\psi}}(u)|x^u}{u} (1-u)^{-\alpha(q^*)\operatorname{Re}(c_{\hat{\psi}})} \cdot (u-\rho_e)^{\alpha(q^*)\operatorname{Re}(\gamma(\eta_e)c_{\hat{\psi}})} du \\ \ll (\log x)^{\epsilon\alpha(q^*)/4} \int_{\tilde{\rho}_e}^1 |G_{\hat{\psi}}(u)| \cdot x^u (1-u)^{-\alpha(q^*)\operatorname{Re}(c_{\hat{\psi}})} du,$$

where we used Proposition 5.3(2) and $(u-\rho_e)^{\alpha(q^*)\operatorname{Re}(\gamma(\eta_e)c_{\hat{\psi}})} \ll (1-\rho_e)^{-\alpha(q^*)} \ll_{\epsilon} (\log x)^{\epsilon\alpha(q^*)/20}$.

Set $\epsilon_0 := \max_{1 < r \leq C_0^{C_0\kappa}} \cos(2\pi/r) < 1$. Since $Q_0 \in (1, C_0^{C_0\kappa}]$ and $c_{\hat{\psi}}$ is a $\varphi(Q_0)$ -th root of unity, we must have $\operatorname{Re}(c_{\hat{\psi}}) \leq \epsilon_0$ or $c_{\hat{\psi}} = 1$. If $\operatorname{Re}(c_{\hat{\psi}}) \leq \epsilon_0$, using Lemma 5.1 and writing $u = 1 - v/\log x$ yields, from (5.15),

$$(5.16) \quad \left| \int_{\Gamma_0} \frac{\mathcal{F}(s)x^s}{s} ds \right| \ll \frac{x(\log_3 x)^3}{(\log x)^{1-\alpha(q^*)(\epsilon_0+\epsilon/4)}} \cdot \Gamma(1-\alpha(q^*)\operatorname{Re}(c_{\hat{\psi}})) \ll \frac{x}{(\log x)^{1-\alpha(q^*)(\epsilon_0+\epsilon/3)}}$$

as the Γ -function is $O(1)$ on the interval $[1-\epsilon_0, 1]$. Inserting (5.16) into (5.14) proves (4.23) when $c_{\hat{\psi}} \neq 1$.

If $c_{\hat{\psi}} = 1$, then $\hat{\psi}(\hat{F}(u)) = 1$ on $\{u \in U_{Q_0} : F(u) \in U_{Q_0}\}$. Applying the definition of $\mathcal{Q}(f_1, \dots, f_K)$ to the characters mod q induced by $\hat{\psi} = (\psi_1, \dots, \psi_K)$, we get $\mathbb{1}_{(f(2^j), q)=1} \hat{\psi}(\hat{f}(2^j)) = -1$ for all $j \geq 1$. Isolating $p = 2$ from the second product in (5.5), we have $G_{\hat{\psi}}(s) = G_{\hat{\psi},1}(s)G_{\hat{\psi},2}(s)$, where $G_{\hat{\psi},2}(s) := (1 + \sum_{j \geq 1} \mathbb{1}_{(f(2^j), q)=1} \hat{\psi}(\hat{f}(2^j))/2^{js}) (1 - c_{\hat{\psi}} \mathbb{1}_{(2F(2), q^*)=1}/2^s)$ is analytic on $\sigma > 0$ and satisfies $G_{\hat{\psi},2}(1) = 0$, and where $G_{\hat{\psi},1}(s)$ is analytic on $\sigma > 1 - 1/\log q^*$ and satisfies $|G_{\hat{\psi},1}(s)| \ll (\log_2 q^*)^3$ uniformly therein. (The assertions on $G_{\hat{\psi},1}$ can be shown by following the arguments in Lemma 5.1.) Hence for all $u \in [\tilde{\rho}_e, 1]$,

$$(5.17) \quad |G_{\hat{\psi}}(u)| \ll (\log_2 q^*)^3 \cdot |G_{\hat{\psi},2}(1) - G_{\hat{\psi},2}(u)| \ll (\log_3 x)^3 \int_u^1 |G'_{\hat{\psi},2}(w)| dw \ll (\log_3 x)^3 (1-u).$$

Inserting this bound into (5.15), and then making the substitution $u = 1 - v/\log x$, we obtain

$$(5.18) \quad \left| \int_{\Gamma_0} \frac{\mathcal{F}(s)x^s}{s} ds \right| \ll \frac{x(\log_3 x)^3}{(\log x)^{2-\alpha(q^*)(1+\epsilon/4)}} \cdot \Gamma(2-\alpha(q^*)) \ll \frac{x}{(\log x)^{1-\alpha(q^*)\epsilon/3}},$$

which by (5.14) completes the proof of (4.23) in the remaining case $c_{\hat{\psi}} = 1$.

Finally, if $(\alpha(q^*), c_{\hat{\psi}}) = (1, 1)$ and ρ_e exists, then our branch cut is only along $(-\infty, \rho_e]$, so we replace all Γ_j and $\bar{\Gamma}_j$ (for $j \geq 5$) by a circle of radius δ_e centered at ρ_e , going from below to above the branch cut; we eventually let $\delta_e \rightarrow 0+$. If ρ_e doesn't exist, then for $(\alpha(q^*), c_{\hat{\psi}}) \neq (1, 1)$ we replace Γ_5 and $\bar{\Gamma}_5$ by straight segments above and below the branch cut (which is along $(-\infty, 1]$), while for $(\alpha(q^*), c_{\hat{\psi}}) = (1, 1)$, we take $\Gamma_1 = \Gamma_2 + \Gamma_3 + \bar{\Gamma}_2 + \bar{\Gamma}_3$ (there is no branch cut in this case). This establishes Theorems 1.2 and 1.3. $\square$

6. OPTIMALITIES IN THEOREMS 1.2 AND 1.3

Lemma 6.1. *Let $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ be multiplicative functions and $\{F_i\}_i \subset \mathbb{Z}[T]$ be nonconstant such that $\prod_{i=1}^K F_i$ is squarefree, and $f_i(p) = F_i(p)$ for all i and all primes p . There exists a constant $C_{\widehat{F}} > 0$ depending only on $\{F_i\}_i$ such that any $q \in \mathbb{Z}^+$ with $P^-(q) > C_{\widehat{F}}$ lies in $\mathcal{Q}(f_1, \dots, f_K)$.*

Proof. If $P^-(q) > D + 1$, then $\alpha(q) = \prod_{\ell|q} (1 - (\ell - 1)^{-1} \#\{u \in U_\ell : \prod_{i=1}^K F_i(u) \equiv 0 \pmod{\ell}\}) \geq \prod_{\ell|q} (1 - D/(\ell - 1)) > 0$. Moreover, if $\mathcal{T}(q) := \{(F_1(u), \dots, F_K(u)) : u \prod_{i=1}^K F_i(u) \in U_q\}$ generates U_q^K , then $q \in \mathcal{Q}(f_1, \dots, f_K)$ vacuously, i.e. there is no tuple of characters $\widehat{\chi} \neq (\chi_0, \dots, \chi_0) \pmod{q}$ for which $\prod_{i=1}^K \chi_i(F_i(u)) = 1$ on $\{u \in U_q : \prod_{i=1}^K F_i(u) \in U_q\}$. Hence, it suffices to show that there exists a constant $C_{\widehat{F}}$ such that $\mathcal{T}(q)$ generates U_q^K for all q with $P^-(q) > C_{\widehat{F}}$.

Now under the isomorphism $U_q^K \cong \prod_{\ell^e \parallel q} U_{\ell^e}^K$, the set $\mathcal{T}(q)$ maps to $\prod_{\ell^e \parallel q} \mathcal{T}(\ell^e)$. Thus if $\mathcal{T}(q)$ doesn't generate U_q^K , then by [23, Lemma 5.13], there is some $\ell^e \parallel q$ and characters $(\psi_1, \dots, \psi_K) \pmod{\ell^e}$, not all trivial, for which $\prod_{i=1}^K \psi_i(F_i(u))$ is constant on $\{u \in U_{\ell^e} : \prod_{i=1}^K F_i(u) \in U_{\ell^e}\}$. The lemma now follows from [21, Lemma 5]. $\square$

We show that the ranges of q in Theorem 1.2 are essentially optimal, in that “ $1 - \epsilon$ ” cannot be replaced by “ $1 + \epsilon$ ” in either subpart. In all our examples below, $\{F_i\}_{i=1}^K$ will be nonconstant and $\prod_{i=1}^K F_i$ will be separable, – guaranteeing that $\{F_i\}_{i=1}^K$ are multiplicatively independent, and that any $q \in \mathbb{Z}^+$ satisfies $IFH(F_1, \dots, F_K; B_0)$ for any $B_0 > 0$. Fix $r \in \mathbb{Z}^+$. Our K_0 will be fixed large enough in terms of $F_1, \dots, F_K$, and we will always have $q \leq (\log x)^{K_0}$.

Optimality in Theorem 1.2(i). Let $F_i(T) := (T - 1)^r + i$, and q be a perfect r -th power with $P^-(q) > \max\{C_{\widehat{F}}, 2K\}$ where $C_{\widehat{F}}$ is as in Lemma 6.1. Then $q \in \mathcal{Q}(f_1, \dots, f_K)$ and any prime $P \equiv 1 \pmod{q^{1/r}}$ satisfies $f_i(P) = F_i(P) \equiv i \pmod{q}$. By Siegel–Walfisz, $\#\{n \leq x : (\forall i) f_i(n) \equiv i \pmod{q}\} \gg x/(q^{1/r} \log x)$ uniformly in $q \leq (\log x)^{K_0}$. The last expression grows faster than the expected main term $\varphi(q)^{-K} \#\{n \leq x : (f(n), q) = 1\}$, once $q > (\log x)^{(1+\epsilon)\alpha(q)(K-1/r)^{-1}} = (\log x)^{(1+\epsilon)\alpha(q)(K-1/D_{\min})^{-1}}$ for any fixed ϵ . (To see this, use (4.1) and the first estimate in Lemma 4.5.)

Stronger optimality in Theorem 1.2(i) for $K > 1, D_{\min} = 1$ (i.e., at least one of the F_i is linear). We show that in this case, the range of q in Theorem 1.2(i) is optimal in a much stronger sense: It is optimal even if **all** the F_i are **any** pairwise coprime linear polynomials. Indeed, if $F_i(T) = c_i T + b_i$, then fixing any $b \in \mathbb{Z} \setminus \{0\}$ with $\prod_{i=1}^K F_i(b) \neq 0$, and taking any q with $P^-(q) > \max\{C_{\widehat{F}}, 1 + |b \prod_{i=1}^K F_i(b)|\}$, we note that any prime $P \equiv b \pmod{q}$ satisfies $f_i(P) \equiv F_i(b) \pmod{q}$. By Siegel–Walfisz, $\#\{n \leq x : (\forall i) f_i(n) \equiv F_i(b) \pmod{q}\} \gg x/\varphi(q) \log x$. This expression grows faster than the expected main term as soon as $q > (\log x)^{(1+\epsilon)\alpha(q)/(K-1)}$.

Optimality in Theorem 1.2(ii). Let $F_i(T) := \prod_{j=1}^r (T - 2j) + 2(2i - 1)$. Each F_i is irreducible by Eisenstein's criterion at 2, hence $\prod_{i=1}^K F_i$ is separable. Now let $q \leq (\log x)^{K_0}$ be a **squarefree** integer having $P^-(q) > \max\{C_{\widehat{F}}, 4Kr\}$. By the Chinese Remainder Theorem, the congruence $\prod_{j=1}^r (v - 2j) \equiv 0 \pmod{q}$ has exactly $r^{\omega(q)}$ distinct solutions $v \in U_q$. Hence by Siegel–Walfisz,

$$(6.1) \quad \sum_{\substack{n \leq x \\ (\forall i) f_i(n) \equiv 2(2i-1) \pmod{q}}} 1 \geq \sum_{\substack{n \leq x: P(n) > q \\ (\forall i) f_i(n) \equiv 2(2i-1) \pmod{q}}} 1 \geq \sum_{\substack{q < P \leq x \\ \prod_{j=1}^r (P-2j) \equiv 0 \pmod{q}}} 1 \gg \frac{r^{\omega(q)} x}{\varphi(q) \log x}.$$

By Lemma 4.5 and (4.1), the rightmost expression grows faster than $\varphi(q)^{-K} \#\{n \leq x : (f(n), q) = 1\}$ as soon as $q^{K-1} D_{\min}^{\omega(q)} = q^{K-1} r^{\omega(q)} > (\log x)^{(1+\epsilon)\alpha(q)}$ for any fixed $\epsilon > 0$.

For completeness, we construct arbitrarily large squarefree q satisfying the last inequality; in fact, we ensure that $r^{\omega(q)} > (\log x)^{(1+\epsilon)\alpha(q)}$. Indeed, let $q := \prod_{\max\{C_{\widehat{F}}, 4Kr\} < \ell \leq Y} \ell$, with $Y \leq (K_0/2) \log_2 x$ a parameter to be chosen later. Then $q \leq (\log x)^{K_0}$, $\omega(q) \geq Y/2 \log Y$, and by the Chinese Remainder Theorem and the Prime Ideal Theorem, we have $\alpha(q) \leq c/\log Y$ for some constant $c > 0$ depending only on $F_1, \dots, F_K$. The inequality $r^{\omega(q)} > (\log x)^{(1+\epsilon)\alpha(q)}$ is then ensured once we choose any $Y \in (4c \log_2 x / \log r, (K_0/2) \log_2 x)$. (We can fix $K_0 > 16c$ at the start.)

Optimality in Theorem 1.3. Note that by the $K = 1$ case of (6.1), the third value of R in Theorem 1.3 is optimal, in that it cannot be reduced to 1. We now show that the second value of R is “nearly optimal”, in that it cannot be reduced to $2K - 1$. With $F_i(T) = \prod_{j=1}^r (T - 2j) + 2(2i - 1)$ as above, we consider multiplicative functions $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ satisfying $f_i(p) = F_i(p)$ and $f_i(p^2) = 1$ for all i and all primes p . Consider any squarefree q as in the previous paragraph, and let $n = (p_1 \cdots p_{K-1})^2 P$ where p_i, P are primes satisfying $q < p_{K-1} < \cdots < p_1 < x^{1/4K} < x^{1/3} < P$ and $\prod_{j=1}^r (P - 2j) \equiv 0 \pmod{q}$. Then $f_i(n) \equiv 2(2i - 1) \pmod{q}$, so that by Siegel–Walfisz,

$$(6.2) \quad \sum_{\substack{n \leq x: P_{2K-1}(n) > q \\ (\forall i) f_i(n) \equiv 2(2i-1) \pmod{q}}} 1 \gg \frac{r^{\omega(q)} x}{\varphi(q) \log x} \sum_{\substack{p_1, \dots, p_{K-1} \in (q, x^{1/4K}) \\ p_1, \dots, p_{K-1} \text{ distinct}}} \frac{1}{(p_1 \cdots p_{K-1})^2},$$

where we have replaced the ordering condition on $p_1, \dots, p_{K-1}$ by a distinctness condition at the cost of $(K-1)!$. Note that $\sum_{p_1, \dots, p_{K-1} \in (q, x^{1/4K})} (p_1 \cdots p_{K-1})^{-2} \gg (\sum_{q < p \leq x^{1/4K}} p^{-2})^{K-1} \gg (q \log q)^{-(K-1)}$. On the other hand, the sum of $1/p_1 \cdots p_{K-1}$ over all primes $p_1, \dots, p_{K-1} > q$ when $p_i = p_j$ for some $i \neq j$, is $\ll (\sum_{p > q} p^{-4}) \cdot (\sum_{p > q} p^{-2})^{K-3} \ll q^{-K}$. Combining, we get

$$(6.3) \quad \sum_{\substack{n \leq x: P_{2K-1}(n) > q \\ (\forall i) f_i(n) \equiv 2(2i-1) \pmod{q}}} 1 \gg \frac{r^{\omega(q)}}{\varphi(q)(q \log q)^{K-1}} \cdot \frac{x}{\log x} \gg \frac{r^{\omega(q)}}{\varphi(q)^K (\log_2 x)^K} \cdot \frac{x}{\log x},$$

where we used $q \ll \varphi(q) \log_2 q$. The last expression above grows faster than $\varphi(q)^{-K} \#\{n \leq x : (f(n), q) = 1\}$ once $r^{\omega(q)} > (\log x)^{(1+\epsilon)\alpha(q)}$ for any fixed $\epsilon > 0$, which is already satisfied by our q .

7. THE GENERAL CASE

The hypothesis “ $\alpha(q) \neq 0$ ” played a key role in Theorem 1.1 and its uniform analogues Theorems 1.2 and 1.3: Not only did it prevent the input sets from becoming too sparse (via Lemma 4.5), but also guaranteed that the “polynomial-type” control on the f_i at the *primes* was most significant in a certain sense. It turns out that if $\alpha(q) = 0$, then the behavior of the f_i at a *higher prime power* becomes more significant. Narkiewicz’s general criterion in [21] allows for this flexibility, and we obtain best possible uniform analogues of his general criterion as well.

7.1. Extending Theorems 1.4 and 1.5 for certain sparse input sets. In this entire subsection, we fix $\nu \in \mathbb{Z}^+$ and $C > 0$. Given functions $h_1, \dots, h_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ and $q \in \mathbb{Z}^+$, we say that $(h_1, \dots, h_K)$ are **ν -supported (with respect to q)** if the ν -free part of any n satisfying $(h_1(n) \cdots h_K(n), q) = 1$ is at most C . Equivalently, any such n can be written as $n_1 n_2$, where $n_1 \leq C$ and n_2 is ν -free. In what follows, we will just write “ $(h_1, \dots, h_K)$ are ν -supported” to mean that they are ν -supported with respect to some q which will always be clear from the context.

Note that if $(h_1, \dots, h_K)$ are ν -supported, then they are also $(\nu - 1)$ -supported. By convention, any functions $(h_1, \dots, h_K)$ are 1-supported. The general paradigm in this entire section is that the control of the h_i at the ν -th powers of primes is most significant: All hypotheses, results and arguments are modified accordingly. In particular, taking $\nu = 1$ here recovers everything before.

In what follows, let $\Psi_\nu(x, z)$ denote the **number of ν -full z -smooth numbers up to x** ; we give a bound for this in Lemma 7.2. Given any ν -supported integer-valued multiplicative functions $(h_1, \dots, h_K)$,

and any $q, r \in \mathbb{Z}^+$ and $z > 0$, we **redefine** $\xi_r(\mathbf{q}; \mathbf{z}) := \xi_r(q; z; \{h_i\}_{i=1}^K) > 0$ be *any* parameter satisfying, for all $x \geq z$ and all coprime residues $a_1, \dots, a_K \pmod q$, the bound

$$(7.1) \quad \#\{P_1 \cdots P_r \leq x : P_i \text{ primes, } P_1 > z, q < P_r < \cdots < P_1, (\forall i) h_i(\mathbf{P}_1^\nu) \cdots h_i(\mathbf{P}_r^\nu) \equiv a_i \pmod q\} \\ \leq \xi_r(q; z) \cdot x(\log \log x)^r / \log(z/q).$$

The following theorem generalizes the main conclusion resulting from Theorems 1.4 and 1.5; the differences have been highlighted. We still use the notation preceding the statement of Theorem 1.4.

Theorem 7.1. *Consider any $q \in \mathbb{Z}^+$ with $q > \mathbf{C}$. Consider any ν -supported multiplicative functions $(h_1, \dots, h_K)$, any $M \geq 1$, $y \geq q$, and a decreasing function $\mathcal{E} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, such that*

$$(7.2) \quad \sum_{y \leq p \leq Y} \widehat{\chi}(\widehat{h}(\mathbf{p}^\nu)) = \alpha_{\widehat{\chi}} \sum_{y \leq p \leq Y} \chi_0(h(\mathbf{p}^\nu)) + O(MY\mathcal{E}(y))$$

for all $\widehat{\chi} \pmod q$ and all $Y \geq y$, where $\alpha_{\widehat{\chi}} \in \mathbb{C}$ lies in the unit disk.

Then for all d dividing q , all $x, z > 0$, and all $J, R \in \mathbb{Z}^+$ satisfying $y \leq z \leq x$ and $J \geq R$, we have

$$(7.3) \quad \sum_{\substack{n \leq x: P_R(n) > q \\ (\forall i) h_i(n) \equiv a_i \pmod q}} 1 - \left(\frac{\varphi(d)}{\varphi(q)} \right)^K \sum_{\substack{n \leq x: (h(n), q) = 1 \\ (\forall i) h_i(n) \equiv a_i \pmod d}} 1 \\ \ll \frac{1}{\varphi(q)^K} \sum_{\substack{\widehat{\chi} \pmod q \\ \dagger(\widehat{\chi}) \nmid d}} |\alpha_{\widehat{\chi}}|^J \sum_{\substack{n \leq x \\ (h(n), q) = 1}} 1 + \Psi_\nu(\mathbf{x}, \mathbf{z}) + J \left(\frac{x}{y} \right)^{1/\nu} + Mx^{1/\nu} \mathcal{E}(y) \log x \\ + \left\{ \left(\frac{\varphi(d)}{\varphi(q)} \right)^K + \xi_{\lceil R/\nu \rceil}(\mathbf{q}; \mathbf{z}) + R^K \sum_{\substack{1 \leq r < R/\nu \\ 1 \leq s < K\nu}} \frac{\xi_r(q; z)}{q^{\max\{\frac{s}{\nu}, \frac{R}{\nu} - r - s\}}} \right\} \frac{x^{1/\nu} (2 \log_2 x)^{R+J}}{\log(z/q)} \exp \left(\sum_{\substack{p \leq y \\ (h(\mathbf{p}^\nu), q) = 1}} \frac{1}{p} \right).$$

The analogue of (1.5) also holds, with the relevant replacements on the right hand side, and with the condition “ $P_J(n) > y$ ” replaced by “ $\mathbf{P}_{J\nu}(\mathbf{n}) > \mathbf{y}$ ”. The implied constants in both estimates depend only on $\nu, K, \mathbf{C}$ and the implied constant in (7.2).

Proof. The proofs are essentially analogous, so we mention the main changes. First, we define n to be “convenient” if the J largest **distinct** prime factors of n all exceed y and each appear exactly to the ν -th power in n . Thus, any convenient n can be uniquely written as $m(P_J \cdots P_1)^\nu$ where (2.1) holds. Once again, to prove the analogue of (1.5), it suffices to show the corresponding estimate with the condition “ $P_{J\nu}(n) > y$ ” replaced by “ n convenient”: This is because any inconvenient n having $P_{J\nu}(n) > y$ must be divisible by the $(\nu + 1)$ -th power of a prime exceeding y , and because

$$(7.4) \quad \sum_{\substack{n \leq x: (h(n), q) = 1 \\ \exists p > y \text{ s.t. } p^{\nu+1} | n}} 1 \leq \sum_{\substack{p > y \\ r \geq \nu+1}} \sum_{\substack{m \leq x/p^r \\ (h(m), q) = 1}} 1 \leq \sum_{\substack{p > y \\ r \geq \nu+1}} \sum_{m_1 \leq C} \sum_{\substack{m_2 \leq x/p^r \\ m_2 \text{ is } \nu\text{-full}}} 1 \ll x^{1/\nu} \sum_{\substack{p > y \\ r \geq \nu+1}} \frac{1}{p^{r/\nu}} \ll \left(\frac{x}{y} \right)^{1/\nu},$$

where we’ve used the Erdős–Székere estimate [10] on ν -full numbers. Now (2.2) and (2.3) hold as stated. The analogues of (2.4) and (2.5) hold with all x/m and $\widehat{\chi}(\widehat{h}(P_i))$ replaced by $x^{1/\nu}/m^{1/\nu}$ and $\widehat{\chi}(\widehat{h}(P_i^\nu))$; the O -term in (2.5) is replaced by $O(J + Mx^{1/\nu} \mathcal{E}(y)/m^{1/\nu} P_1 \cdots P_{i-1} P_{i+1} \cdots P_J)$. Reversing the splitting of convenient n , we see that this O -term summed over m and all other P_i is

$$\ll \sum_{n \leq x/y^\nu: (h(n), q) = 1} 1 + J^{-1} Mx^{1/\nu} \sum_{n \leq x/y^\nu: (h(n), q) = 1} n^{-1/\nu}.$$

The first of the two sums above is $\ll \sum_{n_1 \leq C} \sum_{\substack{n_2 \leq x/y^\nu \\ n_2 \text{ is } \nu\text{-full}}} 1 \ll x^{1/\nu}/y$. The second of the two sums is

$$(7.5) \ll \sum_{\substack{n \leq x \\ (h(n), q)=1}} \frac{1}{n^{1/\nu}} \leq \sum_{n_1 \leq C} \frac{1}{n_1^{1/\nu}} \sum_{\substack{n_2 \leq x \\ n_2 \text{ is } \nu\text{-full}}} \frac{1}{n_2^{1/\nu}} \ll \prod_{p \leq x} \left(1 + \sum_{r \geq \nu} \frac{1}{p^{r/\nu}}\right) \ll \exp\left(\sum_{p \leq x} \frac{1}{p}\right) \ll \log x.$$

We thus find that the analogue of (2.6) holds with all products $P_1 \cdots P_J$ replaced by $(P_1 \cdots P_J)^\nu$, and with all instances of x in the O -term replaced by $x^{1/\nu}$. This proves the analogue of (1.5).

Next to obtain (7.3), we show the analogue of (1.7), which is

$$(7.6) \quad \sum_{\substack{n \leq x: P_{J\nu}(n) \leq y \\ (h(n), q)=1}} 1 \ll \frac{x^{1/\nu}(2 \log_2 x)^J}{\log z} \exp\left(\sum_{p \leq y} \frac{\mathbb{1}_{(h(p^\nu), q)=1}}{p}\right) + \Psi_\nu(x, z) + \left(\frac{x}{y}\right)^{1/\nu}.$$

By (7.4), we may assume that the y -rough part of n is $(\nu+1)$ -free, so that (by definition of ν -supported and $y \geq q > C$) the y -rough part must be a perfect ν -th power. We may also assume that $P(n) > z$ since $\#\{n \leq x : P(n) \leq z, (h(n), q) = 1\} \leq \sum_{n_1 \leq C} \sum_{\substack{n_2 \leq C: P(n_2) \leq z \\ n_2 \text{ is } \nu\text{-full}}} 1 \ll \Psi_\nu(x, z)$. Under these two assumptions, $n = BA^\nu P^\nu$, with $P = P(n) > z$, $P(B) \leq y < P^-(A)$, $\Omega(A) \leq J$ and $(h(B), q) = 1$. Proceeding as in the proof of (1.7), and noting that $\sum_{B: P(B) \leq y} \mathbb{1}_{(h(B), q)=1}/B^{1/\nu} \ll \exp(\sum_{p \leq y} \mathbb{1}_{(h(p), q)=1}/p)$ by the method used in (7.5), we get (7.6).

Finally, we bound the n satisfying $P_R(n) > q$, $P_{J\nu}(n) \leq y$ and $h_i(n) \equiv a_i \pmod{q}$. Again, it suffices to bound the number $\sum_{n \leq x}^* 1$ of all $n \leq x$ with a $(\nu+1)$ -free y -rough part, satisfying $P_R(n) > q$, $P_{J\nu}(n) \leq y$, $P(n) > z$, and $h_i(n) \equiv a_i \pmod{q}$. Let $E_{r,s}$ be the number of n counted in $\sum_{n \leq x}^* 1$ with $\#\{p > q : p^\nu \parallel n\} = r$ and $\#\{p > q : p^{\nu+1} \mid n\} = s$. The analogue of (2.7) holds with “ R ”, “ K ”, “ $p \parallel n$ ”, “ $p^2 \mid n$ ” replaced by “ R/ν ”, “ $K\nu$ ”, “ $p^\nu \parallel n$ ”, “ $p^{\nu+1} \mid n$ ” respectively. (Here we noted that the q -rough part of n is ν -full, as $q > C$ and $h_1, \dots, h_K$ are ν -supported.)

The rest of the proof goes through by adapting the argument for Theorem 1.5: For instance, any n counted in $E_{r,s}$ is of the form $mp_1^{c_1} \cdots p_s^{c_s} A^\nu$, where $P(m) \leq q$, $p_j > q$, $c_j \geq \nu+1$, $c_1 + \cdots + c_s \geq R - \nu r$, and A is squarefree with $P^-(A) > q$, $P(A) > z$, $\Omega(A) = r$. Given $m, p_1, \dots, p_s$ and $c_1, \dots, c_s$, the number of A is at most $\xi_r(q; z) \cdot x^{1/\nu} (\log_2 x)^r / (m^{1/\nu} p_1^{c_1/\nu} \cdots p_s^{c_s/\nu} \log(z/q))$. The sum of $p_1^{-c_1/\nu} \cdots p_s^{-c_s/\nu}$ over all possible $p_j > q$ and over all $c_j \geq \nu+1$ satisfying $c_1 + \cdots + c_s \geq R - \nu r$, is $\ll R^K q^{-\max\{s/\nu, R/\nu - r - s\}}$ by the same argument as given in Theorem 1.5. On the other hand, since $h_1, \dots, h_K$ are ν -supported, we see that the sum of $1/m^{1/\nu}$ over all $P(m) \leq q$ satisfying $(f(m), q) = 1$ is at most $(\sum_{m_1 \leq C} 1/m) \cdot (\sum_{\substack{m_2 \leq x: P(m_2) \leq q \\ m_2 \text{ is } \nu\text{-full}}} 1/m_2) \ll \exp(\sum_{p \leq q} 1/p)$. $\square$

The following lemma gives a general uniform bound on $\Psi_\nu(x, z)$, the number of ν -full z -smooth numbers up to x . This generalizes some known bounds on smooth numbers, such as in [4, p. 15].

Lemma 7.2. *Fix $\nu \in \mathbb{Z}^+$. As $x, z \rightarrow \infty$, we have*

$$\Psi_\nu(x, z) \ll x^{1/\nu} (\log z) \exp(-u\nu^{-1} \log u + O(u \log_2(3u))),$$

uniformly for $(\log x)^{\max\{3, 2\nu\}} \leq z \leq x^{1/2}$. Here $u := \log x / \log z$.

Proof. This is a classic application of Rankin's trick. Let $\eta := (\log u)/(\log z) \leq \min\{1/3, 1/2\nu\}$. Noting that $\sum_p \sum_{r \geq \nu+1} p^{-r(1-\eta)/\nu} \ll_\nu \sum_p p^{-(1-\eta)(1+1/\nu)} \ll_\nu 1$, we have

$$\Psi_\nu(x, z) \leq \sum_{\substack{n: P(n) \leq z \\ n \text{ is } \nu\text{-full}}} \left(\frac{x}{n}\right)^{(1-\eta)/\nu} \leq x^{(1-\eta)/\nu} \prod_{p \leq z} \left(1 + \sum_{r \geq \nu} \frac{1}{p^{r(1-\eta)/\nu}}\right) \ll x^{(1-\eta)/\nu} \exp\left(\sum_{p \leq z} \frac{1}{p^{1-\eta}}\right).$$

Now $\sum_{p \leq z} p^{-(1-\eta)} = \log_2 z + \sum_{p \leq z} (\exp(\eta \log p) - 1)/p + O(1)$. In the last sum, the contribution of all $p \leq 2^{1/\eta}$ is $\ll \eta \sum_{p \leq 2^{1/\eta}} \log p/p \ll 1$, while that of $p \in (2^{1/\eta}, z]$ is at most $(\exp(\eta \log z) - 1) \sum_{2^{1/\eta} < p \leq z} 1/p \leq u(\log_2(3u) + O(1))$. Collecting all these bounds completes the proof. $\square$

We also mention the following generalization of Proposition 1.6 which will be useful to us soon.

Proposition 7.3. *In the setting of Theorem 7.1, we have for any $\hat{\chi} \bmod q$,*

$$(7.7) \quad \sum_{n \leq x} \hat{\chi}(\hat{h}(n)) \ll \frac{x^{1/\nu}}{\log x} \exp\left(\sum_{\substack{p \leq y \\ (h(p^\nu), q)=1}} \frac{1}{p} + |\alpha_{\hat{\chi}}| \sum_{\substack{y < p \leq x \\ (h(p^\nu), q)=1}} \frac{1}{p} + O(M(\log x)^3 \mathcal{E}(y))\right) \\ + \left(\frac{x}{y}\right)^{1/\nu} + x^{1/\nu} (\log x)^2 \exp(-u(2\nu)^{-1} \log u + O(u \log_2(3u))).$$

where $u = \log x / \log y$. The implied constant depends only on the implied constant in (1.4).

Proof. By ν -supportedness (and the fact that $y > C$), the y -rough part of any n on the left of (7.7) is ν -full. By (7.4), it also suffices to restrict the sum on the left of (7.7) to the n 's whose y -rough part is $(\nu+1)$ -free. These restricted n are thus of the form BMA^ν where $P(BM) \leq Y < P^-(A)$, where A is squarefree, M is ν -full, B is ν -free with $B \ll 1$, and $(h(B)h(M)h(A^\nu), q) = 1$.

If $M > x^{1/2}$, then $A \leq (x/BM)^{1/\nu} \leq x^{1/2\nu}/B^{1/\nu}$. Given B and A , the number of M is $\ll (x^{1/\nu}/B^{1/\nu}A) \cdot (\log y) \exp(-u(2\nu)^{-1} \log u + O(u \log_2(3u)))$ by Lemma 7.2, where $u = \log x / \log y$. Since $\sum 1/B \ll 1$ and $\sum 1/A \ll \log x$, the total contribution of n with $M > x^{1/2}$ is absorbed in the right of (7.7). On the other hand, the total contribution of all n having $M \leq x^{1/2}$ to the left hand side of (7.7) equals

$$(7.8) \quad \sum_{\substack{B \leq x: P(B) \leq y \\ B \text{ is } \nu\text{-free}}} \hat{\chi}(\hat{h}(B)) \sum_{\substack{M \leq x^{1/2}: P(M) \leq y \\ M \text{ is } \nu\text{-full}}} \hat{\chi}(\hat{h}(M)) \sum_{A \leq (x/BM)^{1/\nu}} \mu(A)^2 \mathbb{1}_{P^-(A) > y} \hat{\chi}(\hat{h}(A^\nu)).$$

Now we observe that uniformly in all B and M in (7.8), we have

$$(7.9) \quad \sum_{A \leq (x/BM)^{1/\nu}} \mu(A)^2 \mathbb{1}_{P^-(A) > y} \hat{\chi}(\hat{h}(A^\nu)) \ll \frac{(x/BM)^{1/\nu}}{\log x} \exp\left(|\alpha_{\hat{\chi}}| \sum_{\substack{y < p \leq x \\ (h(p^\nu), q)=1}} \frac{1}{p} + O(M(\log x)^3 \mathcal{E}(y))\right).$$

To show this, we proceed as in section 3: Writing $X := (x/BM)^{1/\nu}$, the analogue of (3.4) is

$$\mathcal{D}(X, t) = \sum_{p \leq X} \frac{1 - \operatorname{Re}(\mathbb{1}_{p > y} \hat{\chi}(\hat{h}(p^\nu)) p^{-it})}{p} \geq \log_2 X - |\alpha_{\hat{\chi}}| \sum_{y < p \leq X} \frac{\mathbb{1}_{(h(p^\nu), q)=1}}{p} + O(1 + M \mathcal{E}(y) (\log x)^3).$$

Inserting (7.9) into (7.8), recalling that $B \ll 1$, and observing that $\sum_{\substack{M: P(M) \leq y \\ M \text{ is } \nu\text{-full}}} \mathbb{1}_{(h(M), q)=1} / M^{1/\nu} \leq$

$\prod_{p \leq y} \left(1 + \sum_{r \geq \nu} \mathbb{1}_{(h(p^r), q)=1} / p^{r/\nu}\right) \ll \exp(\sum_{p \leq y} \mathbb{1}_{(h(p^\nu), q)=1} / p)$ completes the proof. $\square$

7.2. Narkiewicz's general criterion and its uniform analogues. As alluded to above, Narkiewicz's criterion allows for the possibility that the behavior of the f_i at the ν -th powers of primes is most significant, for any (fixed) $\nu \in \mathbb{Z}^+$. We thus start with multiplicative functions $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ for which there exist polynomials $\{F_{i,j}\}_{\substack{1 \leq i \leq K \\ 1 \leq j \leq V}} \subset \mathbb{Z}[T]$ satisfying $f_i(p^j) = F_{i,j}(p)$ for all $i \in [K]$, $j \in [V]$, and all primes p . (Thus, $F_{i,j}$ controls the behavior of f_i at the j -th powers of primes.) For each $j \in [V]$, define $\alpha_j(q) := \varphi(q)^{-1} \#\{u \in U_q : \prod_{i=1}^K F_{i,j}(u) \in U_q\}$. For a fixed $\nu \in [V]$, we say that q is **ν -admissible** if $\alpha_\nu(q) \neq 0$ but $\alpha_j(q) = 0$ for all $j < \nu$. (In other words, ν is the minimal index j for which $\prod_{i=1}^K F_{i,j}$ carries some unit mod q to some unit mod q .) Finally, let $\mathcal{Q}(\nu; f_1, \dots, f_K)$ be the set of all **ν -admissible moduli q** satisfying the following property

$$(7.10) \quad \text{For all } \hat{\chi} \neq (\chi_0, \dots, \chi_0) \bmod q \text{ satisfying } \prod_{i=1}^K \chi_i(F_{i,\nu}(u)) = 1 \text{ on its unit support}$$

$$\{u \in U_q : \prod_{i=1}^K F_{i,\nu}(u) \in U_q\}, \text{ there exists a prime } p \text{ satisfying } \sum_{m \geq 0} \hat{\chi}(\hat{f}(p^m)) p^{-m/\nu} = 0.$$

By the triangle inequality, such a prime $p \leq 2^\nu$; thus $\mathcal{Q}(1; f_1, \dots, f_K)$ is precisely the set $\mathcal{Q}(f_1, \dots, f_K)$ that we had been working with before. We now state Narkiewicz's general criterion.

Theorem 7.4. [21] *Fix ν -admissible q . Then $(f_1, \dots, f_K)$ are WUD mod q iff $q \in \mathcal{Q}(\nu; f_1, \dots, f_K)$.*

As the reader may expect by now, the polynomials $(F_{1,\nu}, \dots, F_{K,\nu})$ play the role of $(F_1, \dots, F_K)$ from the previous sections. Our first complete uniform analogue of Theorem 7.4 is thus the following generalization of Theorem 1.1, giving uniformity without any restrictions on inputs. All implied constants in what follows depend only on the polynomials $\{F_{i,j}\}_{\substack{1 \leq i \leq K \\ 1 \leq j \leq \nu}}$.

Theorem 7.5. *Fix $K_0 > 0$ and $\epsilon \in (0, 1)$, and assume that $F_{1,\nu}, \dots, F_{K,\nu}$ are multiplicatively independent. Then $(f_1, \dots, f_K)$ are WUD mod $q \leq (\log x)^{K_0}$ lying in $\mathcal{Q}(\nu; f_1, \dots, f_K)$ and satisfying $IFH(F_{1,\nu}, \dots, F_{K,\nu}; B_0)$, provided at least one of the following two conditions holds:*

- (i) $q \leq (\log x)^{(1-\epsilon)\alpha_\nu(q)(K-1/D_{\min})^{-1}}$, where $D_{\min} := \min\{\deg F_{i,\nu} : 1 \leq i \leq K\}$, **or**
- (ii) q is squarefree and $q^{K-1} D_{\min}^{\omega(q)} \leq (\log x)^{(1-\epsilon)\alpha_\nu(q)}$.

Again, both these ranges are essentially optimal. If $K > 1$ and $D_{\min} = 1$, then optimality holds in the stronger sense, i.e. even if $F_{1,\nu}, \dots, F_{K,\nu}$ are **any** pairwise coprime linear polynomials. The examples are analogous to those constructed in section 6; we discuss the changes in subsection § 7.5. As in Theorem 1.3, we need to restrict our inputs to restore full ‘‘Siegel–Walfisz uniformity’’.

Theorem 7.6. *Fix $K_0 > 0$ and assume that $F_{1,\nu}, \dots, F_{K,\nu}$ are multiplicatively independent. Then (1.2) holds uniformly in moduli $q \leq (\log x)^{K_0}$ lying in $\mathcal{Q}(\nu; f_1, \dots, f_K)$ and satisfying $IFH(F_{1,\nu}, \dots, F_{K,\nu}; B_0)$, and uniformly in $a_i \in U_q$. Here for a general q , we can take*

$$(7.11) \quad R = \begin{cases} \nu(KD + 1), & \text{if } \nu < D := \sum_{i=1}^K \deg F_{i,\nu}. \\ \text{the least integer exceeding } \nu(1 + (\nu + 1)(K - 1/D)), & \text{if } \nu \geq D. \end{cases}$$

If q is squarefree, we can take $R = \nu(K\nu + K - \nu + 1) + 1$, and we can also improve this to

$$(7.12) \quad R = \begin{cases} 2, & \text{if } K = \nu = 1 \text{ and } F_{1,\nu} \text{ is not squarefull.} \\ \nu(K\nu + K - \nu) + 1, & \text{if } \nu > 1 \text{ and at least one of } \{F_{i,\nu}\}_{i=1}^K \text{ is not squarefull.} \end{cases}$$

Both values of R in (7.12) are optimal in general; we discuss this in subsection § 7.5. The *second* value of R in (7.11) is also optimal, as shown by the example of $\sigma(n)$ in the discussion below. Note that the second values of R in (7.11) and (7.12) had no analogues in Theorem 1.3.

We mention concrete examples which can only be addressed with this greater generality. Recall the definition of $\mathcal{Q}(\nu; f_1, \dots, f_K)$ given in (7.10).

- Śliwa [34] showed that $\sigma(n)$ is WUD modulo *fixed* q iff $6 \nmid q$; in fact, $\mathcal{Q}(1; \sigma) = \{q : 2 \nmid q\}$ and $\mathcal{Q}(2; \sigma) = \{q : 2 \mid q, 3 \nmid q\}$. While Theorem 1.2 only dealt with odd q , Theorem 7.5 shows that $\sigma(n)$ is WUD modulo all $q \leq (\log x)^{(2-\delta)\tilde{\alpha}(q)}$ in $\mathcal{Q}(2; \sigma)$, as well as modulo all squarefree $q \leq (\log x)^{K_0}$ in $\mathcal{Q}(2; \sigma)$ satisfying $2^{\omega(q)} \leq (\log x)^{(1-\delta)\tilde{\alpha}(q)}$, where $\tilde{\alpha}(q) = \prod_{\ell|q: \ell \equiv 1 \pmod{3}} (1 - 2/(\ell - 1))$. (Here $\nu = D = 2$ and IFH is trivial as $T^2 + 1$ is separable.) The restriction on squarefree q is optimal by the example in [31, subsection 7.2].
- Moreover, Theorem 7.6 shows that $\sigma(n)$ is WUD modulo all $q \leq (\log x)^{K_0}$ in $\mathcal{Q}(2; \sigma)$ if we restrict to inputs n with $P_5(n) > q$ (respectively, $P_3(n) > q$ for squarefree q). By the examples constructed in [31, subsections 6.1 and 7.1], both these restrictions are optimal. (Note that our inputs n must be of the form m^2 or $2m^2$ since $\sigma(n)$ is odd, q being even.)
- Narkiewicz [22] studied the weak equidistribution of $\sigma_3(n) = \sum_{d|n} d^3$. He showed that $\mathcal{Q}(1; \sigma_3) = \{q : (q, 14) = 1\}$, $\mathcal{Q}(2; \sigma_3) = \{q : (q, 6) = 2\}$, and $\mathcal{Q}(\nu; \sigma_3) = \emptyset$ for all $\nu > 2$. Our results extend this to show that $\sigma(n)$ is WUD modulo all $q \leq (\log x)^{(3/2-\delta)\alpha(q)}$ in $\mathcal{Q}(1; \sigma_3)$, as well as modulo all squarefree $q \leq (\log x)^{K_0}$ in $\mathcal{Q}(1; \sigma_3)$ that satisfy $3^{\omega(q)} \leq (\log x)^{(1-\delta)\tilde{\alpha}(q)}$, where $\tilde{\alpha}(q) = \prod_{\ell \equiv 1 \pmod{3}} \ell|q (1 - 3/(\ell - 1)) \cdot \prod_{\ell \not\equiv 1 \pmod{3}} \ell|q (1 - 1/(\ell - 1))$. Uniformity is restored modulo all $q \leq (\log x)^{K_0}$ in $\mathcal{Q}(1; \sigma_3)$ by restricting to inputs n with $P_4(n) > q$ (resp. $P_2(n) > q$ for squarefree $q \leq (\log x)^{K_0}$ in $\mathcal{Q}(1; \sigma_3)$). Analogous results can be given for varying q in $\mathcal{Q}(2; \sigma_3)$; this genuinely requires the additional generality in this section.

Our results apply to any $\sigma_r(n)$ (for which all involved polynomials are separable), as well as to families like $(n, \varphi(n), \sigma(n))$ ($\sigma, \sigma_2, \sigma_3$), $(\varphi, \sigma, \sigma_2, \sigma_3)$, etc. In general, once we have an explicit description of the sets $\{\mathcal{Q}(\nu; f_1, \dots, f_K)\}_{\nu=1}^{\infty}$ (which is a “fixed-modulus” problem for each fixed ν), we have explicit analogues of Siegel–Walfisz for $(f_1, \dots, f_K)$ with optimal arithmetic restrictions. However, this problem of giving explicit descriptions of $\{\mathcal{Q}(\nu; f_1, \dots, f_K)\}_{\nu=1}^{\infty}$ is not solved in general. In fact, even just for the single function σ_r (with $r > 1$ fixed), the sets $\{\mathcal{Q}(\nu; \sigma_r)\}_{\nu=1}^{\infty}$ have only been computed upto $r \leq 200$ in works of Narkiewicz and Rayner [22, 28, 29]. Narkiewicz [20, 23] shows that for $(f_1, \dots, f_K)$ satisfying some natural additional constraints, we have $\mathcal{Q}(\nu; f_1, \dots, f_K) = \emptyset$ for all $\nu \gg 1$; he also gives algorithms to compute the nonempty $\mathcal{Q}(\nu; f_1, \dots, f_K)$.

7.3. Additional ingredients for Theorems 7.5 and 7.6. From now on, we are in the setting of Theorem 7.4 (as described before the statement of the theorem). Our first key observation:

(7.13) If q is ν -admissible, then $(f_1, \dots, f_K)$ are ν -supported.⁸

To see this, let $\mathcal{A}$ be the set of primes ℓ which satisfy $\alpha_j(\ell) = 0$ for some $j < \nu$. Note that since $\alpha_j(\ell) \geq 1 - (\ell - 1)^{-1} \sum_{i=1}^K \deg F_{i,j}$, the primes in the set $\mathcal{A}$ are no more than $1 + \max_{j \in [\nu]} \sum_{i=1}^K \deg F_{i,j} \ll 1$. To show (7.13), it thus suffices to show that for any n satisfying $(f(n), q) = 1$, the primes dividing the ν -free part of n must lie in $\mathcal{A}$. So assume by way of contradiction that $p^j \parallel n$ for some $p \notin \mathcal{A}$ and $j < \nu$. Then $\prod_{i=1}^K F_{i,j}(p) = f(p^j)$ divides $f(n)$. Since $j < \nu$ and q is ν -admissible, we have $\alpha_j(q) = 0$, so that by the Chinese Remainder Theorem, $\alpha_j(\ell_0) = 0$ for some prime $\ell_0 \mid q$. Then $\ell_0 \in \mathcal{A}$, so that $p \in U_{\ell_0}$. This forces $\ell_0 \mid \prod_{i=1}^K F_{i,j}(p) = f(p^j)$ as $\alpha_j(\ell_0) = 0$. Hence $\ell_0 \mid f(n)$, violating the fact that $(f(n), q) = 1$. This establishes (7.13).

⁸where we can take the “ C ” in the definition of ν -supported to be some constant depending only on $\{F_{i,j}\}_{1 \leq i \leq K, 1 \leq j \leq \nu}$.

We intend to apply Theorem 7.1. To set the stage, we need two additional ingredients. The first is an estimate on the input sets that generalizes and plays the role of the first estimate in Lemma 4.5.

Proposition 7.7. *As $x \rightarrow \infty$, we have uniformly in ν -admissible $q \leq (\log x)^{K_0}$,*

$$(7.14) \quad \#\{n \leq x : (f(n), q) = 1\} = \frac{x^{1/\nu}}{(\log x)^{1-\alpha_\nu(q)}} \exp(O((\log_2(3q))^{O(1)})).$$

Proof. The lower bound follows just by looking at the n of the form m^ν and applying the first estimate in Lemma 4.5 to the multiplicative function $m \mapsto f(m^\nu)$. To show the upper bound implied in (7.14), define $Y := \exp(\sqrt{\log x})$, and note that by the arguments leading to (7.8), it suffices to count the n of the form BMA^ν where $P(BM) \leq Y < P^-(A)$, where A is squarefree, M is ν -full and $M \leq x^{1/2}$, where B is ν -free with $B \ll 1$, and where $(f(B)f(M)f(A^\nu), q) = 1$. By [13, Theorem 01, p. 2], given B and M , the total number of A is

$$\sum_{A \leq (x/BM)^{1/\nu}} \mu(A)^2 \mathbb{1}_{P^-(A) > Y} \mathbb{1}_{(f(A^\nu), q) = 1} \ll \frac{(x/BM)^{1/\nu}}{\log x} \exp\left(\sum_{Y < p \leq x} \frac{\mathbb{1}_{(f(p^\nu), q) = 1}}{p}\right).$$

Bounding $\sum 1/M^{1/\nu}$ as at the end of the proof of Proposition 7.3, we see that the number of such n is $\ll (x^{1/\nu}/\log x) \exp(\sum_{p \leq x} \mathbb{1}_{(f(p^\nu), q) = 1}/p)$. This is absorbed in the right of (7.14) because

$$(7.15) \quad \sum_{\substack{p \leq X \\ (f(p^\nu), q) = 1}} \frac{1}{p} = \sum_{\substack{p \leq X \\ (F_{1,\nu}(p) \cdots F_{K,\nu}(p), q) = 1}} \frac{1}{p} = \alpha_\nu(q) \log_2 X + O((\log_2(3q))^{O(1)}) \text{ uniformly in } X \geq 3q,$$

by [27, Lemma 2.4]. This establishes the upper bound implied in (7.14). $\square$

The second ingredient is a partial improvement of (4.6) in the case $r = 3$.

Lemma 7.8. *Fix $\varepsilon > 0$. If at least one of $\{F_{i,\nu}\}_{i=1}^K$ is not squarefull (in $\mathbb{C}$), then*

$$(7.16) \quad \xi_3(q; z) \ll q^{-2+\varepsilon}, \text{ uniformly in squarefree } q \text{ and in } z \in (q, x].$$

Proof. The analogue of (4.15) holds with $f_i(P_j)$ replaced by $f_i(P_j^\nu)$ (and with $\mathcal{V}_{r,K}(q, \widehat{a})$ redefined accordingly), so we need only show that $\#\mathcal{V}_{3,K}(q; \widehat{a}) \ll \varphi(q)^{1+\varepsilon}$. As $\#\mathcal{V}_{3,K}(q; \widehat{a}) = \prod_{\ell|q} \#\mathcal{V}_{3,K}(\ell; \widehat{a})$ and $\omega(q) \ll \log q / \log_2 q$, it suffices to show (to complete the proof of the lemma) that

$$(7.17) \quad \#\mathcal{V}_{3,K}(\ell; \widehat{a}) \ll \varphi(\ell) \text{ uniformly in primes } \ell \gg 1 \text{ and } \widehat{a} = (a_1, a_2, a_3) \in U_\ell^3.$$

Assume w.l.o.g. that $F_{1,\nu}$ is not squarefull (in $\mathbb{C}$). As in the last bound in (4.6), our idea will be to embed $\mathcal{V}_{3,K}(\ell; \widehat{a})$ into the set $V_\ell(\mathbb{F}_\ell)$ of all $\mathbb{F}_\ell$ -rational points of the variety

$$V_\ell := \{(X, Y, Z) \in \overline{\mathbb{F}}_\ell^3 : F_{1,\nu}(X)F_{1,\nu}(Y)F_{1,\nu}(Z) - a_1 = F_{2,\nu}(X)F_{2,\nu}(Y)F_{2,\nu}(Z) - a_2 = 0\}.$$

Now, any sufficiently large prime ℓ satisfies all the following properties:

- (i) ℓ doesn't divide the leading coefficients of $F_{1,\nu}$ or $F_{2,\nu}$;
- (ii) $F_{1,\nu}$ is not squarefull in $\overline{\mathbb{F}}_\ell$;
- (iii) If $F_{1,\nu}^{c_1} \cdot F_{2,\nu}^{c_2}$ is constant in $\overline{\mathbb{F}}_\ell(T)$ for some $c_1, c_2 \in \mathbb{Z}$, then $\ell \mid (c_1, c_2)$.

We ensure (ii) by arguing as in the first paragraph of the proof of Lemma 4.3. To ensure (iii), we proceed as in the proof of Lemma 4.3(2) to see that the polynomials $(F'_{1,\nu}F_{2,\nu}, F_{1,\nu}F'_{2,\nu})$ must be $\mathbb{F}_\ell$ -linearly independent for all $\ell \gg 1$. (Basically, ℓ shouldn't divide the invariant factors of the 2-column matrix listing the coefficients of $F'_{1,\nu}F_{2,\nu}$ and $F_{1,\nu}F'_{2,\nu}$.) Now if $F_{1,\nu}^{c_1} \cdot F_{2,\nu}^{c_2}$ is constant in $\overline{\mathbb{F}}_\ell(T)$ for any such ℓ , then from its derivative, we get $c_1 F'_{1,\nu}F_{2,\nu} + c_2 F_{1,\nu}F'_{2,\nu} = 0$, forcing $\ell \mid (c_1, c_2)$.

Our next important observation is that if ℓ is large enough to satisfy (i)-(iii) above, then

$$(7.18) \quad F_1^*(X, Y, Z) := F_{1,\nu}(X)F_{1,\nu}(Y)F_{1,\nu}(Z) - a_1 \text{ is irreducible over } \overline{\mathbb{F}}_\ell$$

and does **not** divide $F_2^*(X, Y, Z) := F_{2,\nu}(X)F_{2,\nu}(Y)F_{2,\nu}(Z) - a_2$ in $\overline{\mathbb{F}}_\ell[X, Y, Z]$.

The first assertion can be shown by simply replicating the arguments given for the irreducibility of “ $F_1(X)F_1(Y) - w$ ” in the proof of Proposition 4.4(3). Next, assume for the sake of contradiction $F_2^* = F_1^*H_1$ for some $H_1 \in \overline{\mathbb{F}}_\ell[T]$. Comparing the coefficients of the monomial $Y^{r_1}Z^{r_2}$ of maximum total degree $r_1 + r_2$ on both sides of the last identity, we obtain $F_{1,\nu}(X) \mid F_{2,\nu}(X)$. Hence $F_{2,\nu} = F_{1,\nu}^m \cdot H$ for some $m \in \mathbb{Z}^+$ and $H \in \overline{\mathbb{F}}_\ell[X]$ such that $F \nmid H$. We will show that H must be constant.

To see this, note that by an easy induction argument, we have for all $t \in \{0, 1, \dots, m\}$,

$$(7.19) \quad F_1^* \text{ divides } G_t(X, Y, Z) := F_{1,\nu}(X)^{m-t}F_{1,\nu}(Y)^{m-t}F_{1,\nu}(Z)^{m-t}H(X)H(Y)H(Z) - a_1^{-t}a_2 \text{ in } \overline{\mathbb{F}}_\ell[X, Y, Z].$$

Indeed, since $G_0 = F_2^*$, this is tautological for $t = 0$. If (7.19) holds for some $t \leq m - 1$, then writing $G_t = F_1^*Q_t$ shows that $F_{1,\nu}(X)F_{1,\nu}(Y)F_{1,\nu}(Z) \mid (Q_t(X, Y, Z) - a_1^{-(t+1)}a_2)$. Defining Q_{t+1} by the relation $Q_t(X, Y, Z) - a_1^{-(t+1)}a_2 = F_{1,\nu}(X)F_{1,\nu}(Y)F_{1,\nu}(Z)Q_{t+1}(X, Y, Z)$, the identity $G_t = F_1^*Q_t$ leads to $G_{t+1} = F_1^*Q_{t+1}$, completing the induction and proving (7.19). Applying (7.19) for $t = m$, we get $H(X)H(Y)H(Z) - a_1^{-m}a_2 = F_1^*(X, Y, Z)Q(X, Y, Z)$ for some $Q \in \overline{\mathbb{F}}_\ell[X, Y, Z]$. Now if H were not constant, then again comparing the coefficient of the monomial $Y^{r_1}Z^{r_2}$ with maximal $r_1 + r_2$ on both sides of the last identity would show that $F \mid H$, a contradiction. Hence $F_{1,\nu}^{-m} \cdot F_{2,\nu} = H$ must be constant in $\overline{\mathbb{F}}_\ell(T)$, violating property (iii) from before. This proves (7.18).

The final stretch of the argument involves some key inputs from commutative algebra and algebraic geometry. Note that by (7.18), (F_2^*, F_1^*) form an $\overline{\mathbb{F}}_\ell[X, Y, Z]$ -regular sequence⁹: This is because if F_1^*Q lies in the ideal (F_2^*) , then (7.18) forces $Q \in (F_2^*)$. By [3, Proposition 1.2.14], it thus follows that the ideal (F_2^*, F_1^*) has height at least 2, so that the variety V_ℓ has Krull dimension at most 1 (since the ambient ring $\overline{\mathbb{F}}_\ell[X, Y, Z]$ has Krull dimension 3). Finally, the version of the Lang–Weil bound in [9, Claim 7.2] yields $\#V_\ell(\overline{\mathbb{F}}_\ell) \ll \ell$ for all $\ell \gg 1$, yielding the desired (7.17). $\square$

We are now ready to apply Theorem 7.1 in the context of Theorems 7.5 and 7.6. **Defining $\alpha_{\hat{\chi}}$ and y as in subsection § 4.1** (but with $F_{i,\nu}$ playing the role of “ F_i ” there), we see that (7.2) holds with same M and $\mathcal{E}(y)$ as in § 4.1. We thus apply Theorem 7.1 with the same J, z as in § 4.1, and with $d := Q_0$ (with Q_0 defined as after (4.4).) Proposition 4.4 (which just required multiplicative independence) continues to hold as stated, with $F_{i,\nu}$ playing the role of F_i , and with the $\xi_r(q, z)$ (re)defined in (7.1). This proposition and Lemma 7.8 bound all the relevant quantities on the right of (7.3). Lemma 7.2 and (7.15) estimate $\Psi_\nu(x, z)$ and $\sum_{p \leq y} \mathbb{1}_{(f(p^\nu), q)=1}/p$ respectively.

Collecting all these observations shows that the right of (7.3) is negligible compared to the main term $\varphi(q)^{-K} \#\{n \leq x : (f(n), q) = 1\}$. (For instance, to show that the values of R in (7.11) work, we just need to check that $\max\{s/\nu + r/D, R/\nu - r + r/D - s\} > K$. If $s/\nu + r/D \leq K$, then $R/\nu - r + r/D - s \geq R/\nu - K\nu + ((\nu + 1)/D - 1)r$, and this last quantity attains its minimum over all $r \in [1, R/\nu]$, either at $r = 1$ or $r = R/\nu$, depending on whether $\nu \geq D$ or $\nu < D$. The corresponding check for squarefree q is just more mechanically tedious; one needs only use the best available bound out of (7.16) and the analogues of (4.6) for each $r \in [1, R/\nu]$.)

⁹Following [3, Chapter 1], given a module M over a ring R , we say that $x_1, \dots, x_n \in R$ form an M -regular sequence if $M(x_1, \dots, x_n) \neq M$, and if for each $i \geq 1$, the multiplication-by- x_i map is injective in the quotient module $M/(x_1, \dots, x_{i-1})M$. (For $i = 1$, this means that the map $M \rightarrow M : \lambda \mapsto x_i\lambda$ is injective.)

Hence, proving Theorems 7.5 and 7.6 once again comes down to showing the following analogue of (4.23): There exists a constant $\delta_0 := \delta_0(C_0, \kappa) < 1$ such that

$$(7.20) \quad \sum_{n \leq x} \mathbb{1}_{(f(n), q)=1} \widehat{\psi}(\widehat{f}(n)) \ll \frac{x^{1/\nu}}{(\log x)^{1-\delta_0 \alpha(q)}} \text{ for all nontrivial tuples } \widehat{\psi} \bmod Q_0,$$

uniformly in $q \leq (\log x)^{K_0}$ lying in $\mathcal{Q}(f_1, \dots, f_K)$, having $Q_0 > 1$ and having sufficiently large radical. Once again, Proposition 7.3 yields (7.20) when $\prod_{i=1}^K \psi_i(F_{i,\nu}(u))$ is not constant on its unit support $\{u \in U_{Q_0} : \prod_{i=1}^K F_{i,\nu}(u) \in U_{Q_0}\}$. Hence, we need only show (7.20) when $\prod_{i=1}^K \psi_i(F_{i,\nu}(u))$ takes a constant value $c_{\widehat{\psi}}$ on its unit support. To do this, we just need to modify the arguments in section 5 as follows.

7.4. The additional analytic ingredients. Our first (and most nontrivial) modification is that in order to get the analogue of (5.3), we need the following observation.

Lemma 7.9. *For any $x \geq 2$, there exists an integer $X \in [x, x + x/\log^2 x]$ such that*

$$\sum_{\substack{3X/4 < n < 5X/4 \\ n \neq X}} \frac{\mathbb{1}_{(f(n), q)=1}}{|\log(X/n)|} \ll X^{1/\nu} \log X.$$

Proof. The lemma would follow once we show that with $h := x/\log^2 x$, we have

$$(7.21) \quad \sum_{X \in \mathbb{Z} \cap [x, x+h)} \sum_{\substack{3X/4 < n < 5X/4 \\ n \neq X}} \frac{\mathbb{1}_{(f(n), q)=1}}{|\log(X/n)|} \ll x^{1/\nu} h \log x.$$

We write the total double sum on the left as $S_1 + S_2$, where S_1 denotes the contribution of $n \in (3X/4, X-1]$. Then for any n in S_1 , we can write $n = X - r$, where $r \in [1, X/4] \subset [1, (x+h)/4]$. By (7.13), we have $n = n_1 n_2$ where $n_1 \ll 1$ and n_2 is ν -full. Hence, $n_2 = (X - r)/n_1 \in [(x - r)/n_1, (x + h - r)/n_1]$. Also $|\log(X/n)| = -\log(1 - r/X) \gg r/X \gg r/x$. Combining these,

$$S_1 \leq x \sum_{1 \leq r \leq \frac{x+h}{4}} \frac{1}{r} \sum_{n_1 \ll 1} \sum_{\substack{\frac{x-r}{n_1} \leq n_2 < \frac{x+h-r}{n_1} \\ n_2 \text{ is } \nu\text{-full}}} 1 \ll x \log x \left(x^{1/\nu} \cdot \frac{h}{x} + x^{1/(\nu+1)} \right) \ll x^{1/\nu} h \log x,$$

where we have noted that the sum on n_2 above is $\ll x^{1/\nu} \{(1 + O(h/x))^{1/\nu} - 1\} + x^{1/(\nu+1)}$ by the Erdős-Szekeres estimate [10]. Likewise, we have $S_2 \ll x^{1/\nu} h \log x$, establishing (7.21). $\square$

Note that $\sum_{x < n \leq x + x/\log^2 x} \mathbb{1}_{(f(n), q)=1} \ll x^{1/\nu} / \log^2 x$ (write $n = n_1 n_2$ and proceed as above). Hence, it suffices to show (7.20) with “ x ” replaced by the “ X ” in Lemma 7.9. Now, our Dirichlet series $\mathcal{F}(s) = \sum_{n \geq 1} \mathbb{1}_{(f(n), q)=1} \widehat{\psi}(\widehat{f}(n)) / n^s$ absolutely converges on $\text{Re}(s) > 1/\nu$, since in the Euler product $\mathcal{F}(s) = \prod_p (1 + \sum_{r \geq 1} \mathbb{1}_{(f(p^r), q)=1} / p^{rs})$, all but finitely many of the factors are of the form $1 + \sum_{r \geq \nu} \mathbb{1}_{(f(p^r), q)=1} / p^{rs}$ by ν -admissibility. Thus by Perron’s formula [35, Theorem II.2.3],

$$\sum_{n \leq X} \mathbb{1}_{(f(n), q)=1} \widehat{\psi}(\widehat{f}(n)) = \frac{1}{2\pi i} \int_{\frac{1}{\nu}(1+\frac{1}{\log X}) - iT}^{\frac{1}{\nu}(1+\frac{1}{\log X}) + iT} \frac{\mathcal{F}(s) X^s}{s} ds + O\left(\frac{X^{1/\nu} \log X}{T}\right).$$

To bound the error term from Perron, we used Lemma 7.9, and the technique in (5.2) which gave

$$\sum_{\substack{n \geq 1 \\ n \notin (3X/4, 5X/4)}} \frac{\mathbb{1}_{(f(n), q)=1}}{n^{\frac{1}{\nu}(1+\frac{1}{\log X})} |\log(X/n)|} \ll \sum_{\substack{n_2 \geq 1 \\ n_2 \text{ is } \nu\text{-full}}} \frac{1}{n_2^{\frac{1}{\nu}(1+\frac{1}{\log X})}} \ll \prod_p \left(1 + \sum_{r \geq \nu} \frac{1}{p^{\frac{r}{\nu}(1+\frac{1}{\log X})}} \right) \ll \log X.$$

The rest of the argument in section 5 goes through only by scaling things by ν appropriately. Branch cut conventions are analogous to those before the statement of Lemma 5.1: Branch cuts up to $\sigma \leq 1$ (respectively, $\sigma \leq \rho_e$) there are replaced by those up to $\sigma \leq 1/\nu$ (resp. $\sigma \leq \rho_e/\nu$). Redefining $\mathcal{L}_{q^*}(t) = \log(Q(|t\nu| + 1))$, Lemma 5.1 holds with $F_{\hat{\psi}}(s)$ replaced by

$$F_{\hat{\psi}}(s\nu) := \left(\prod_{\eta \bmod Q} L(s\nu, \eta)^{\gamma(\eta)} \right)^{\alpha_\nu(Q)c_{\hat{\psi}}} , \quad \text{with } \gamma(\eta) := \frac{1}{\alpha_\nu(Q)\varphi(Q)} \sum_{\substack{u \in U_{q^*} \\ \prod_{i=1}^K F_{i,\nu}(u) \in U_{q^*}}} \bar{\eta}(u),$$

so that $G_{\hat{\psi}}(s)$ (redefined accordingly) is analytic on $\sigma > \nu^{-1}(1 - 1/\log q^*)$ and satisfies $G_{\hat{\psi}}(s) \ll (\log_2 q^*)^3$ uniformly therein. (The analogue of (5.5) holds with $1 - c_{\hat{\psi}} \cdot \mathbb{1}_{p \prod_{i=1}^K F_{i,\nu}(p) \in U_{q^*}}/p^{s\nu}$ replacing $1 - c_{\hat{\psi}} \cdot \mathbb{1}_{pF(p) \in U_{q^*}}/p^s$ there.) We redefine our contour Γ_1 (Figure 1) by replacing $1 + 1/\log x$, 1 , $\tilde{\rho}_e, \rho_e$, by $\nu^{-1}(1 + 1/\log X)$, $1/\nu$, $\tilde{\rho}_e/\nu$, ρ_e/ν , and by taking $\vartheta(t) := \nu^{-1}(1 - c_1/4\mathcal{L}_{q^*}(t))$, $T = \exp(\sqrt{\log x})$.

We redefine $H_{\hat{\psi}}(s)$ to $H_{\hat{\psi}}(s) := F_{\hat{\psi}}(s\nu) (s - 1/\nu)^{\alpha_\nu(Q)c_{\hat{\psi}}} (s - \rho_e/\nu)^{-\alpha_\nu(Q)\gamma(\eta_e)c_{\hat{\psi}}}$ which analytically continues into $\sigma > \nu^{-1}(1 - c_1/4\mathcal{L}_{q^*}(t))$. The analogue of Proposition 5.3 holds with s in subpart (2) lying in $[\nu^{-1}(1 - c_1/4\log q^*), \nu^{-1}]$. Its proof is analogous, the main changes being:

- (5.7) holds with “ $1 - c_1/2\mathcal{L}_{q^*}(t) \leq \operatorname{Re}(s)$ ” replaced by “ $\nu^{-1}(1 - c_1/2\mathcal{L}_{q^*}(t)) \leq \operatorname{Re}(s)$ ”.
- In the proof of subpart (1), we redefine $\mu(t)$ to $\nu^{-1}(1 + c_1/7\mathcal{L}_{q^*}(t))$. Hence the mention of $\zeta(\mu(t))$ is replaced by $\zeta(\nu\mu(t))$.

Proposition 5.4 and everything thereafter goes through with analogous arguments; the analogue of Γ_0 in (5.14) is the union of segments joining $\tilde{\rho}_e/\nu$ and $1/\nu$ above and below the branch cut. At the very end (in the case when $c_{\hat{\psi}} = 1$), we can write $G_{\hat{\psi}}(s) = G_{\hat{\psi},1}(s)G_{\hat{\psi},2}(s)$, where $G_{\hat{\psi},2}(s) := \prod_{p \leq 2\nu} (1 + \sum_{r \geq 1} \mathbb{1}_{(f(p^r), q)=1}/p^{rs})$, and where $G_{\hat{\psi},1}(s)$ is analytic on $\sigma > \nu^{-1}(1 - 1/\log q^*)$ and satisfies $G_{\hat{\psi},1}(s) \ll (\log_2 q^*)^3$ uniformly therein. By definition of $\mathcal{Q}(\nu; f_1, \dots, f_K)$ (i.e., (7.10) and its ensuing observation), we have $G_{\hat{\psi},2}(1/\nu) = 0$, allowing us to adapt (5.17) and (5.18). This establishes the desired (7.20), completing the proofs of Theorems 7.5 and 7.6. $\square$

7.5. Optimality in Theorems 7.5 and 7.6. The arguments for Lemma 6.1 show that given $\hat{F} := (F_{1,\nu}, \dots, F_{K,\nu})$ with $\prod_{i=1}^K F_{i,\nu}$ squarefree, there exists a constant $C_{\hat{F}} > 0$ such that any q with $P^-(q) > C_{\hat{F}}$ has $\alpha_\nu(q) \neq 0$ and satisfies (7.10) vacuously (there is **no** tuple of characters $\hat{\chi} \neq (\chi_0, \dots, \chi_0) \bmod q$ for which $\prod_{i=1}^K \chi_i(F_{i,\nu}(u))$ is constant on $\{u \in U_q : \prod_{i=1}^K F_{i,\nu}(u) \in U_q\}$). Henceforth, we fix $\nu \in \mathbb{Z}^+$, a sufficiently large prime $\ell_0 > C_{\hat{F}}$, and $\{F_{i,j}\}_{1 \leq i \leq K, 1 \leq j \leq \nu-1} \subset \mathbb{Z}[T]$ with **all** coefficients divisible by ℓ_0 . Our $\{F_{i,\nu}\}_{i=1}^K$ play the role of the $\{F_i\}_{i=1}^K$ in section 6. Our moduli $q \leq (\log x)^{K_0}$ will have $P^-(q) = \ell_0$, so that by this discussion, $q \in \mathcal{Q}(\nu; f_1, \dots, f_K)$.¹⁰

The ranges (i) and (ii) in Theorem 7.5 are both optimal, and if $D_{\min} = 1$, the range in (i) is optimal for $\{F_{i,\nu}\}_{i=1}^K$ being **any** pairwise coprime linear polynomials. All this can be shown by using the set-up in the previous paragraph, and having the prime powers P^ν play the role of the prime inputs “ P ” in section 6. (Example: To prove optimality of Theorem 7.5(i), take $F_{i,\nu}(T) := (T - 1)^r + i$, and q a perfect r -th power with $P^-(q) = \ell_0 > \max\{C_{\hat{F}}, 2K\}$. Then any prime $P \equiv 1 \pmod{q^{1/r}}$ satisfies $f_i(P^\nu) \equiv i \pmod{q}$, showing that $\#\{n \leq x : (\forall i) f_i(n) \equiv i \pmod{q}\} \gg x^{1/\nu}/(q^{1/r} \log x)$.)

¹⁰The construction of $\{F_{i,j}\}_{1 \leq i \leq K, 1 \leq j \leq \nu-1}$ guarantees that $\alpha_j(q) = 0$ for all $j < \nu$, and the definition of $C_{\hat{F}}$ guarantees both that $\alpha_\nu(q) \neq 0$ and that q satisfies (7.10).

Finally, we show the optimality of the values of R in (7.12); hereafter q is **squarefree**. The value $R = 2$ is optimal by the constructions in section 6. To show that $R = \nu(K\nu + K - \nu) + 1$ is optimal in the setting of (7.12), we adapt the last construction in section 6 as follows:

Take $F_{i,\nu}(T) := \prod_{j=1}^r (T - 2j) + 2(2i - 1)$, and continue with the set-up in the first paragraph of this subsection (with $P^-(q) = \ell_0 > \max\{C_{\widehat{F}}, 4Kr\}$). Let $f_1, \dots, f_K : \mathbb{Z}^+ \rightarrow \mathbb{Z}$ be any multiplicative functions satisfying $f_i(p^j) = F_{i,j}(p)$ for all $i \in [K]$ and all $j \in [\nu]$, as well as $f_i(p^{\nu+1}) := 1$ for all i . Let $n = (p_1 \dots p_{(K-1)\nu})^{\nu+1} \cdot P^\nu$, with $p_1, \dots, p_{(K-1)\nu}, P$ being distinct primes satisfying $q < p_{K-1} < \dots < p_1 < x^{1/8K\nu^2} < x^{1/3\nu} < P \leq (x/(p_1 \dots p_{(K-1)\nu})^{\nu+1})^{1/\nu}$ and $\prod_{j=1}^r (P - 2j) \equiv 0 \pmod{q}$. (Recall that there are exactly $r^{\omega(q)}$ many possible coprime residue classes mod q that P could lie in.) Then $n \leq x$, and $f_i(n) = F_{i,\nu}(P) \equiv 2(2i - 1) \pmod{q}$ for all i , so that a computation entirely analogous to that done in (6.2) and (6.3) shows that

$$\sum_{\substack{n \leq x: P_{\nu(K\nu+K-\nu)}(n) > q \\ (\forall i) f_i(n) \equiv 2(2i-1) \pmod{q}}} 1 \gg \frac{r^{\omega(q)}}{\varphi(q)^K (\log_2 x)^{\nu(K-1)+1}} \cdot \frac{x^{1/\nu}}{\log x}$$

As in section 6, the right can be made to grow much faster than our expected main term.

ACKNOWLEDGEMENTS

I would like to thank Paul Pollack for fruitful discussions and for his continued support and encouragement. I am grateful to Martin Cech, Siu Hang Man, Neil Lyall, Akos Magyar and Giorgis Petridis, for helpful comments on the exposition. I am also grateful to the Department of Mathematics at the University of Georgia for their support and hospitality. I would like to thank William Banks, Régis de la Bretèche, Caroline Turnage-Butterbaugh, Karl Dilcher, Kevin Ford, Michael Filaseta, Larry Guth, Tsz Ho Chan, Jeffrey Lagarias, Robert Lemke-Oliver, Florian Luca, Yu-Ru Liu, Amita Malik, Carl Pomerance, Yuta Suzuki, Lola Thompson, Enrique Trevino, Ognian Trifonov, Lee Troupe, Asif Zaman and Tamar Ziegler for their helpful advice and/or their interest in my work. I acknowledge support from the Charles University Grant PRIMUS/25/SCI/008.

REFERENCES

1. A. Akande, *Uniform distribution of polynomially-defined additive functions to varying moduli*, submitted.
2. W. D. Banks and I. E. Shparlinski, *Congruences and exponential sums with the Euler function*, High primes and misde-meanours: lectures in honour of the 60th birthday of Hugh Cowie Williams, Fields Inst. Commun., vol. 41, Amer. Math. Soc., Providence, RI, 2004, pp. 49–59.
3. W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics, vol. 39, Cambridge University Press, Cambridge, 1998.
4. E. R. Canfield, P. Erdős, and C. Pomerance, *On a problem of Oppenheim concerning “factorisatio numerorum”*, J. Number Theory **17** (1983), 1–28.
5. T. Cochrane, *Exponential sums modulo prime powers*, Acta Arith. **101** (2002), 131–149.
6. H. Delange, *On integral-valued additive functions*, J. Number Theory **1** (1969), 419–430.
7. ———, *On integral-valued additive functions, II*, J. Number Theory **6** (1974), 161–170.
8. T. Dence and C. Pomerance, *Euler’s function in residue classes*, Ramanujan J. **2** (1998), 7–20.
9. Z. Dvir, J. Kollár, and S. Lovett, *Variety evasive sets*, Comput. Complexity **23** (2014).
10. P. Erdős and G. Szekeres, *Über die Anzahl der Abelschen Gruppen gegebener Ordnung und über ein verwandtes zahlen-theoretisches Problem*, Acta Univ. Szeged **7** (1934–1935), 95–102.
11. O. M. Fomenko, *The distribution of values of multiplicative functions with respect to a prime modulus (Russian)*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **93** (1980), 218–224.
12. A. Granville and K. Soundararajan, *Pretentious multiplicative functions and an inequality for the zeta-function*, Anatomy of integers, CRM Proc. Lecture Notes, vol. 46, Amer. Math. Soc., Providence, RI, 2008, pp. 191–197.
13. R. R. Hall and G. Tenenbaum, *Divisors*, Cambridge Tracts in Mathematics, vol. 90, Cambridge University Press, Cambridge, 1988.
14. S. Konyagin, *Letter to the editors: “The number of solutions of congruences of the n th degree with one unknown”*, Mat. Sb. (N.S.) **110(152)** (1979), 158.

15. ———, *The number of solutions of congruences of the n th degree with one unknown*, Mat. Sb. (N.S.) **109(151)** (1979), 171–187, 327.
16. E. Landau, *Lösung des Lehmer'schen Problems*, American J. Math. **31** (1909), 86–102.
17. N. Lebowitz-Lockard, P. Pollack, and A. Singha Roy, *Distribution mod p of Euler's totient and the sum of proper divisors*, Michigan Math. J. **74** (2024), 143–166.
18. D. B. Leep and C. C. Yeomans, *The number of points on a singular curve over a finite field*, Arch. Math. (Basel) **63** (1994), 420–426.
19. W. Narkiewicz, *On distribution of values of multiplicative functions in residue classes*, Acta Arith. **12** (1967), 269–279.
20. ———, *Euler's function and the sum of divisors*, J. Reine Angew. Math. **323** (1981), 200–212.
21. ———, *On a kind of uniform distribution for systems of multiplicative functions*, Litovsk. Mat. Sb. **22** (1982), no. 1, 127–137.
22. ———, *Distribution of coefficients of Eisenstein series in residue classes*, Acta Arith. **43** (1983), 83–92.
23. ———, *Uniform distribution of sequences of integers in residue classes*, Lecture Notes in Mathematics, vol. 1087, Springer-Verlag, Berlin, 1984.
24. W. Narkiewicz and F. Rayner, *Distribution of values of $\sigma_2(n)$ in residue classes*, Monatsh. Math. **94** (1982), 133–141.
25. P. Pollack and A. Singha Roy, *Joint distribution in residue classes of polynomial-like multiplicative functions*, Acta Arith. **202** (2022), 89–104.
26. ———, *Benford behavior and distribution in residue classes of large prime factors*, Canad. Math. Bull. **66** (2023), no. 2, 626–642.
27. ———, *Distribution in coprime residue classes of polynomially-defined multiplicative functions*, Math. Z. **303** (2023), no. 4, 20 pages.
28. F. Rayner, *Weak uniform distribution for divisor functions. I*, Math. Comp. **50** (1988), 335–342.
29. ———, *Weak uniform distribution for divisor functions. II*, Math. Comp. **51** (1988), 331–337.
30. E. J. Scourfield, *A uniform coprimality result for some arithmetic functions*, J. Number Theory **20** (1985), 315–353.
31. A. Singha Roy, *Anatomical mean value bounds on multiplicative functions and the distribution of the sum of divisors*, submitted.
32. ———, *Joint distribution in residue classes of families of polynomially-defined additive functions*, submitted.
33. ———, *Joint distribution in residue classes of families of polynomially-defined multiplicative functions II*, submitted.
34. J. Śliwa, *On distribution of values of $\sigma(n)$ in residue classes*, Colloq. Math. **27** (1973), 283–291, 332.
35. G. Tenenbaum, *Introduction to analytic and probabilistic number theory*, third ed., Graduate Studies in Mathematics, vol. 163, American Mathematical Society, Providence, RI, 2015.
36. D. Wan, *Generators and irreducible polynomials over finite fields*, Math. Comp. **66** (1997), no. 219, 1195–1212.

CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF ALGEBRA, SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

Email address: akash01s.roy@gmail.com