
a
Median calibrated ρ
PRISM
GDSC
CTRP
Baseline RNA
Protein, MS (CCLE)
Protein, MS (ProCan)
Protein, RPPA
Tahoe response
Response + RNA
+0.04
+0.00
+0.22*
+0.13
+0.07
+0.18*
+0.05
+0.29*
+0.28*
+0.10
+0.17*
+0.18*
-0.05
+0.10
+0.10*
+0.02
+0.08
+0.23*
calibrated ρ
* FDR < 0.05 (Wilcoxon > 0)
b
Independent replication in CTRP (46 drugs)
+0.22*
+0.18*
+0.18*
+0.10*
+0.23*
+0.11*
Calibrated ρ per drug
Response < RNA: FDR 0.040
RPPA < RNA: FDR 0.0035
RNA
Protein (MS)
RPPA
Tahoe response
Response + RNA
Response + MS
c
Paired differences vs baseline RNA
Tahoe response − RNA
RPPA − RNA
MS (CCLE) − RNA
MS (ProCan) − RNA
Response + RNA − RNA
Median paired difference in calibrated ρ
PRISM
GDSC
CTRP
FDR < 0.05
d
Robust to model choice
Median paired difference
Ridge LOO (frozen)
Ridge-GCV
Kernel ridge
Elastic net
Random forest
Response adds to RNA
RPPA − RNA
PRISM
GDSC
e
Tahoe response: top vs middle dose
Calibrated ρ, middle dose
Calibrated ρ, top dose
PRISM: r = 0.12
GDSC: r = 0.08