
a
PRISM: median calibrated ρ by model family
RNA +0.04 +0.01 +0.06 -0.03 +0.11
Protein (MS) +0.13 +0.13 +0.11 +0.01 +0.09
RPPA +0.10 -0.03 +0.04 -0.04 -0.01
Tahoe response -0.05 +0.03 +0.07 -0.01 +0.04
Response + RNA +0.02 +0.04 +0.06 +0.02 +0.08
Response + MS +0.09 +0.06 +0.08 +0.03 +0.11
Ridge LOO (frozen)
Ridge-GCV
Kernel ridge
Elastic net
Random forest
b
GDSC: median calibrated ρ by model family
RNA +0.00 +0.07* +0.10 +0.11* +0.12*
Protein (MS) +0.07 +0.06 +0.01 -0.02 +0.05
RPPA +0.17* +0.09 +0.18 +0.06 +0.16
Tahoe response +0.10 +0.08 +0.01 +0.09 +0.07
Response + RNA +0.08 +0.14* +0.13* +0.15* +0.17*
Response + MS +0.11 +0.09 +0.08 +0.02 +0.17*
Ridge LOO (frozen)
Ridge-GCV
Kernel ridge
Elastic net
Random forest
median calibrated ρ
0.20
0.15
0.10
0.05
0.00
−0.05
−0.10
−0.15
−0.20
* FDR < 0.05 (Wilcoxon > 0, BH within model and screen). Frozen = Amendment 1 ridge LOO, PCA-5, α = 1, 500 permutations; best effort = 5-fold CV within drug, PCA-10, 50 permutations.
c
RPPA-target concordance (46 units)
Concordance (mean percentile of target-matched antibodies)
1.0
0.8
0.6
0.4
0.2
0.0
tucatinib
alpelisib
erlotinib
gefitinib
afatinib
osimertinib
PRISM (n = 25)
GDSC (n = 21)
d
Pooled median vs permutation null
observed 0.511
chance 0.5
one-sided p = 0.147
(not supported)
Permutations
800
600
400
200
0
0.44
0.46
0.48
0.50
0.52
0.54
0.56
Median concordance