

International Journal of Computing and Optimization
Vol. 5, 2018, no. 1, 33 - 38
HIKARI Ltd, www.m-hikari.com
<https://doi.org/10.12988/ijco.2018.846>

Some New Fuzzy Entropy Formulae

Geoffrey O. Barini

Institute of Basic Sciences, Technology and Innovation
Pan African University, P.O. Box 62000-00200, Nairobi-Kenya

Livingstone M. Ngoo

Department of Electrical and Telecommunication Engineering
Multimedia University of Kenya, P.O. Box 15653-00503, Nairobi-Kenya

Ronald M. Waweru

Department of Computing
Jomo Kenyatta University of Agriculture and Technology
P.O. Box 62000-00200, Nairobi-Kenya

Copyright © 2018 Geoffrey O. Barini, Livingstone M. Ngoo and Ronald M. Waweru.
This article is distributed under the Creative Commons Attribution License, which permits
unrestricted use, distribution, and reproduction in any medium, provided the original work
is properly cited.

Abstract

Fuzzy entropy plays a significant role in characterization and quantification of fuzzy uncertainty. This study introduces some new formulae of fuzzy entropy. Unlike most of the existing geometrical measures of fuzziness, the measures proposed here are based on the distance between a fuzzy set and the centre of a fuzzy unit hypercube. Their structure is particularly simple and intuitive.

Keywords: Fuzzy entropy, Distance Measure, Fuzzy hypercube

1 Introduction

Since its inception [11], the theory of fuzzy sets has continued to play a leading role in modeling non-statistical uncertainty-more specifically, its contribution

to characterization and quantification of fuzzy uncertainty is undeniable [12, 9]. Fuzzy sets provide a mathematical framework within which vague concepts can be quantitatively defined and analyzed more naturally without much loss of information.

Fuzziness is measured using fuzzy entropy [3]. Motivated by the classical Shannon entropy, Deluca and Termin introduced the now widely adopted definition of fuzzy entropy [3]. Since then, numerous researchers have successfully formulated new fuzzy entropy formulae satisfying this definition ([7],[8],[6],[10]). This study proposes some new distance based fuzzy entropy formulae. The structure of the formulae is based the distance between a fuzzy set and the centre of the a fuzzy unit hypercube.

The following section provides definitions and notations. The third section reviews some existing measures of fuzzy entropy. Then some new formulae of fuzzy entropy are given in the last section.

2 Preliminaries

Let $\zeta = \{x_1, x_2, \dots, x_n\}$ be a universal set. A set of all crisp and fuzzy set over ζ is denoted by 2^ζ and $F(2^\zeta)$ respectively. A fuzzy subset $A = \{(x_1, a_1), (x_2, a_2), \dots, (x_n, a_n)\}$ is characterized by its membership function m_A , a mapping

$$m_A : \zeta \rightarrow [0, 1] \quad (1)$$

such that $m_A(x_i) = a_i \in [0, 1]$ is a membership (or fit) value of x_i in A . Fuzzy set A represented by a fit vector (a_1, a_2, \dots, a_n) is regarded as a point in an n -dimensional fuzzy unit hypercube $[0, 1] \times [0, 1] \times \dots \times [0, 1] = I^n$ [8]. A constant $A_{[a]}$ fuzzy set on ζ is such that $m_A(x_i) = a \forall x_i \in \zeta$. The complement \bar{A} of A is fuzzy set with membership function $m_{\bar{A}}(x) = 1 - m_A(x)$. The cardinality of A is a non-negative real scalar $c(A) = \sum_{i=1}^n m_A(x_i)$. A fuzzy set A^* is said to be a sharpened form of A if $m_A(x) \geq \frac{1}{2}$ then $m_{A^*}(x) \geq m_A(x)$ and if $m_A(x) \leq \frac{1}{2}$ then $m_{A^*}(x) \leq m_A(x)$.

For any $A \in F(2^\zeta)$, denote respectively by A_{near} and A_{far} the nearest and farthest crisp sets with membership functions

$$A_{near} = \begin{cases} 1, & \text{if } m_A(x) \geq \frac{1}{2} \\ 0, & \text{if } m_A(x) < \frac{1}{2} \end{cases} \quad (2)$$

and

$$A_{far} = \begin{cases} 0, & \text{if } m_A(x) \geq \frac{1}{2} \\ 1, & \text{if } m_A(x) < \frac{1}{2} \end{cases} \quad (3)$$

A general class of distance measure, namely the Minkowski distance between fuzzy sets A and B is defined as,

$$\gamma_p(A, B) = \left(\sum_{i=1}^n |\mu_A(x_i) - \mu_B(x_i)|^p \right)^{\frac{1}{p}}, \quad p \geq 1 \quad (4)$$

3 Fuzzy Entropy

Fuzzy entropy is a set to point mapping [3]

$$E : F(2^\zeta) \rightarrow [0, 1] \quad (5)$$

given by

$$E(A) = -k \{m_A(x_i) \log m_A(x_i) + (1 - m_A(x_i)) \log (1 - m_A(x_i))\} \quad (6)$$

with $k \geq 0$, satisfying the following properties.

$$(A1) \quad E(A) = 0 \forall A \in 2^\zeta$$

$$(A2) \quad E(A) \text{ attains its maximum at } A_{[\frac{1}{2}]}$$

$$(A3) \quad \text{if } A^* \text{ is a sharpened version of any } A \in F(2^\zeta) \text{ then } E(A^*) \leq E(A)$$

$$(A4) \quad \text{For any } A \in F(2^\zeta), E(A) = E(\overline{A})$$

Based on this axiomatic definition, Kaufman [6] proposed a measure of fuzziness based the distance between fuzzy set A and the nearest crisp set A_{near} as

$$E_{ka}(A) = \left(\sum_{i=1}^n |m_A(x_i) - m_{A_{near}}(x_i)|^p \right)^{\frac{1}{p}} \quad (7)$$

For $p = 1$ and $p = 2$ this measure collapses to special cases namely,

$$E_{ka}(A) = \sum_{i=1}^n |m_A(x_i) - m_{A_{near}}(x_i)| \quad (8)$$

and

$$E_{ka}(A) = \left(\sum_{i=1}^n |m_A(x_i) - m_{A_{near}}(x_i)|^2 \right)^{\frac{1}{2}} \quad (9)$$

the Hamming and Euclidean distances respectively. Viewing fuzziness as indistinguishability between fuzzy set A and its opposite \overline{A} , Yager suggested a measure of fuzziness given by the formula[10],

$$E_y(A) = 1 - \frac{\gamma_p(A, \overline{A})}{n^{\frac{1}{p}}} \quad (10)$$

where

$$\gamma_p(A, \bar{A}) = \left(\sum_{i=1}^n |m_A(x_i) - m_{\bar{A}}(x_i)|^p \right)^{\frac{1}{p}} \quad (11)$$

For $p = 1$, this measure is of particular simplicity,

$$E_y(A) = 1 - \hat{\gamma}_1(A, \bar{A}) \quad (12)$$

where $\hat{\gamma}_1(A, \bar{A})$ is the normalized Hamming distance.

Kosko's fuzzy entropy formula[8],

$$E_{k1}(A) = \frac{\gamma_p(A, A_{near})}{\gamma_p(A, A_{far})} \quad (13)$$

was largely motivated by the geometrical representation of fuzzy sets. Kosko formulated another fuzzy entropy measure based on the fuzzy operators of the union and intersection as[8]

$$E_{k2}(A) = \frac{c(A \cap \bar{A})}{c(A \cup \bar{A})} \quad (14)$$

The formula given in is sometimes known as the fuzzy entropy theorem [8].

Distance induced fuzzy entropy is also introduced in [5]. Bhandari and Pal [1] have suggested fuzzy entropy of a fuzzy set with respect to some fixed fuzzy set.

4 Some new fuzzy entropy formulae

For any $A \in F(2^\zeta)$ it is directly that,

$$\begin{aligned} \gamma_p\left(A, A_{\left[\frac{1}{2}\right]}\right) &= \gamma_p\left(\bar{A}, A_{\left[\frac{1}{2}\right]}\right) = \left(\sum_{i=1}^n |m_A(x_i) - \frac{1}{2}|^p \right)^{\frac{1}{p}} \\ &\leq \frac{n^{\frac{1}{p}}}{2} \end{aligned} \quad (15)$$

The metric $\gamma_p\left(A, A_{\left[\frac{1}{2}\right]}\right)$ inversely expresses the degree of fuzziness of A . Thus,

$$E_1(A) = 1 - \frac{n^{\frac{1}{p}}}{2} \gamma_p\left(A, A_{\left[\frac{1}{2}\right]}\right) \quad (16)$$

gives a measure of fuzzy degree of set A . We can also intuitively introduce a measure of entropy of fuzzy set A as,

$$E_2(A) = \frac{n^{\frac{1}{p}} - 2\gamma_p\left(A, A_{\left[\frac{1}{2}\right]}\right)}{n^{\frac{1}{p}} + 2\gamma_p\left(A, A_{\left[\frac{1}{2}\right]}\right)} \quad (17)$$

is a fuzzy entropy measure.

Recognize that this measure can be expressed as,

$$E_2(A) = \frac{n^{\frac{1}{p}} - 2 \left(\sum_{i=1}^n |m_A(x_i) - \frac{1}{2}|^p \right)^{\frac{1}{p}}}{n^{\frac{1}{p}} + 2 \left(\sum_{i=1}^n |m_A(x_i) - \frac{1}{2}|^p \right)^{\frac{1}{p}}} \quad (18)$$

Similarly (A1), (A2) and (A4) are easy to check. The proof for (A3) follows from the fact that fuzziness of A is inversely related to the distance between A and the centre of the hypercube.

5 Conclusion

We have intuitively introduced new fuzzy entropy measures based on the distance between a fuzzy set and the centre of the fuzzy unit hypercube. These measures offer more choices from which we can choose in modeling uncertainty due to fuzziness

Acknowledgements. This study was supported by PAUISTI, Under the Commission of the African Union.

References

- [1] D. Bhandari, N. R. Pal, Some new information measures for fuzzy sets, *Information Sciences*, **67** (1993), no. 3, 209-228.
[https://doi.org/10.1016/0020-0255\(93\)90073-u](https://doi.org/10.1016/0020-0255(93)90073-u)
- [2] V. Carr, J. Tah, A fuzzy approach to construction project risk assessment and analysis: construction project risk management system, *Advances in Engineering Software*, **32** (2001), no. 10, 847-857.
[https://doi.org/10.1016/s0965-9978\(01\)00036-9](https://doi.org/10.1016/s0965-9978(01)00036-9)
- [3] A. De Luca, S. Termini, A definition of a non probabilistic entropy in the setting of fuzzy set theory, *Information and Control*, **20** (1972), no. 4, 301-312. [https://doi.org/10.1016/s0019-9958\(72\)90199-4](https://doi.org/10.1016/s0019-9958(72)90199-4)
- [4] D. Dubois, H. Prade, *Fundamentals of Fuzzy Sets*, Vol. 20, Springer Science and Business, 2012.
- [5] J. Fan, W. Xie, Distance measure and induced fuzzy entropy, *Fuzzy Sets and Systems*, **104** (1999), no. 1, 305-314.
[https://doi.org/10.1016/s0165-0114\(99\)80011-6](https://doi.org/10.1016/s0165-0114(99)80011-6)

- [6] A. Kaufman, Introduction to the theory of fuzzy subsets, *Fuzzy Sets and Systems*, **1** (1986).
- [7] B. Kosko, Fuzziness entropy and conditioning, *Information Sciences*, **40** (1986), no. 2, 165-174. [https://doi.org/10.1016/0020-0255\(86\)90006-x](https://doi.org/10.1016/0020-0255(86)90006-x)
- [8] B. Kosko, Fuzziness vs. probability, *International Journal of General Systems*, **17** (1990), no. 2-3, 211-240.
<https://doi.org/10.1080/03081079008935108>
- [9] N. R. Pal, S. K. Pal, Higher order fuzzy entropy and hybrid entropy of a set, *Information Sciences*, **61** (1992), no. 3, 211-231.
[https://doi.org/10.1016/0020-0255\(92\)90051-9](https://doi.org/10.1016/0020-0255(92)90051-9)
- [10] R. R. Yager, On the measure of fuzziness and negation part I: membership in the unit interval, *International Journal of General Systems*, **5** (1979), no. 4, 221-229. <https://doi.org/10.1080/03081077908547452>
- [11] L. A. Zadeh, Fuzzy sets, *Information and Control*, **8** (1965), no. 3, 338-353. [https://doi.org/10.1016/s0019-9958\(65\)90241-x](https://doi.org/10.1016/s0019-9958(65)90241-x)
- [12] L. A. Zadeh, The role of fuzzy sets in the management of uncertainty in expert systems, *Fuzzy Sets and Systems*, **11** (1983), no. 1-3, 199-227.
[https://doi.org/10.1016/s0165-0114\(83\)80081-5](https://doi.org/10.1016/s0165-0114(83)80081-5)

Received: May 11, 2018; Published: October 3, 2018