

Sanjivani University
School of Engineering and Technology

Applied Science (24UETBS102)
Unit 1
Introduction to Fundamental Physics

Content:

Waves and Oscillation: Types of Waves, Superposition of Waves, Standing Waves (Nodes and Antinodes), , and Its Applications.

SHM: Wave Equations, Simple Pendulum (Problems).

Electrodynamics: Faraday's Laws, Lenz's Law, Biot-Savart Law, Physics of Divergence, Gradient, and Curl, Magnetic Field, Vector Potential, Maxwell Equations (Qualitative).

Semiconductor Electronics: Band theory of solids, Energy band diagram, Semiconductor: Properties, Type, P-N Junction diode and applications, I-V characteristics, P-N-P and N-P-N transistors,

Modern Physics: Photoelectric Effect, Einstein's Photoelectric equation, Photocell(working), Solar cell.

Ancient Astronomical Instruments: Chakra, Dhanuryantra, Yasti and Phalak Yantra etc.

1. Waves and Oscillation

Introduction:

Wave is nothing but an oscillatory disturbance produced in the medium.

A means by which disturbances are transmitted from one region to another is called as **wave motion**.

Example

Sound spread in a hall

Stone dropped in a pond having still surface.

In wave motion only energy is transferred however no part of medium is transported in the medium.

Waves can be classified into different types based on their characteristics and the medium through which they travel.

Mechanical Waves: Mechanical waves require a medium (such as air, water, or solids) to travel through.

Example sound waves

Electromagnetic waves: Waves which do not require material medium for their propagation

Example: Light Waves

Progressive Wave: The wave which travel continuously in a given direction without change of form is called as progressive wave.

There are two types of progressive waves

Sr. No	Transverse wave	Longitudinal wave
1	A wave in which particles of the medium vibrate in a direction perpendicular to the direction of propagation of wave.	A wave in which particles of the medium vibrate in a direction parallel to the direction of propagation of wave.
2	Light wave	Sound wave
3	Do not require medium	Requires medium Medium should possess property of elasticity of volume i.e. Bulk modulus

Surface Waves: These waves travel along the surface of a medium and have characteristics of both transverse and longitudinal waves.

Example

Water waves.

Recall concepts

Amplitude: Maximum displacement from mean position.

SI unit m

Frequency: Number of oscillations performed per second.

SI unit Hz

Time Period: Time required to complete one oscillation

SI Unit second

$$T = 1/n$$

Wavelength: Distance covered by wave in one complete oscillation. SI Unit m

Or

Distance between two successive compression or rarefactions. (in longitudinal wave)

Or

Distance between two successive crests and troughs (In transverse wave)

Velocity of wave: Distance traveled by wave in one second.

SI unit m/s

Relation between velocity wavelength and frequency

λ

$$V = n \lambda$$

=wavelength

V= velocity

n= frequency

Electromagnetic Waves: Waves which do not require a medium .

They are created by the oscillation of electric and magnetic fields and include:

Light waves: Travel with a velocity of $3 \times 10^8 \text{ m/s}$

Radio Waves: Used in communication systems, such as radio and television broadcasts.

Microwaves: Used in microwave ovens and radar technology.

Infrared Waves: Experienced as heat and used in remote controls and thermal imaging.

Visible Light: The portion of the electromagnetic spectrum visible to the human eye.

Ultraviolet Light: Beyond the visible spectrum, used in sterilization and fluorescent lighting.

X-Rays: Used in medical imaging and security scanners.

Gamma Rays: Emitted by radioactive materials and used in cancer treatment and imaging.

Superposition of waves:

Principle:

When two waves propagate simultaneously in a medium the displacement of the particles in the medium at any instant of time is the algebraic sum of the displacement caused by the individual waves.

Example :

If two musical instruments are played simultaneously on stage like sitar and tabla , the audience can clearly distinguish the sound of these instruments.

Principle of Superposition can be used to study

Interference


Beats

Stationary waves

Interference:

When two sound waves of same amplitude and frequency travel simultaneously in a medium, the resulting wave has an intensity different from that of the individual waves. It is louder at some place whereas there is complete silence at some place. This is called as interference.

<https://www.acs.psu.edu/drussell/Demos/superposition/superposition.html>




Interference of waves

Beats:

When two sound waves of same amplitude but slightly different frequencies travel in a medium along the same direction, the resultant amplitude is alternately maximum and minimum. They produce resultant sound waves of varying intensities (waxing and waning).

The phenomenon of waxing and waning of sound is called as beats.




Formation of Beats (<https://byjus.com/physics/beats/>)

Stationary Waves:

When the two identical progressive waves travel through a medium along the same line in opposite directions with the equal velocities, they superimpose over each other and produce a new type of wave called as stationary waves or standing waves along which there is no transfer of energy.

Nodes : The points half a wavelength apart in a medium which are permanently at rest are called as *Nodes*.

Antinodes: The points in medium midway between the nodes, where the displacement is maximum is called as *Antinodes*.



Stationary waves

Periodic motion:

Motion which repeats itself in equal intervals of time is called as periodic motion.

Oscillations: To and fro motion along the same path is called as oscillatory motion

Example

Motion of hands of clock

Motion of simple pendulum

Motion of hands of a man walking

Simple Harmonic Motion:

A motion in which force (acceleration) is always directed towards mean position and magnitude is directly proportional to displacement from mean position.

e.g. Motion of needle of sewing machine

Vibration of string

Motion of prongs of tuning fork. etc

Let us consider a particle moving along a circle of radius a with a constant speed v

Wave Equations:

Sr. No	Particular		Mean Position	Extreme Position
1	Equation of displacement	$y = a \sin(\omega t + \alpha)$	$t = 0, \alpha = 0$ $y = 0$ Minimum	$t = 0, \alpha = \frac{\pi}{2} \text{ or } \frac{3\pi}{2}$ $y = \pm a$ Maximum
2	Equation of Velocity	$v = \frac{dy}{dt} = \frac{d}{dt}(a \sin(\omega t + \alpha)) =$ $v = a\omega \cos(\omega t + \alpha)$ $v = \omega \sqrt{a^2 - y^2}$	$t = 0, \alpha = 0$ $v = a\omega$ maximum	$t = 0, \alpha = \frac{\pi}{2} \text{ or } \frac{3\pi}{2}$ $v = 0$ minimum
3.	Equation of acceleration	$acceleration = \frac{dv}{dt} = a\omega \cos(\omega t + \alpha)$ $acceleration = -a\omega^2 \sin(\omega t + \alpha)$ $acceleration = -\omega^2 y$	$t = 0, \alpha = 0$ Acceleration=0 Minimum	$t = 0, \alpha = \frac{\pi}{2} \text{ or } \frac{3\pi}{2}$ $acceleration = \pm a\omega^2$ Maximum

Where,

$y = \text{displacement}$

$v = \text{velocity}$

$a = \text{amplitude}$

$\alpha = \text{epoch}$


$\omega = \text{angular velocity}$

Simple Pendulum:

Simple pendulum is the simplest form of oscillatory motion.

Ideal simple pendulum is a heavy particle suspended by a massless, inextensible flexible string from a rigid support.

Practical simple pendulum is a small heavy sphere suspended by a light and inextensible string from a rigid support.



Period for simple pendulum is...

$$T = 2\pi\sqrt{\frac{L}{g}}$$

Simple pendulum performs linear simple harmonic motion for small amplitude.

Semiconductor Electronics:

Introduction:

Semiconductors are an essential component of electronic devices, enabling advances in communications, computing, healthcare, military systems, transportation, clean energy, and countless other applications. Due to their role in the fabrication of electronic devices, semiconductors are an important part of our lives. Imagine life without electronic devices. There would be no smartphones, radios, TVs, computers, video games, or advanced medical diagnostic equipment.

Band Theory of solids

➤ **Introduction** :- Atom is the fundamental unit of matter. It consists of a centrally placed nucleus which consists of protons and neutrons in it and negatively charged electrons are revolving around the nucleus in an elliptical orbit. Electrons in the inner shell (orbit) are bound to the nucleus and hence called bound electrons. The electrons in the outermost orbit are called valence electrons.

➤ **Formation of Energy bands in solids:**

There are millions of electrons belonging to the first orbit of atoms in the solid having different energy levels which are closely packed. Thus closely spaced energy levels of millions of first orbit electrons form a continuous band of energy and second orbit electrons form the second band.




Fig. 1 Energy level diagram of Atom




Fig. 2 Energy band diagram of solid

Q. Define energy band, valence band, conduction band, forbidden energy gap?

Energy band: The range of energies possessed by electrons of different atoms in solid is called as energy band.

Valence band: The range of energies possessed by electrons in the outermost orbit of an atom is called as valence band.

Conduction band: Valence electrons are loosely attached to the nucleus and can be detached and become free electrons. These free electrons in a conductor are responsible to carry current. They are called as conducting electrons.

“The range of energies possessed by conducting electrons is known as conduction band.”

Forbidden energy Gap: The band (energy gap) between valence band and conduction band is called forbidden gap.

In order to lift the electrons from valence band to conduction band .Band gap energy (E_g) must be provided.

Energy of electron in an atom: In an atom various electrons are distributed in different orbits. In an orbit certain permissible values of energy apart from these energy levels.

Draw the energy band diagram for conductors, semiconductors and insulators.

Ans. **Conductor**: Metal (aluminium, silver, copper) contains large number of free electrons at room temperature. Hence conduction band and valence bands are overlapping each other. Hence forbidden energy gap is absent.

Semiconductors: Conductivity of semiconductor is greater than insulator and less than conductors.

Valence band is full and conduction band is almost empty, forbidden gap is very small of the order of 1eV.




eg. Si and germanium.

Insulators: The forbidden energy gap is very high greater than 5 eV .The electrons cannot jump from valence band to conduction band.

Eg Glass, wood, rubber, plastic,etc.

Name two semiconductor which are commonly used.

Ans: **Silicon** and **germanium** are commonly used semiconductors.




Define intrinsic semiconductor.

Ans: The semiconductor material in its purest form is called as intrinsic semiconductor.

(at 0 K)


Explanation: At intrinsic semiconductor like pure silicon or germanium has four electrons in its outermost orbit of its atom. In order to fill valence shell atoms arrange themselves in such a way that each atom is surrounded by four atoms. This pattern is called as crystal. Each atom forms covalent bond with its four adjacent atom.



Intrinsic semiconductor At 0^0K

At absolute zero temperature there are no free electrons available for electrical conduction. **Semiconductor behaves as a perfect insulator at absolute zero temperature.**

At room temperature:-



Intrinsic semiconductor at room temperature

Room temperature is sufficient to break few covalent bonds. Thus few electrons are free to move in the crystal. When an electron moves away from an atom, a vacancy is created, i.e., a hole in the bond, which has a positive charge. Hence free electrons and holes are generated simultaneously (in pairs). This is called as thermal generation.

What is the effect of temperature on conductivity of semiconductor?

Ans: As the temperature of semiconductor increases, its conductivity increases.

Define Extrinsic semiconductor.

Ans: When a suitable impurity is added to a pure semiconductor, the type of semiconductor is called as an extrinsic semiconductor.


What is doping?

Ans: The process of adding impurity atoms in an intrinsic semiconductor to increase its conductivity is known as doping.

➤ **p type extrinsic semiconductor:**

Ans: When a pure semiconductor is doped with the trivalent impurity it is called as P type semiconductor. (gallium, indium, aluminium).

Consider a pure silicon crystal. If trivalent impurity like Al, Ga is added to it then three valence electrons from each Si atom forms three covalent bands with Ga atom and instead of fourth bond electron a hole is created.



p-type semiconductor


The **majority** carriers in p type are **holes** and impurity is called as **acceptor** impurity.

➤ **N type Extrinsic semiconductor**

When semiconductor is doped with pentavalent impurity it is called as n type semiconductor. (arsenic, antimony, phosphor) .

Consider a pure Si crystal If it is doped with pentavalent impurity like Arsenic (As) Antimony etc. which have five electrons in the outermost orbit, arsenic (impurity) atom forms four covalent bonds with Si atoms and fifth electron is extra and free in crystal.


The **majority** carriers in N type semiconductor are **electrons** and impurity is called as **donor** impurity.



n- type semiconductor

PN junction Diode

When P Type semiconductor and N type semiconductor are joined together the structure so formed is called as **PN junction diode** . (Diode is a device which has two terminals)




In P type semiconductor majority charge carriers are holes and in N type majority carriers are electrons.

When PN junction is formed, some electrons from N region cross the junction and diffuse into the P region. Recombination of holes and electrons takes place at the junction region creating +ve ions on N region and -ve ions on P region. After some time the process of recombination stops because of the formation of +ve and -ve ions on both sides of junction and that region is called as depletion region which is depleted of electrons and holes.

Due to the formation of +ve and -ve ions of PN junction an electric field V_B is produced at depletion layer called as **barrier potential** which acts as a small battery and opposes further diffusion of holes and electrons.(positive towards n-type and negative towards p-type)

Barrier potential for **silicon** is **0.7** volts and for **germanium** is **0.3volt** .




P-N junction diode

Biasing:--

Application of external DC supply to PN junction diode is called as biasing. There are two types of biasing on PN junction diode .

Forward biased PN junction Diode:

If P region is connected to positive terminal of battery and N region is connected to negative terminal of battery the diode is said to be forward biased.



Forward Biased PN junction Diode

Working:-

Holes in the P region get repelled by +ve of battery and cross the junction and electrons in N region get repelled by - ve of battery and crosses the junction because of which diode conducts current.

The barrier potential V_B opposes the external battery. In order to flow current through diode, the external voltage supplied must be greater than V_B i.e. 0.7 V for Si and 0.3 V for Ge. Because of pushing of charge carriers towards the junction, width of depletion layer decreases.