

Patient ID	10132070	Patient Name	Hanady Serage Hassan Eisa
Age / Gender	40Y 5M / Female	Referring Physician	Dr^Alshimaa^Rezk
DOB	1986-01-01	Procedure	ULTRASOUND THYROID
Study Time	2026-06-20 16:40:35	Reported DateTime	2026-06-21 16:36:04

US Neck:

Both thyroid glandlobes are seen within average size however showing diffuse heterogeneous parenchymal echotexture. On applying colour Doppler a rather hypervascularity was noted.

Right lobe measures 3.5 x 1 x 2.1 cm in its longitudinal, transverse, and AP diameter, with estimated volume of 3.9 ml. It shows an ill-defined wider than taller nodule with indeterminate composition, indeterminate echogenicity due to overlying coarse calcific foci, approximately measuring 4.3 x 2.8 mm, TI-RAD 4

Left lobe measures 4.4 x 1.6 x 1.8 cm with estimated volume of 6.4 ml. It shows a very ill-defined calcified nodule similar to that seen on the right lobe, TI-RAD 4 approximately measuring 4.7 x 2.6 mm

Another ill-defined wider than taller isoechoic lesion is noted at its posterior margin approximately measuring 13.6 x 9.8 mm showing smooth margins with no calcific foci likely representing a pseudo nodule


Isthmus measures 8 mm

Normal appearance of both submandibular glands with few intraparenchymal lymph nodes. Bilateral cervical likely reactionary enlarged lymph nodes were noted. No other remarkable abnormality.

Conclusion:

Ultrasound picture impressive of thyroiditis, to be correlated clinically and with laboratory results

Bilateral thyroid nodules, TI-RAD 4 as detailed to be correlated with findings of previously done biopsy



Dr. Fatmaalzhaa Mahmoud Ahmed
Specialist Radiologist