

Assignment 12

Akhila Kumbha,CS21BTECH11031

June 16, 2022

Outline

1 Question

2 Solution

Question

Let $P_n(z)$ represent the Levinson polynomial of the first kind.

(a) If one of the roots of $P_n(z)$ lie on the unit circle, then show that all other roots of $P_n(z)$ are simple and lie on the unit circle.

(b) If the reflection coefficient $s_k \neq 0$, then show that $P_k(z)$ and $P_{k+1}(z)$ have no roots in common.

Solution

(a) Let $z = e^{j\theta_1}$ represent one of the roots of the Levinson Polynomial $P_n(z)$ that lie on the unit circle. In that case

$$P_n(e^{j\theta_1}) = 0 \quad (1)$$

and substituting this into the recursion equation..

$$\sqrt{1 - s_n^2} P_n(z) = P_{n-1}(z) - z s_n \tilde{P}_{n-1}(z) \quad (2)$$

We get

$$|s_n| = \left| \frac{P_{n-1}(e^{j\theta_1})}{\tilde{P}_{n-1}(e^{j\theta_1})} \right| = 1 \quad (3)$$

so that

$$s_n = e^{j\alpha} \quad (4)$$

Let

$$P_{n-1}(e^{j\theta}) = R(\theta)e^{j\psi(\theta)} \quad (5)$$

and since $P_{n-1}(z)$ is free of zeros in $|z| \leq 1$, we have $R(\theta) > 0$, $0 < \theta < 2\pi$, and once again substituting these into

$$\sqrt{1 - s_n^2} P_n(z) = P_{n-1}(z) - z s_n \tilde{P}_{n-1}(z) \quad (6)$$

We obtain

$$\sqrt{1 - s_n^2} P_n(e^{j\theta}) = R(\theta)e^{j\psi(\theta)} - e^{j(\theta+\alpha)} e^{j(n-1)\theta} R(\theta)e^{-j\psi(\theta)} \quad (7)$$

$$= R(\theta)[e^{j\psi(\theta)} - e^{j(n\theta+\alpha)} e^{-j\psi(\theta)}] \quad (8)$$

$$= 2jR(\theta)e^{j(n\theta+\alpha)/2} \sin\left(\psi(\theta) - \frac{n\theta}{2} - \frac{\alpha}{2}\right) \quad (9)$$

Due to the strict Hurwitz nature of $P_{n-1}(z)$, as θ varies from 0 to 2π , there is no net increment in the phase term $\psi(\theta)$, and the entire argument of the sine term above increases by $n\pi$.

Consequently $P_n(e^{j\theta})$ equals zero atleast at n distinct points

$\theta_1, \theta_2, \dots, \theta_n$, $0 < \theta_i < 2\pi$.

However $P_n(z)$ is a polynomial odd degree n in z and can have atmost n zeros. Thus all the above zeros are simple and they all lie on the unit circle.

(b) Suppose $P_n(z)$ and $P_{n-1}(z)$ has a common zero at $z = z_0$. Then $|z_0| > 1$ and we obtain

$$z_0 s_n \tilde{P}_{n-1}(z_0) = 0 \quad (10)$$

which gives $s_n = 0$, since $\tilde{P}_{n-1}(z_0) = 0$, ($\tilde{P}_{n-1}(z)$ has all its zeros in $|z| < 1$).

Hence $s_n \neq 0$ implies $P_n(z)$ and $P_{n-1}(z)$ do not have a common zero.