

# MATH3075/3975

## Financial Derivatives

School of Mathematics and Statistics  
University of Sydney

Semester 2, 2020

---

### Tutorial sheet 7

---

**Background: Chapter 3 – Multi-Period Market Models.**

**Exercise 1** We consider the conditional expectation  $\mathbb{E}_{\mathbb{P}}(X | \mathcal{G})$  where  $\mathcal{G}$  is generated by a finite partition  $(A_i)_{i \in I}$  of the sample space  $\Omega = \{\omega_1, \dots, \omega_k\}$ . Specifically, let  $k = 5$  and

$$A_1 = \{\omega_1, \omega_2\}, \quad A_2 = \{\omega_3\}, \quad A_3 = \{\omega_4, \omega_5\}.$$

Let the probability measure  $\mathbb{P}$  be given by

$$\mathbb{P}(\omega_1) = \mathbb{P}(\omega_2) = 0.1, \quad \mathbb{P}(\omega_3) = 0.3, \quad \mathbb{P}(\omega_4) = 0.2, \quad \mathbb{P}(\omega_5) = 0.3.$$


Consider the random variable  $X : \Omega \rightarrow \mathbb{R}$  given by  $X(\omega_i) = i$  for  $i = 1, \dots, 5$ .

- (a) Find the probability distribution of the random variable  $X$ .
- (b) Compute the conditional expectation  $\mathbb{E}_{\mathbb{P}}(X | \mathcal{G})$ .
- (c) Find the probability distribution of the random variable  $Y := \mathbb{E}_{\mathbb{P}}(X | \mathcal{G})$ .
- (d) Show that  $\mathbb{E}_{\mathbb{P}}(X) = \mathbb{E}_{\mathbb{P}}(\mathbb{E}_{\mathbb{P}}(X | \mathcal{G}))$ .

**Exercise 2** Consider the two-period market model  $\mathcal{M} = (B, S)$  with the savings account  $B$  given by

$$B_0 = 1, \quad B_1 = 1 + r, \quad B_2 = (1 + r)^2$$

with  $r = 0.25$  and the stock price  $S$  evolving according to the following diagram



- (a) Compute the probabilities of the states  $\omega_1, \omega_2, \omega_3, \omega_4$ .
- (b) Compute the conditional expectation  $\mathbb{E}_{\mathbb{P}}(S_2 | \mathcal{F}_1)$ :
- (b1) using the formula

$$\mathbb{E}_{\mathbb{P}}(S_2 | \mathcal{F}_1) = \sum_{i=1}^m \frac{\mathbb{1}_{A_i}}{\mathbb{P}(A_i)} \sum_{\omega \in A_i} S_2(\omega) \mathbb{P}(\omega),$$

- (b2) using directly the conditional probabilities.
- (c) Compute  $\mathbb{E}_{\mathbb{P}}(S_2)$  directly and using the equality

$$\mathbb{E}_{\mathbb{P}}(S_2) = \mathbb{E}_{\mathbb{P}}(\mathbb{E}_{\mathbb{P}}(S_2 | \mathcal{F}_1)).$$

**Exercise 3 (MATH3975)** Consider a finite probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and an arbitrary  $\sigma$ -field  $\mathcal{G} \subset \mathcal{F}$ . Let  $X$  be any  $\mathcal{F}$ -measurable random variable.

- (a) Show that the conditional expectation  $\mathbb{E}_{\mathbb{P}}(X | \mathcal{G})$  satisfies

$$\sum_{\omega \in G} X(\omega) \mathbb{P}(\omega) = \sum_{\omega \in G} \mathbb{E}_{\mathbb{P}}(X | \mathcal{G})(\omega) \mathbb{P}(\omega), \quad \forall G \in \mathcal{G}.$$

Deduce from this equality that  $\mathbb{E}_{\mathbb{P}}(X) = \mathbb{E}_{\mathbb{P}}(\mathbb{E}_{\mathbb{P}}(X | \mathcal{G}))$ .

- (b) Let  $\eta$  be a random variable such that  $\eta$  is  $\mathcal{G}$ -measurable and

$$\sum_{\omega \in G} X(\omega) \mathbb{P}(\omega) = \sum_{\omega \in G} \eta(\omega) \mathbb{P}(\omega), \quad \forall G \in \mathcal{G}.$$

Show that  $\eta = \mathbb{E}_{\mathbb{P}}(X | \mathcal{G})$ .

**Exercise 4 (MATH3975)** Let  $\mathbb{P}$  and  $\mathbb{Q}$  be two equivalent probability measures on a (finite) probability space  $(\Omega, \mathcal{F})$ . Let  $\mathbb{F}$  be an arbitrary filtration. For any fixed  $t = 0, 1, \dots, T$ , we denote by  $L_t$  the Radon-Nikodym density of  $\mathbb{Q}$  with respect to  $\mathbb{P}$  when  $\mathbb{Q}$  and  $\mathbb{P}$  are restricted to the  $\sigma$ -field  $\mathcal{F}_t$ .

- (a) Show that  $\mathbb{E}_{\mathbb{P}}(L_s | \mathcal{F}_t) = L_t$  for every  $0 \leq t \leq s \leq T$ . You may use part (b) in Exercise 3.
- (b) Using the abstract Bayes formula, establish the following equality, for an arbitrary  $\mathcal{F}_s$ -measurable random variable  $Y$  and for every  $0 \leq t \leq s$

$$\mathbb{E}_{\mathbb{Q}}(Y | \mathcal{F}_t) = (L_t)^{-1} \mathbb{E}_{\mathbb{P}}(Y L_s | \mathcal{F}_t).$$

- (c) Let  $M$  be a process such that  $M_t$  is  $\mathcal{F}_t$ -measurable for every  $t$ . Show that the following conditions are equivalent:
  - (i)  $\mathbb{E}_{\mathbb{Q}}(M_s | \mathcal{F}_t) = M_t$  for every  $0 \leq t \leq s \leq T$ ,
  - (ii)  $\mathbb{E}_{\mathbb{P}}(L_s M_s | \mathcal{F}_t) = L_t M_t$  for every  $0 \leq t \leq s \leq T$ .

If a process  $M$  is such that  $M_t$  is  $\mathcal{F}_t$ -measurable for every  $t$ , then we say that  $M$  is  $\mathbb{F}$ -adapted. If for an  $\mathbb{F}$ -adapted process  $M$  the equality  $\mathbb{E}_{\mathbb{Q}}(M_s | \mathcal{F}_t) = M_t$  is satisfied for every  $0 \leq t \leq s \leq T$ , then we say that  $M$  is an  $\mathbb{F}$ -martingale under  $\mathbb{Q}$ . Hence it was shown in part (c) that the following conditions are equivalent for an  $\mathbb{F}$ -adapted process  $M$ :

- (i) the process  $M$  is an  $\mathbb{F}$ -martingale under  $\mathbb{Q}$ ,
  - (ii) the process  $LM$  is an  $\mathbb{F}$ -martingale under  $\mathbb{P}$ .
- Note also that it was shown in (a) that the Radon-Nikodym density process  $L$  of  $\mathbb{Q}$  with respect to  $\mathbb{P}$  is an  $\mathbb{F}$ -martingale under  $\mathbb{P}$ .

**Exercise 5 (MATH3975)** Using the tower property of conditional expectation, show that if  $M$  is an  $\mathbb{F}$ -adapted process, then the following conditions are equivalent:

- (i) the process  $M$  is a martingale under  $\mathbb{P}$ ,
- (ii)  $\mathbb{E}_{\mathbb{P}}(M_{t+1} | \mathcal{F}_t) = M_t$  for every  $0 \leq t \leq T - 1$ ,
- (iii)  $\mathbb{E}_{\mathbb{P}}(M_T | \mathcal{F}_t) = M_t$  for every  $0 \leq t \leq T$ .

Deduce that if  $X$  an  $\mathcal{F}_T$ -measurable random variable, then the process  $M_t := \mathbb{E}_{\mathbb{P}}(X | \mathcal{F}_t)$  is the unique martingale under  $\mathbb{P}$  with the terminal value  $M_T = X$ .

**Exercise 6 (MATH3975)** Consider the process  $S$  from Exercise 2.

- (a) Show that  $S$  is not a martingale under  $\mathbb{P}$ .
- (b) Find the unique probability measure  $\mathbb{Q}$  on  $(\Omega, \mathcal{F}_2)$  such that  $S$  is a martingale under  $\mathbb{Q}$ .
- (c) Find the Radon-Nikodym density process  $L$  of  $\mathbb{Q}$  with respect to  $\mathbb{P}$  and show that  $L$  is a martingale under  $\mathbb{P}$ .