

# Exploration of non-resonant/chaotic divertor configurations at W7-X

R. Davies<sup>1</sup>, A. Kharwandikar<sup>1</sup>, K. Garcia<sup>2</sup>, D. Boeyaert<sup>2</sup>, J. Geiger<sup>1</sup>, O. Schmitz<sup>2</sup>, S. Henneberg<sup>1</sup>

<sup>1</sup>Max-Planck-Institut für Plasmaphysik, Greifswald, Germany <sup>2</sup>University of Wisconsin-Madison, Madison, USA







This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 — EUROfusion). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

This research was supported in part by the Helmholtz International Laboratory for Optimized Advanced Divertors in Stellarators (HILOADS).


email: robert.davies@ipp.mpg.de









# heat deposition in (phi, theta) (EMC3-Lite)









# deposition in (phi, theta) (EMC3-Lite)













# deposition in (phi, theta) (EMC3-Lite)







Coil A: 0.33 \* 1.109484MA = **0.3661MA** Coil B: 0.33 \* 1.109484MA = **0.3661MA** 



