



Unusually chaotic W7-X configurations at high iota

R. Davies¹, Y. Feng¹, J. Geiger¹, D. Boeyaert², K. A. Garcia², O Schmitz², J. Fellinger¹, S. Henneberg¹

¹Max-Planck-Institut für Plasmaphysik, Greifswald, Germany

²University of Wisconsin-Madison, Madison, Wisconsin

email: robert.davies@ipp.mpg.de



This work has been carried out within the framework of the EUROfusion Consortium, funded by the European Union via the Euratom Research and Training Programme (Grant Agreement No 101052200 — EUROfusion). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Commission. Neither the European Union nor the European Commission can be held responsible for them.

This research was supported in part by the Helmholtz International Laboratory for Optimized Advanced Divertors in Stellarators (HILOADS).

This work received funding from the Department of Energy (DoE) under grant numbers DE-SC0014210 and DE-FG02-93ER54222, and by the Research Forward program of the Vice Chancellor for Research and Graduate Education at UW Madison, WI, USA.



Contents

- Motivation & Background
- Configurations selected for study
- Understanding the chaos:
 - **(1) Magnetic field line trajectories (“videos”)**
 - **(2) A Target-Independent Heat Metric**
- Conclusions, Speculations and Future Work




Motivation & Background

- Magnetic field chaos* can occur in W7-X configurations
 - Vacuum fields at high iota
 - Finite beta and/or toroidal current
- What are the consequences of this chaos?
 - How does transport (heat, ionised particles) change?
 - Consequences for neutral particles/neutral exhaust?
 - What about ExB flows, recycling, resilience,

Is chaos good or bad in W7-X?

*chaos indicated by regions where good flux surfaces do not exist




(W7-X new configuration GYM000+1750)




Configurations selected for study

- “Unusually chaotic” vacuum configurations found at very high iota
 - Increase ratio of planar coil current/ non-planar coil current beyond the normal limit
 - Possible when operating at low field strength
- Planned experiment in next W7-X campaign
 - Proposal: rsd_001 (“*Exploration of non-resonant/chaotic divertor configurations at W7-X*”)
 - Configurations: GYM000+1750 , JYM+1730, TYM+1701



(W7-X new configuration GYM000+1750)

Configurations selected for study - a look at GYM





- GYM: what we learn from Poincare + connection length
 - PFCs no longer intersected by good flux surfaces
 - Web-like structures in Poincare and connection length
 - Fine structure in connection length
- Can we understand these features?

Understanding the chaos (1): Magnetic field line trajectories



- Method: follow trajectories of selected magnetic field lines




Understanding the chaos (1): Magnetic field line trajectories




- Method: follow trajectories of selected magnetic field lines


$\phi = 0^\circ$




$\phi = 18^\circ$




$\phi = 36^\circ$



$\phi = 54^\circ$



$\phi = 72^\circ$



- Leading order motion: poloidal (anticlockwise)
- Same direction as the motion of "sharp corner", but slower


Understanding the chaos (1): Magnetic field line trajectories



poloidal
motion


field periods = 4

island
dynamics




- Leading order motion: poloidal (anticlockwise)
- Next order motion: island dynamics (5/4 island chain dominant) (anticlockwise)

Understanding the chaos (1): Magnetic field line trajectories




- Additional “irregular” motion (responsible for flux surface breaking)
- Caused by trajectories making large radial excursions (flux expansion) and rotational transform increasing (“surfing the sharp corner”)

Understanding the chaos (1): Magnetic field line trajectories




- Irregular motion creates void regions/field lines “bunch up”

Understanding the chaos (1): Magnetic field line trajectories



- Irregular motion creates void regions/field lines “bunch up”

Understanding the chaos (1): Magnetic field line trajectories




- After many field periods:
 - Initial blob spread out / decorrelated (chaos: extreme sensitivity to initial (R, Z))
 - Points created near X-points start to separate.
- Theory*: Near the X-point, we have 2 **stable** and 2 **unstable** manifolds
 - Magnetic field lines on the stable manifold asymptotically approach the X-point
 - Field lines on the unstable manifold exponentially move away from the X-point.
- So by tracing field lines near the X-point we can trace out the unstable manifolds of the X-points
- **Tracing the field lines backwards** (negative phi) we can also see the **stable** manifolds

*With many thanks to C. Smiet for useful discussions

Understanding the chaos (1): Magnetic field line trajectories




A little aside: manifolds in chaotic & non-chaotic fields (homo/heteroclinic “tangle”)



Non-chaotic case: The unstable manifolds of one X-point lies on top of the stable manifold of the other, making a flux surface.




Chaotic case: The unstable manifolds of one X-point does not lie on top of the stable manifold of the other. Instead of forming a flux surface, we have a chaotic region.

Understanding the chaos (1): Magnetic field line trajectories



With many thanks to C. Smiet for useful discussions





Understanding the chaos (1): Magnetic field line trajectories



(2) Tracing the manifolds for a little bit more

With many thanks to C. Smiet for useful discussions

Understanding the chaos (1): Magnetic field line trajectories



Understanding the chaos (2): a Target-Independent Heat Metric



“Accessible”


- **Idea:** PFC-independent metric describing where heat can easily access, assuming diffusive transport

- Method:


- Discretise volume into flux tubes
 - Create “random walk” Monte Carlo particles on LCFS and track them, **ignoring** PFC interactions
 - Score each flux tube:

metric $\sim (\#MC \text{ particles per flux tube}) * (1 - \#\text{flux tubes travelled per particle})$

- More details: W7-X DCD meeting 13/02/2024
- Important physics parameter: Ratio of parallel to perpendicular diffusivity
 - $\chi = \text{perp. diffusivity}$



Understanding the chaos (2): a Target-Independent Heat Metric



- Fine structure visible at low perp. diffusivity χ
- “Inside island” structure washed out by χ , likely unimportant
- “Outside island” structure persistent with χ . Likely important for heat transport
- Exhaust performance probably resembles island divertor but with funny-shaped separatrix

Probably-unimportant
fine structure

Probably-important fine
structure (~separatrix)

Conclusions, Speculations and Future Work




- Conclusions from FL tracing (“videos”):
 - A clear hierarchy of magnetic field motion:
 - **Toroidal**
 - **Poloidal (main rotational transform)**
 - **Island dynamics (island rotational transform)**
 - **“Irregular motion” (field line surfing, bunching up)**
 - Stable/unstable manifolds emerge from X-points
 - **Creates something like an island separatrix, but chaotic**
- Conclusions from Target-Independent Heat Metric:
 - **Island separatrix appears important**
 - **“Inside island” fine structure probably washed out by perp. diffusivity**



Conclusions, Speculations and Future Work

- **Speculation:** Might be helpful to create the label “**Chaotic Island Divertor**”: PFC heat deposition not governed by good magnetic surfaces, but there’s a dominant island chain
 - Like an island divertor, but with a funny-looking “separatrix”
- In “Chaotic Island Divertors”, the important properties affecting performance are likely to be:
 - Island rotational transform
 - “Separatrix” shape (X-point manifolds)



Conclusions, Speculations and Future Work



Future work

- Similar analysis for other “Chaotic Island Divertor” candidates
- Improved methods of manifold calculation¹
- EMC3-EIRENE simulations
- Experiment planning / experiment
 - **All collaborations welcome!**

Thanks for listening

¹L. Rais & C. Smiet, EPFL. More information at CWGM NRD JA meeting on 13/06 (ask Bob for details).



APPENDIX



Target-Independent Heat Metric for high iota at finite beta

