

17.4 Gauges, Support Functions and Dual Norms

Chapter 15 §01 observed that the closed unit ball of every norm is convex and symmetric, and left the converse alone: does every convex, symmetric set arise as the unit ball of some norm? This section answers yes, by reading a norm off a convex set from the inside, the **gauge**. It then reads a function off a convex set from the outside, the **support function**, which turns §03's description of a closed convex set by half-spaces into a formula. Applied to a unit ball, the support function is the **dual norm**, the measurement Chapter 15 never gave to a linear functional, and the last part of the section proves that taking the dual twice returns the norm we started with.

Throughout, the field is $\mathbb{R}$, and we work in $\mathbb{R}^n$, $n \geq 1$, with the dot product $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{y}$ and its norm $\|\cdot\|_2$. So everything proved in §03 is available, with $V = \mathbb{R}^n$. A symbol $\|\cdot\|$ without a subscript is an **arbitrary norm on $\mathbb{R}^n$ in the sense of Definition 15.1.1, and B is its closed unit ball (Definition 15.1.2). Interior points are meant in the sense of Definition 17.3.2, which does not depend on the norm. For a set K and a real t , we write $tK = \{t\mathbf{x} : \mathbf{x} \in K\}$.**

17.4.1 Gauges

Every norm can be read off its unit ball. By (N2), for $t > 0$ we have $\|\mathbf{x}\| \leq t$ exactly when $\|\mathbf{x}/t\| \leq 1$, that is, when $\mathbf{x} \in tB$. Hence

$$\|\mathbf{x}\| = \inf\{t > 0 : \mathbf{x} \in tB\},$$

the inflation factor by which B has to be blown up before it reaches $\mathbf{x}$. The right-hand side makes sense for sets that are not unit balls, and this is the formula to study.

The gauge of a convex set measures a vector by how far the set must be inflated to reach it.

Definition 17.4.1 — Gauge

Let $K \subseteq \mathbb{R}^n$ be convex, with $\mathbf{0}$ an **interior point** of K . The **gauge** (or **Minkowski functional**) of K is the function $p_K : \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$p_K(\mathbf{x}) = \inf\{t > 0 : \mathbf{x} \in tK\}.$$

The infimum exists. Choose $r > 0$ with every $\mathbf{y}$ satisfying $\|\mathbf{y}\|_2 < r$ in K . If $t > \|\mathbf{x}\|_2/r$, then $\|\mathbf{x}/t\|_2 < r$, so $\mathbf{x}/t \in K$ and $\mathbf{x} \in tK$. So the set of admissible t is non-empty, and it is bounded below by 0; by **the completeness of $\mathbb{R}$, fact (A1) of Chapter 15's introduction**, it has a greatest lower bound, and

$$0 \leq p_K(\mathbf{x}) \leq \|\mathbf{x}\|_2/r \quad \text{for every } \mathbf{x} \in \mathbb{R}^n.$$

The interior point is what makes p_K finite. For the segment $[-\mathbf{e}_1, \mathbf{e}_1] \subseteq \mathbb{R}^2$, which has no interior points, the vector $\mathbf{e}_2$ lies in no multiple tK , and the formula would give the infimum of the empty set.

Proposition 17.4.1 — Properties of the Gauge

Let $K \subseteq \mathbb{R}^n$ be convex with $\mathbf{0}$ an interior point, and let $\mathbf{x} \in \mathbb{R}^n$.

- (a) $\mathbf{x} \in tK$ for **every** $t > p_K(\mathbf{x})$.
- (b) p_K is sublinear (Definition 17.3.3).
- (c) $\{\mathbf{x} : p_K(\mathbf{x}) < 1\} \subseteq K \subseteq \{\mathbf{x} : p_K(\mathbf{x}) \leq 1\}$.
- (d) If K is closed, then $K = \{\mathbf{x} : p_K(\mathbf{x}) \leq 1\}$.

Idea. Everything comes from one observation, (a): since $\mathbf{0} \in K$ and K is convex, shrinking a point of K towards $\mathbf{0}$ keeps it in K , so the admissible t form a whole half-line. Subadditivity is then convexity applied to $\mathbf{x}/s$ and $\mathbf{y}/t$ with weights proportional to s and t .

Proof.

- (a) Let $t > p_K(\mathbf{x})$. By the definition of an infimum there is s with $p_K(\mathbf{x}) \leq s < t$ and $\mathbf{x} \in sK$, so $\mathbf{x}/s \in K$. Since $\mathbf{0} \in K$ and K is convex,

$$\frac{\mathbf{x}}{t} = \frac{s}{t} \cdot \frac{\mathbf{x}}{s} + \left(1 - \frac{s}{t}\right) \mathbf{0} \in K,$$

that is, $\mathbf{x} \in tK$.

- (b) Let $\lambda > 0$. Since $\lambda\mathbf{x} \in tK$ exactly when $\mathbf{x} \in (t/\lambda)K$, the admissible t for $\lambda\mathbf{x}$ are λ times those for $\mathbf{x}$, and $p_K(\lambda\mathbf{x}) = \lambda p_K(\mathbf{x})$. For $\lambda = 0$, $\mathbf{0} \in tK$ for every $t > 0$, so $p_K(\mathbf{0}) = 0$. For subadditivity, let $\varepsilon > 0$, $s = p_K(\mathbf{x}) + \varepsilon$ and $t = p_K(\mathbf{y}) + \varepsilon$. By (a), $\mathbf{x}/s$ and $\mathbf{y}/t$ lie in K , so by convexity

$$\frac{\mathbf{x} + \mathbf{y}}{s + t} = \frac{s}{s + t} \cdot \frac{\mathbf{x}}{s} + \frac{t}{s + t} \cdot \frac{\mathbf{y}}{t} \in K.$$

Hence $p_K(\mathbf{x} + \mathbf{y}) \leq s + t = p_K(\mathbf{x}) + p_K(\mathbf{y}) + 2\varepsilon$, for every $\varepsilon > 0$, and so $p_K(\mathbf{x} + \mathbf{y}) \leq p_K(\mathbf{x}) + p_K(\mathbf{y})$.

- (c) If $p_K(\mathbf{x}) < 1$, then $\mathbf{x} \in 1 \cdot K$ by (a). If $\mathbf{x} \in K$, then $t = 1$ is admissible, so $p_K(\mathbf{x}) \leq 1$.
- (d) Let $p_K(\mathbf{x}) \leq 1$. By (a), $\mathbf{x} \in (1 + 1/k)K$ for every $k \geq 1$, so the points $\mathbf{x}/(1 + 1/k)$ lie in K . They converge to $\mathbf{x}$, and K is closed, so $\mathbf{x} \in K$. With (c) this gives $K = \{\mathbf{x} : p_K(\mathbf{x}) \leq 1\}$.

■

The gauge of a unit ball is its norm, by the computation that opened this section. A second example: for the rectangle $K = [-2, 2] \times [-1, 1] \subseteq \mathbb{R}^2$, a vector lies in tK exactly when $|x_1| \leq 2t$ and $|x_2| \leq t$, so

$$p_K(\mathbf{x}) = \max\left(\frac{1}{2}|x_1|, |x_2|\right),$$

a weighted ∞ -norm of the kind in Example (Inner products, weights, and a function space) (b). At the degenerate extreme, $K = \mathbb{R}^n$ lies in every tK , and p_K is identically 0: sublinear, but no norm.

The next theorem says exactly when a gauge is a norm. Its part (b) records the facts about unit balls that make the match perfect.

Theorem 17.4.2 — Convex Bodies and Norms (a) Let $K \subseteq \mathbb{R}^n$ be closed, bounded, convex and symmetric ($-K = K$), with $\mathbf{0}$ an interior point. Then p_K is a norm on $\mathbb{R}^n$, and its closed unit ball is K .

(b) Conversely, the closed unit ball B of any norm $\|\cdot\|$ on $\mathbb{R}^n$ is closed, bounded, convex and symmetric, with $\mathbf{0}$ an interior point, and $p_B = \|\cdot\|$.

So the norms on $\mathbb{R}^n$ correspond one to one with the sets described in (a).

Idea. Sublinearity is two thirds of a norm, and Proposition 17.4.1 already has it. The missing third is absolute homogeneity for a scalar $c < 0$, which is what symmetry buys, and positive definiteness, which is what boundedness buys: a non-zero vector cannot fit inside every shrunken copy tK of a bounded set. Closedness makes the unit ball of p_K equal to K rather than squeezed between the two sets of Proposition 17.4.1 (c).

Proof.

- (a) By Proposition 17.4.1 (b), p_K is sublinear, which gives (N3) and (N2) for $c \geq 0$. For $c < 0$, symmetry gives $\mathbf{x} \in tK$ if and only if $-\mathbf{x} \in t(-K) = tK$, so $p_K(-\mathbf{x}) = p_K(\mathbf{x})$ and $p_K(c\mathbf{x}) = |c|p_K(-\mathbf{x}) = |c|p_K(\mathbf{x})$. For (N1), $p_K \geq 0$ by definition. Let R be such that $\|\mathbf{y}\|_2 \leq R$ for all $\mathbf{y} \in K$. If $p_K(\mathbf{x}) = 0$, then $\mathbf{x} \in tK$ for every $t > 0$ by Proposition 17.4.1 (a), so $\|\mathbf{x}\|_2 \leq tR$ for every $t > 0$, and $\mathbf{x} = \mathbf{0}$. So p_K is a norm, and its closed unit ball $\{p_K \leq 1\}$ is K by Proposition 17.4.1 (d).
- (b) Chapter 15 §01 showed, right after Definition 15.1.2, that B is convex and symmetric. By Theorem 15.2.2 there are constants $0 < m \leq M$ with $m\|\mathbf{x}\|_2 \leq \|\mathbf{x}\| \leq M\|\mathbf{x}\|_2$. If $\|\mathbf{x}\| \leq 1$, then $\|\mathbf{x}\|_2 \leq 1/m$, so B is bounded. If $\|\mathbf{x}\|_2 < 1/M$, then $\|\mathbf{x}\| < 1$, so $\mathbf{0}$ is an interior point. If $\mathbf{x}_k \in B$ and $\mathbf{x}_k \rightarrow \mathbf{x}$, then $|\|\mathbf{x}_k\| - \|\mathbf{x}\|| \leq \|\mathbf{x}_k - \mathbf{x}\| \leq M\|\mathbf{x}_k - \mathbf{x}\|_2 \rightarrow 0$ by Lemma 15.1.1, and $\|\mathbf{x}\| \leq 1$ because a non-strict inequality survives the limit; so B is closed. Finally, $\mathbf{x} \in tB$ exactly when $t \geq \|\mathbf{x}\|$, for $t > 0$, so $p_B(\mathbf{x}) = \|\mathbf{x}\|$ (for $\mathbf{x} = \mathbf{0}$, the infimum of all $t > 0$ is 0). This proves the theorem. ■

This is the converse of Chapter 15’s “every unit ball is convex”. There, a norm produced a convex symmetric set, the ball of Definition 15.1.2, and convexity was what the triangle inequality bought. Here a convex symmetric set produces a norm, and the triangle inequality is what convexity buys: the subadditivity step in Proposition 17.4.1 (b) is the convexity of K , used once. Norms and closed bounded symmetric convex sets with $\mathbf{0}$ inside are the same objects seen two ways, one analytic, one geometric.

Each hypothesis in (a) carries one clause of Definition 15.1.1, and dropping one breaks exactly that clause.

- **Drop symmetry.** The triangle $T = \{\mathbf{x} : x_1 \geq -1, x_2 \geq -1, x_1 + x_2 \leq 1\}$, with vertices $(-1, -1)$, $(2, -1)$, $(-1, 2)$, is closed, bounded and convex, and contains the disc of radius $\frac{1}{2}$ about $\mathbf{0}$. A vector lies in tT exactly when $-x_1 \leq t$, $-x_2 \leq t$ and $x_1 + x_2 \leq t$, so

$$p_T(\mathbf{x}) = \max(-x_1, -x_2, x_1 + x_2),$$

which is never negative, since the three numbers add up to 0. This is sublinear and satisfies (N1), but $p_T((1, 1)) = 2$ while $p_T((-1, -1)) = 1$, so (N2) fails at $c = -1$.

- **Drop boundedness.** The strip $S = \{\mathbf{x} \in \mathbb{R}^2 : |x_2| \leq 1\}$ is closed, convex and symmetric, and $p_S(\mathbf{x}) = |x_2|$. Now (N2) and (N3) hold but (N1) fails: $p_S(\mathbf{e}_1) = 0$. This is the seminorm of Example (Three near misses) (b).

Quick check. Let K be the open unit disc $\{\mathbf{x} \in \mathbb{R}^2 : \|\mathbf{x}\|_2 < 1\}$, which is bounded, symmetric and convex (if $\|\mathbf{x}\|_2, \|\mathbf{y}\|_2 < 1$ and $0 \leq t \leq 1$, then $\|(1-t)\mathbf{x} + t\mathbf{y}\|_2 \leq (1-t)\|\mathbf{x}\|_2 + t\|\mathbf{y}\|_2 < 1$) but not closed. What is p_K , and what is its closed unit ball?

Solution. For $t > 0$, $\mathbf{x} \in tK$ exactly when $\|\mathbf{x}\|_2 < t$, so the admissible t form the interval $(\|\mathbf{x}\|_2, \infty)$ and $p_K(\mathbf{x}) = \|\mathbf{x}\|_2$. So p_K is a norm after all, but its closed unit ball is the closed disc, not K . Closedness is used only for the last claim of [Theorem 17.4.2](#) (a), through [Proposition 17.4.1](#) (d).

17.4.2 Support functions

§03 described a closed convex set from the outside: by Corollary 17.3.3 it is the intersection of the closed half-spaces containing it. For a fixed normal vector $\mathbf{y}$, the half-spaces $\{\mathbf{x} : \langle \mathbf{x}, \mathbf{y} \rangle \leq c\}$ containing K are those with $c \geq \langle \mathbf{x}, \mathbf{y} \rangle$ for every $\mathbf{x} \in K$, and the tightest of them has $c = \sup_{\mathbf{x} \in K} \langle \mathbf{x}, \mathbf{y} \rangle$. That number, as a function of the direction $\mathbf{y}$, deserves a name.

The support function of a set records, for each direction, how far the set reaches in that direction.

Definition 17.4.2 — Support Function

Let $K \subseteq \mathbb{R}^n$ be **non-empty**. The **support function** of K is

$$h_K(\mathbf{y}) = \sup_{\mathbf{x} \in K} \langle \mathbf{x}, \mathbf{y} \rangle, \quad \mathbf{y} \in \mathbb{R}^n,$$

with the value $+\infty$ when the numbers $\langle \mathbf{x}, \mathbf{y} \rangle$, $\mathbf{x} \in K$, are unbounded above.

When the numbers are bounded above, the supremum exists by (A1). If K is **bounded**, say $\|\mathbf{x}\|_2 \leq R$ on K , then $\langle \mathbf{x}, \mathbf{y} \rangle \leq R\|\mathbf{y}\|_2$ by Theorem 10.1.5, so h_K is real-valued. If K is **compact**, the supremum is a maximum by **the extreme value theorem, fact (A4) of Chapter 15's introduction**, since $\mathbf{x} \mapsto \langle \mathbf{x}, \mathbf{y} \rangle$ is continuous: $|\langle \mathbf{x}, \mathbf{y} \rangle - \langle \mathbf{x}', \mathbf{y} \rangle| \leq \|\mathbf{x} - \mathbf{x}'\|_2 \|\mathbf{y}\|_2$. For a unit vector $\mathbf{y}$, the hyperplane $\{\langle \cdot, \mathbf{y} \rangle = h_K(\mathbf{y})\}$ is the one with normal $\mathbf{y}$ that touches K from outside, and $h_K(\mathbf{y})$ is its signed distance from $\mathbf{0}$.

Some examples, each computed from the definition.

- **A point.** $h_{\{\mathbf{x}_0\}}(\mathbf{y}) = \langle \mathbf{x}_0, \mathbf{y} \rangle$, a linear function.
- **The Euclidean ball.** $h_{B_2}(\mathbf{y}) = \|\mathbf{y}\|_2$: at most that by Theorem 10.1.5, and attained at $\mathbf{x} = \mathbf{y}/\|\mathbf{y}\|_2$ when $\mathbf{y} \neq \mathbf{0}$.
- **The standard simplex** $\text{conv}\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$. By Theorem 17.1.3 its points are $\sum_i t_i \mathbf{e}_i$ with $t_i \geq 0$ and $\sum_i t_i = 1$, and $\sum_i t_i y_i \leq \max_i y_i$, with equality at a vertex $\mathbf{e}_k$ where y_k is largest. So $h(\mathbf{y}) = \max_i y_i$.
- **A segment.** $h_{[-\mathbf{e}_1, \mathbf{e}_1]}(\mathbf{y}) = |y_1|$ in $\mathbb{R}^2$.
- **An unbounded set.** For the half-line $\{t\mathbf{e}_1 : t \geq 0\}$, $h(\mathbf{y}) = 0$ if $y_1 \leq 0$ and $h(\mathbf{y}) = +\infty$ if $y_1 > 0$.

Proposition 17.4.3 — Properties of the Support Function

Let $K, L \subseteq \mathbb{R}^n$ be non-empty and bounded.

- (a) h_K is sublinear.
- (b) If $K \subseteq L$, then $h_K \leq h_L$.
- (c) $h_{\text{conv } K} = h_K$.

Idea. A supremum of linear functions of $\mathbf{y}$ inherits whatever inequalities each of them satisfies, which gives (a) and (b). For (c), a linear function takes on a convex combination a weighted average of its values, and an average never exceeds the largest value.

Proof.

- (a) For $\mathbf{x} \in K$, $\langle \mathbf{x}, \mathbf{y} + \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{z} \rangle \leq h_K(\mathbf{y}) + h_K(\mathbf{z})$, and taking the supremum over $\mathbf{x}$ gives $h_K(\mathbf{y} + \mathbf{z}) \leq h_K(\mathbf{y}) + h_K(\mathbf{z})$. For $t > 0$, $\langle \mathbf{x}, t\mathbf{y} \rangle = t\langle \mathbf{x}, \mathbf{y} \rangle$, and a positive factor comes out of a supremum; for $t = 0$ both sides are 0, as $K \neq \emptyset$.
- (b) A supremum over a larger set is at least as large.
- (c) Since $K \subseteq \text{conv } K$, (b) gives $h_K \leq h_{\text{conv } K}$, once we know $\text{conv } K$ is bounded: it lies in the ball $\{\|\mathbf{x}\|_2 \leq R\}$, which is convex and contains K (Definition 17.1.4). Conversely, by Theorem 17.1.3 a point of $\text{conv } K$ is $\mathbf{x} = \sum_i t_i \mathbf{x}_i$ with $\mathbf{x}_i \in K$, $t_i \geq 0$, $\sum_i t_i = 1$, and then

$$\langle \mathbf{x}, \mathbf{y} \rangle = \sum_i t_i \langle \mathbf{x}_i, \mathbf{y} \rangle \leq \sum_i t_i h_K(\mathbf{y}) = h_K(\mathbf{y}).$$

So $h_{\text{conv } K} \leq h_K$. ■

Warning. A support function cannot see gaps. The two-point set $\{\mathbf{e}_1, -\mathbf{e}_1\}$ and the segment $[-\mathbf{e}_1, \mathbf{e}_1]$ in $\mathbb{R}^2$ have the same support function $|y_1|$, by Proposition 17.4.3 (c). So h_K does not determine K in general; it determines only the convex hull, and the next theorem shows that among closed convex sets it determines the set.

Here is the promised formula. It is Corollary 17.3.3 with the family of half-spaces indexed by their normals.

Theorem 17.4.4 — The Support Function Determines a Closed Convex Set

Let $K, L \subseteq \mathbb{R}^n$ be non-empty, closed and convex. Then

- (a) $K = \{\mathbf{x} \in \mathbb{R}^n : \langle \mathbf{x}, \mathbf{y} \rangle \leq h_K(\mathbf{y}) \text{ for every } \mathbf{y} \in \mathbb{R}^n\}$;
- (b) $K \subseteq L$ if and only if $h_K(\mathbf{y}) \leq h_L(\mathbf{y})$ for every $\mathbf{y}$; in particular $K = L$ if and only if $h_K = h_L$.

Here h_K and h_L may take the value $+\infty$, with $a \leq +\infty$ for every a .

Idea. Part (a) is Corollary 17.3.3, with the half-space for the normal $\mathbf{y}$ chosen as tight as possible, at level $h_K(\mathbf{y})$. Part (b) then compares the two families of half-spaces normal by normal.

Proof.

- (a) Every $\mathbf{x} \in K$ satisfies $\langle \mathbf{x}, \mathbf{y} \rangle \leq h_K(\mathbf{y})$ by definition. Conversely let $\mathbf{x} \notin K$. By Theorem 17.3.2 there is $\mathbf{a} \neq \mathbf{0}$ with $\langle \mathbf{a}, \mathbf{z} \rangle \leq \langle \mathbf{a}, \mathbf{x} \rangle - \|\mathbf{a}\|_2^2$ for all $\mathbf{z} \in K$. Hence $h_K(\mathbf{a}) \leq \langle \mathbf{a}, \mathbf{x} \rangle - \|\mathbf{a}\|_2^2 < \langle \mathbf{x}, \mathbf{a} \rangle$, and $\mathbf{x}$ violates the condition at $\mathbf{y} = \mathbf{a}$.
- (b) If $K \subseteq L$, then $h_K \leq h_L$ because a supremum over a larger set is at least as large. Conversely, suppose $h_K \leq h_L$ and let $\mathbf{x} \in K$. For every $\mathbf{y}$, $\langle \mathbf{x}, \mathbf{y} \rangle \leq h_K(\mathbf{y}) \leq h_L(\mathbf{y})$, so $\mathbf{x} \in L$ by (a) applied to L . The last statement follows by applying this in both directions. ■

17.4.3 Polar sets

The condition $h_K(\mathbf{y}) \leq 1$ cuts out a set of directions, and that set turns out to be a mirror image of K .

Definition 17.4.3 — Polar Set

The **polar** of a set $K \subseteq \mathbb{R}^n$ is

$$K^\circ = \{\mathbf{y} \in \mathbb{R}^n : \langle \mathbf{x}, \mathbf{y} \rangle \leq 1 \text{ for every } \mathbf{x} \in K\}.$$

For non-empty K , $K^\circ = \{\mathbf{y} : h_K(\mathbf{y}) \leq 1\}$. The polar is always closed, convex and contains $\mathbf{0}$, being an intersection of closed half-spaces $\{\mathbf{y} : \langle \mathbf{x}, \mathbf{y} \rangle \leq 1\}$, one for each $\mathbf{x} \in K$. Two examples:

- $(B_2)^\circ = B_2$, because $h_{B_2}(\mathbf{y}) = \|\mathbf{y}\|_2$.
- For the single point $K = \{\mathbf{e}_1\}$ in $\mathbb{R}^2$, $K^\circ = \{\mathbf{y} : y_1 \leq 1\}$, a half-plane. Its polar consists of the $\mathbf{x}$ with $x_1 y_1 + x_2 y_2 \leq 1$ whenever $y_1 \leq 1$. Letting y_2 run over $\mathbb{R}$ forces $x_2 = 0$, letting $y_1 \rightarrow -\infty$ forces $x_1 \geq 0$, and $y_1 = 1$ forces $x_1 \leq 1$. So $K^{\circ\circ}$ is the segment $[\mathbf{0}, \mathbf{e}_1]$, which is not K .

The second example shows what the theorem must assume.

Theorem 17.4.5 — Bipolar Theorem

Let $K \subseteq \mathbb{R}^n$ be closed and convex with $\mathbf{0} \in K$. Then $K^{\circ\circ} = K$.

Idea. One inclusion is the definition read twice. For the other, cut a point $\mathbf{x} \notin K$ off with Theorem 17.3.2, and rescale the normal so that K lies in $\{\langle \cdot, \mathbf{b} \rangle \leq 1\}$ while $\mathbf{x}$ does not. That rescaled normal lies in K° and witnesses $\mathbf{x} \notin K^{\circ\circ}$; the rescaling needs $\mathbf{0} \in K$.

Proof. ($\supseteq$) Let $\mathbf{x} \in K$. For every $\mathbf{y} \in K^\circ$, $\langle \mathbf{x}, \mathbf{y} \rangle \leq 1$ by the definition of K° . Hence $\mathbf{x} \in K^{\circ\circ}$.

($\subseteq$) Let $\mathbf{x} \notin K$. By Theorem 17.3.2 there is $\mathbf{a} \neq \mathbf{0}$ with $\langle \mathbf{a}, \mathbf{z} \rangle \leq \langle \mathbf{a}, \mathbf{x} \rangle - \|\mathbf{a}\|_2^2$ for all $\mathbf{z} \in K$. Put $s = \sup_{\mathbf{z} \in K} \langle \mathbf{a}, \mathbf{z} \rangle$, which exists by (A1); since $\mathbf{0} \in K$, $s \geq 0$, and $s < \langle \mathbf{a}, \mathbf{x} \rangle$. If $s > 0$, let $\mathbf{b} = \mathbf{a}/s$. Then $\langle \mathbf{z}, \mathbf{b} \rangle \leq 1$ for all $\mathbf{z} \in K$, so $\mathbf{b} \in K^\circ$, while $\langle \mathbf{x}, \mathbf{b} \rangle > 1$. If $s = 0$, then $\langle \mathbf{a}, \mathbf{x} \rangle > 0$; let $\mathbf{b} = 2\mathbf{a}/\langle \mathbf{a}, \mathbf{x} \rangle$. Then $\langle \mathbf{z}, \mathbf{b} \rangle \leq 0 \leq 1$ on K , so $\mathbf{b} \in K^\circ$, while $\langle \mathbf{x}, \mathbf{b} \rangle = 2 > 1$. In both cases $\mathbf{x} \notin K^{\circ\circ}$. ■

The hypothesis $\mathbf{0} \in K$ was used to make $s \geq 0$, so that dividing by s keeps the direction of the inequality; the counterexample $\{\mathbf{e}_1\}$ above is exactly a closed convex set without it.

17.4.4 Dual norms

Chapter 15 measured vectors (Definition 15.1.1) and matrices (Definition 15.3.2). A third kind of object sits between them and was never measured: a linear functional on $\mathbb{R}^n$, which by Theorem 4.1.1 is $\mathbf{x} \mapsto \mathbf{y}^\top \mathbf{x} = \langle \mathbf{x}, \mathbf{y} \rangle$ for exactly one $\mathbf{y}$. How big is it?

When $\mathbf{x}$ is measured by $\|\cdot\|_2$, Theorem 10.1.5 answers: $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2$, and the constant $\|\mathbf{y}\|_2$ is the best possible. When $\mathbf{x}$ is measured by another norm, the question is the same — what is the smallest c with $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq c\|\mathbf{x}\|$ for every $\mathbf{x}$? — and the answer is the largest value of $\langle \mathbf{x}, \mathbf{y} \rangle$ over the unit ball. That is the support function of B , and it is also the operator norm of the $1 \times n$ matrix $\mathbf{y}^\top$. An expression that turns up three ways deserves a name.

The dual norm of $\mathbf{y}$ is the most that the functional $\langle \cdot, \mathbf{y} \rangle$ can extract from a vector of norm at most one.

Definition 17.4.4 — Dual Norm

Let $\|\cdot\|$ be a norm on $\mathbb{R}^n$, with closed unit ball B . The **dual norm** of $\|\cdot\|$ is the function $\|\cdot\|_*: \mathbb{R}^n \rightarrow \mathbb{R}$ given by

$$\|\mathbf{y}\|_* = \max\{\langle \mathbf{x}, \mathbf{y} \rangle : \mathbf{x} \in \mathbb{R}^n, \|\mathbf{x}\| \leq 1\} = h_B(\mathbf{y}).$$

The maximum exists. By Theorem 17.4.2 (b), B is non-empty, closed and bounded, hence compact by Corollary 15.2.5, which rests on **fact (A3) of Chapter 15's introduction**; and $\mathbf{x} \mapsto \langle \mathbf{x}, \mathbf{y} \rangle$ is continuous, so by **the extreme value theorem, fact (A4)**, it attains its maximum on B .

In words: fix $\mathbf{y}$, let $\mathbf{x}$ range over the vectors that the **given** norm calls short, and record the largest value of $\langle \mathbf{x}, \mathbf{y} \rangle$. The inner product is part of the definition, and so is the norm; changing either changes $\|\cdot\|_*$. No absolute value is needed in the maximum: B is symmetric, so replacing $\mathbf{x}$ by $-\mathbf{x}$ shows that the largest value of $\langle \mathbf{x}, \mathbf{y} \rangle$ on B is also the largest value of $|\langle \mathbf{x}, \mathbf{y} \rangle|$.

Proposition 17.4.6 — Properties of the Dual Norm

Let $\|\cdot\|$ be a norm on $\mathbb{R}^n$, with closed unit ball B .

- (a) $\|\cdot\|_*$ is a norm on $\mathbb{R}^n$.
- (b) $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\| \|\mathbf{y}\|_*$ for all $\mathbf{x}, \mathbf{y}$, and $\|\mathbf{y}\|_*$ is the smallest constant c with $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq c\|\mathbf{x}\|$ for every $\mathbf{x}$.
- (c) $\|\mathbf{y}\|_*$ is the operator norm (Definition 15.3.2) of the $1 \times n$ matrix $\mathbf{y}^\top$, with $\|\cdot\|$ on $\mathbb{R}^n$ and $|\cdot|$ on $\mathbb{R}$.
- (d) The closed unit ball of $\|\cdot\|_*$ is the polar B° .

Idea. Each norm axiom for $\|\cdot\|_*$ is checked one $\mathbf{x} \in B$ at a time and then survives the maximum; (N1) needs a good test vector, $\mathbf{y}$ itself rescaled into B . For (b), rescale any $\mathbf{x} \neq \mathbf{0}$ into B . Parts (c) and (d) only compare definitions.

Proof.

- (a) (N1): $\mathbf{x} = \mathbf{0} \in B$ gives $\|\mathbf{y}\|_* \geq 0$. If $\mathbf{y} \neq \mathbf{0}$, then $\mathbf{x} = \mathbf{y}/\|\mathbf{y}\| \in B$ gives $\|\mathbf{y}\|_* \geq \|\mathbf{y}\|_2^2/\|\mathbf{y}\| > 0$. (N2): for $c \geq 0$, $\max_B \langle \mathbf{x}, c\mathbf{y} \rangle = c \max_B \langle \mathbf{x}, \mathbf{y} \rangle$; for $c < 0$, $\langle \mathbf{x}, c\mathbf{y} \rangle = |c| \langle -\mathbf{x}, \mathbf{y} \rangle$, and $-\mathbf{x}$ runs over B as $\mathbf{x}$ does, by (N2) for $\|\cdot\|$. (N3): for $\mathbf{x} \in B$, $\langle \mathbf{x}, \mathbf{y} + \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{z} \rangle \leq \|\mathbf{y}\|_* + \|\mathbf{z}\|_*$; take the maximum over $\mathbf{x}$.
- (b) For $\mathbf{x} = \mathbf{0}$ there is nothing to prove. For $\mathbf{x} \neq \mathbf{0}$, $\mathbf{x}/\|\mathbf{x}\| \in B$, so $\pm \langle \mathbf{x}, \mathbf{y} \rangle / \|\mathbf{x}\| = \langle \pm \mathbf{x} / \|\mathbf{x}\|, \mathbf{y} \rangle \leq \|\mathbf{y}\|_*$, using that B is symmetric. If c is any constant with $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq c\|\mathbf{x}\|$ for all $\mathbf{x}$, then $\langle \mathbf{x}, \mathbf{y} \rangle \leq c$ on B , so $\|\mathbf{y}\|_* \leq c$.
- (c) By Definition 15.3.2 and the equivalent descriptions that follow it in Chapter 15 §03, the operator norm of $\mathbf{y}^\top$ is $\max\{\|\mathbf{y}^\top \mathbf{x}\| : \|\mathbf{x}\| \leq 1\}$, which is $\|\mathbf{y}\|_*$ by the remark after Definition 17.4.4.
- (d) $\|\mathbf{y}\|_* \leq 1$ exactly when $\langle \mathbf{x}, \mathbf{y} \rangle \leq 1$ for every $\mathbf{x} \in B$, which is the definition of $\mathbf{y} \in B^\circ$ (Definition 17.4.3).

■

Part (b) is Cauchy–Schwarz for an arbitrary norm, with the price that the two factors are measured differently. Now the three norms of Chapter 15.

Proposition 17.4.7 — Duals of the 1-, 2- and Infinity-Norms

On $\mathbb{R}^n$ with the dot product,

- (a) the dual of $\|\cdot\|_1$ is $\|\cdot\|_\infty$;
- (b) the dual of $\|\cdot\|_\infty$ is $\|\cdot\|_1$;
- (c) the dual of $\|\cdot\|_2$ is $\|\cdot\|_2$.

Idea. Each part is an upper bound and a witness, the same pattern as Theorem 15.3.3. For the 1-norm, all of the budget $\|\mathbf{x}\|_1 \leq 1$ goes on the coordinate where $|y_k|$ is largest. For the ∞ -norm, every coordinate can be spent at once, each with the sign of y_i , so nothing cancels. For the 2-norm, the witness is $\mathbf{y}$ itself, rescaled.

Proof.

- (a) For $\|\mathbf{x}\|_1 \leq 1$, $\langle \mathbf{x}, \mathbf{y} \rangle \leq \sum_i |x_i| |y_i| \leq \|\mathbf{y}\|_\infty \sum_i |x_i| \leq \|\mathbf{y}\|_\infty$. Choose k with $|y_k| = \|\mathbf{y}\|_\infty$, and let $\mathbf{x} = \pm \mathbf{e}_k$ with the sign of y_k (either sign if $y_k = 0$). Then $\|\mathbf{x}\|_1 = 1$

- and $\langle \mathbf{x}, \mathbf{y} \rangle = |y_k| = \|\mathbf{y}\|_\infty$.
- (b) For $\|\mathbf{x}\|_\infty \leq 1$, $\langle \mathbf{x}, \mathbf{y} \rangle \leq \sum_i |x_i| |y_i| \leq \sum_i |y_i| = \|\mathbf{y}\|_1$. Let $x_i = 1$ if $y_i \geq 0$ and $x_i = -1$ if $y_i < 0$. Then $\|\mathbf{x}\|_\infty = 1$ and $\langle \mathbf{x}, \mathbf{y} \rangle = \sum_i |y_i| = \|\mathbf{y}\|_1$.
- (c) For $\|\mathbf{x}\|_2 \leq 1$, $\langle \mathbf{x}, \mathbf{y} \rangle \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2 \leq \|\mathbf{y}\|_2$ by Theorem 10.1.5. For $\mathbf{y} \neq \mathbf{0}$, $\mathbf{x} = \mathbf{y}/\|\mathbf{y}\|_2$ gives equality; for $\mathbf{y} = \mathbf{0}$ both sides are 0.

■

So the 1- and ∞ -norms are each other's duals, and the Euclidean norm is its own. In the language of polars, (a) and (b) say $(B_1)^\circ = B_\infty$ and $(B_\infty)^\circ = B_1$: the diamond and the square of Chapter 15's picture are polar to each other, and the disc is polar to itself. Part (b) of Proposition 17.4.6 now reads $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq \|\mathbf{x}\|_1 \|\mathbf{y}\|_\infty$, the simplest of a family of inequalities. The general member pairs $\|\cdot\|_p$ with $\|\cdot\|_q$, $\frac{1}{p} + \frac{1}{q} = 1$; it is **Hölder's inequality**, and §11 of this chapter proves it and deduces that the dual of $\|\cdot\|_p$ is $\|\cdot\|_q$ for every $1 < p < \infty$. The degenerate case $n = 1$ is worth one line: every norm on $\mathbb{R}$ is $\|x\| = a|x|$ with $a > 0$, its ball is $[-1/a, 1/a]$, and $\|y\|_* = |y|/a$.

A non-example by minimal change. Keep the formula $\max_K \langle \mathbf{x}, \mathbf{y} \rangle$, and replace the symmetric ball by the triangle $K = \text{conv}\{(-1, -1), (2, -1), (-1, 2)\}$, which is compact and convex and has $\mathbf{0}$ inside. By Proposition 17.4.3 (c), $h_K(\mathbf{y})$ is the largest of $-y_1 - y_2$, $2y_1 - y_2$ and $-y_1 + 2y_2$. It is sublinear by Proposition 17.4.3 (a), and positive at every $\mathbf{y} \neq \mathbf{0}$, since the three numbers add up to 0 and are not all zero. But $h_K(\mathbf{e}_1) = 2$ and $h_K(-\mathbf{e}_1) = 1$, so (N2) fails at $c = -1$. Symmetry of the ball is exactly what makes a dual norm a norm.

Why this definition, and not another? The measurement of a functional must be compatible with the measurement of its inputs, in the sense that $|\langle \mathbf{x}, \mathbf{y} \rangle| \leq c\|\mathbf{x}\|$ for every $\mathbf{x}$, and Proposition 17.4.6 (b) says that $\|\mathbf{y}\|_*$ is the **smallest** compatible c . Any larger choice would be a bound, not a measurement. Abstractly, the dual norm lives on the dual space: for $\varphi \in (\mathbb{R}^n)^*$ one would put $\|\varphi\| = \max_{\|\mathbf{x}\| \leq 1} \varphi(\mathbf{x})$, and the dot product is used only to name φ by a vector, through Theorem 4.1.1. That is why a different inner product on $\mathbb{R}^n$ gives a different formula for “the same” dual norm. The name is standard: the dual space carries the dual norm.

Warning. **A larger norm has a smaller dual.** If $\|\mathbf{x}\|_a \leq \|\mathbf{x}\|_b$ for all $\mathbf{x}$, then the ball of $\|\cdot\|_b$ lies inside the ball of $\|\cdot\|_a$, and a maximum over a smaller set is smaller: $\|\mathbf{y}\|_{b*} \leq \|\mathbf{y}\|_{a*}$. For instance $N(\mathbf{x}) = 2\|\mathbf{x}\|_2$ has ball of radius $\frac{1}{2}$ and dual $N_*(\mathbf{y}) = \frac{1}{2}\|\mathbf{y}\|_2$, not $2\|\mathbf{y}\|_2$. Scaling a norm up scales its dual down.

Chapter 15 measured vectors and matrices but never a linear functional; Definition 17.4.4 fills that gap, and Proposition 17.4.6 (c) shows it is exactly an operator norm of Chapter 15, the one for $1 \times n$ matrices. The one question it leaves open is whether the construction loses information.

17.4.5 The dual of the dual

Apply the construction twice. Since $\|\cdot\|_*$ is a norm, it has a dual $\|\cdot\|_{**}$, and the natural guess is that we are back where we started. For the three norms of Proposition 17.4.7 this is visible: $1 \rightarrow \infty \rightarrow 1$ and $2 \rightarrow 2 \rightarrow 2$. In general it is a theorem, and its proof is a supporting hyperplane.

Theorem 17.4.8 — The Dual of the Dual Norm

For every norm $\|\cdot\|$ on $\mathbb{R}^n$, $\|\cdot\|_{**} = \|\cdot\|$.

Idea. One inequality is Cauchy–Schwarz for dual norms. For the other, take $\mathbf{x}$ on the unit sphere of $\|\cdot\|$. It is on the edge of the ball B , so §03 provides a supporting hyperplane of B at $\mathbf{x}$. Its normal $\mathbf{a}$ is a direction in which B reaches no further than $\mathbf{x}$ does, so the maximum defining $\|\mathbf{a}\|_*$ is attained at $\mathbf{x}$; rescaling $\mathbf{a}$ produces a functional of dual norm 1 that takes the value 1 at $\mathbf{x}$.

Proof. Let $\mathbf{x} \in \mathbb{R}^n$. For every $\mathbf{y}$ with $\|\mathbf{y}\|_* \leq 1$, Proposition 17.4.6 (b) gives $\langle \mathbf{x}, \mathbf{y} \rangle \leq \|\mathbf{x}\| \|\mathbf{y}\|_* \leq \|\mathbf{x}\|$. By Definition 17.4.4 applied to $\|\cdot\|_*$, whose unit ball is $\{\mathbf{y} : \|\mathbf{y}\|_* \leq 1\}$, this says $\|\mathbf{x}\|_{**} \leq \|\mathbf{x}\|$.

For the reverse inequality, both sides vanish at $\mathbf{x} = \mathbf{0}$, and both are absolutely homogeneous, so it suffices to take $\|\mathbf{x}\| = 1$. Then $\mathbf{x} \in B$, but $\mathbf{x}$ is not an interior point of B : the vectors $(1 + 1/k)\mathbf{x}$ have norm $1 + 1/k > 1$ and $\|(1 + 1/k)\mathbf{x} - \mathbf{x}\|_2 = \|\mathbf{x}\|_2/k \rightarrow 0$. By Theorem 17.3.6, applied to the convex set B and the point $\mathbf{x}$, there is $\mathbf{a} \neq \mathbf{0}$ with $\langle \mathbf{z}, \mathbf{a} \rangle \leq \langle \mathbf{x}, \mathbf{a} \rangle$ for every $\mathbf{z} \in B$. Since $\mathbf{x} \in B$, the maximum in Definition 17.4.4 is attained at $\mathbf{x}$:

$$\|\mathbf{a}\|_* = \max_{\mathbf{z} \in B} \langle \mathbf{z}, \mathbf{a} \rangle = \langle \mathbf{x}, \mathbf{a} \rangle.$$

By Proposition 17.4.6 (a), $\|\mathbf{a}\|_* > 0$. Let $\mathbf{y} = \mathbf{a}/\|\mathbf{a}\|_*$. Then $\|\mathbf{y}\|_* = 1$ and $\langle \mathbf{x}, \mathbf{y} \rangle = 1$. Hence $\|\mathbf{x}\|_{**} \geq 1 = \|\mathbf{x}\|$. This proves the theorem. ■

Alternatively, the vector $\mathbf{y}$ can be obtained from Hahn–Banach. Corollary 17.3.9, applied to $U = \text{span}(\mathbf{x})$ and $f(t\mathbf{x}) = t\|\mathbf{x}\|$, gives a functional F with $F(\mathbf{x}) = \|\mathbf{x}\|$ and $|F| \leq \|\cdot\|$; writing $F = \langle \cdot, \mathbf{y} \rangle$ by Theorem 4.1.1, the second property says $\|\mathbf{y}\|_* \leq 1$ by Proposition 17.4.6 (b), so $\|\mathbf{x}\|_{**} \geq \langle \mathbf{x}, \mathbf{y} \rangle = \|\mathbf{x}\|$. So the analytic and the geometric forms of §03 each prove the theorem, which is one more sign that they are the same fact.

Read with the definition of $\|\cdot\|_{**}$ unwound, the theorem is a formula for the original norm.

Corollary 17.4.9 — A Norm Is a Maximum of Linear Functions

For every norm $\|\cdot\|$ on $\mathbb{R}^n$ and every $\mathbf{x} \in \mathbb{R}^n$,

$$\|\mathbf{x}\| = \max\{\langle \mathbf{x}, \mathbf{y} \rangle : \|\mathbf{y}\|_* \leq 1\},$$

and when $\mathbf{x} \neq \mathbf{0}$ the maximum is attained at some $\mathbf{y}$ with $\|\mathbf{y}\|_* = 1$.

Proof. The right-hand side is $\|\mathbf{x}\|_{**}$ by Definition 17.4.4 applied to $\|\cdot\|_*$, and it equals $\|\mathbf{x}\|$ by Theorem 17.4.8. For $\mathbf{x} \neq \mathbf{0}$, the proof of Theorem 17.4.8, applied to $\mathbf{x}/\|\mathbf{x}\|$, produced $\mathbf{y}$ with $\|\mathbf{y}\|_* = 1$ and $\langle \mathbf{x}/\|\mathbf{x}\|, \mathbf{y} \rangle = 1$, so $\langle \mathbf{x}, \mathbf{y} \rangle = \|\mathbf{x}\|$. ■

So every norm is the support function of the unit ball of its dual, $\|\cdot\| = h_{\{\mathbf{y} : \|\mathbf{y}\|_* \leq 1\}}$: a maximum of the linear functions $\mathbf{x} \mapsto \langle \mathbf{x}, \mathbf{y} \rangle$, one for each $\mathbf{y}$ in a compact set. §10 of this chapter shows that a pointwise maximum of linear functions is always convex, and norms (their convexity was already the triangle inequality) are among its first examples.

Quick check. For $\mathbf{x} = (3, -4)$, find a $\mathbf{y}$ with $\|\mathbf{y}\|_\infty = 1$ and $\langle \mathbf{x}, \mathbf{y} \rangle = \|\mathbf{x}\|_1$, and a $\mathbf{y}$ with $\|\mathbf{y}\|_1 = 1$ and $\langle \mathbf{x}, \mathbf{y} \rangle = \|\mathbf{x}\|_\infty$.

Solution. The dual of $\|\cdot\|_1$ is $\|\cdot\|_\infty$ (Proposition 17.4.7 (a)), so the first $\mathbf{y}$ is a sign vector: $\mathbf{y} = (1, -1)$ gives $3 + 4 = 7 = \|\mathbf{x}\|_1$. The dual of $\|\cdot\|_\infty$ is $\|\cdot\|_1$, so the second $\mathbf{y}$ puts all its weight on the largest coordinate: $\mathbf{y} = (0, -1)$ gives $4 = \|\mathbf{x}\|_\infty$. Each is the witness from the proof of Proposition 17.4.7, with the roles of $\mathbf{x}$ and $\mathbf{y}$ exchanged.

17.4.6 Operator norms as maxima of bilinear values

The corollary also rewrites Chapter 15's operator norm. Fix a norm $\|\cdot\|_\alpha$ on $\mathbb{R}^n$ and a norm $\|\cdot\|_\beta$ on $\mathbb{R}^m$, and write $\|\mathbf{A}\|_{\alpha \rightarrow \beta}$ for the operator norm of $\mathbf{A} \in M_{m \times n}(\mathbb{R})$ that they induce (Definition 15.3.2). By the equivalent descriptions after that definition, $\|\mathbf{A}\|_{\alpha \rightarrow \beta} = \max\{\|\mathbf{Ax}\|_\beta : \|\mathbf{x}\|_\alpha \leq 1\}$.

Proposition 17.4.10 — Operator Norms through Dual Norms

Let $\mathbf{A} \in M_{m \times n}(\mathbb{R})$, and let $\|\cdot\|_\alpha, \|\cdot\|_\beta$ be norms on $\mathbb{R}^n$ and $\mathbb{R}^m$, with duals $\|\cdot\|_{\alpha*}$ and $\|\cdot\|_{\beta*}$. Then

- (a) $\|\mathbf{A}\|_{\alpha \rightarrow \beta} = \max\{\mathbf{y}^\top \mathbf{Ax} : \|\mathbf{x}\|_\alpha \leq 1, \|\mathbf{y}\|_{\beta*} \leq 1\}$;
- (b) $\|\mathbf{A}^\top\|_{\beta* \rightarrow \alpha*} = \|\mathbf{A}\|_{\alpha \rightarrow \beta}$.

Idea. Write the outer norm $\|\mathbf{Ax}\|_\beta$ as a maximum over its dual ball by Corollary 17.4.9, and the operator norm becomes one maximum over pairs $(\mathbf{x}, \mathbf{y})$. Since $\mathbf{y}^\top \mathbf{Ax} = \mathbf{x}^\top \mathbf{A}^\top \mathbf{y}$, the same double maximum, taken with $\mathbf{x}$ innermost, is the operator norm of $\mathbf{A}^\top$, which gives (b).

Proof.

- (a) For $\|\mathbf{x}\|_\alpha \leq 1$, Corollary 17.4.9 in $\mathbb{R}^m$ gives $\|\mathbf{Ax}\|_\beta = \max\{\mathbf{y}^\top \mathbf{Ax} : \|\mathbf{y}\|_{\beta*} \leq 1\}$. So every value $\mathbf{y}^\top \mathbf{Ax}$ in the set is at most $\|\mathbf{Ax}\|_\beta \leq \|\mathbf{A}\|_{\alpha \rightarrow \beta}$. Conversely, choose $\mathbf{x}_0$ with $\|\mathbf{x}_0\|_\alpha \leq 1$ and $\|\mathbf{Ax}_0\|_\beta = \|\mathbf{A}\|_{\alpha \rightarrow \beta}$, which exists by Lemma 15.3.1, and then $\mathbf{y}_0$ attaining the maximum for $\|\mathbf{Ax}_0\|_\beta$. The value $\mathbf{y}_0^\top \mathbf{Ax}_0$ equals $\|\mathbf{A}\|_{\alpha \rightarrow \beta}$, so the maximum exists and has this value.
- (b) By the equivalent descriptions after Definition 15.3.2, and then Definition 17.4.4 for $\|\cdot\|_{\alpha*}$,

$$\|\mathbf{A}^\top\|_{\beta* \rightarrow \alpha*} = \max_{\|\mathbf{y}\|_{\beta*} \leq 1} \|\mathbf{A}^\top \mathbf{y}\|_{\alpha*} = \max_{\|\mathbf{y}\|_{\beta*} \leq 1} \max_{\|\mathbf{x}\|_\alpha \leq 1} \langle \mathbf{x}, \mathbf{A}^\top \mathbf{y} \rangle.$$

Since $\langle \mathbf{x}, \mathbf{A}^\top \mathbf{y} \rangle = \mathbf{x}^\top \mathbf{A}^\top \mathbf{y} = \mathbf{y}^\top \mathbf{Ax}$, this is the maximum in (a).

■

With $\alpha = \beta = \infty$, part (b) and Proposition 17.4.7 say $\|\mathbf{A}^\top\|_1 = \|\mathbf{A}\|_\infty$: the largest column

sum of $\mathbf{A}^\top$ is the largest row sum of $\mathbf{A}$, which is also what Theorem 15.3.3 gives directly. With $\alpha = \beta = 2$ it says $\|\mathbf{A}^\top\|_2 = \|\mathbf{A}\|_2$, that is, $\sigma_1(\mathbf{A}^\top) = \sigma_1(\mathbf{A})$. For the matrix $\mathbf{A} = \begin{pmatrix} 3 & 0 \\ 4 & 5 \end{pmatrix}$ of Example (One matrix, four numbers), $\|\mathbf{A}\|_1 = 7$ and $\|\mathbf{A}\|_\infty = 9$. Its transpose $\begin{pmatrix} 3 & 4 \\ 0 & 5 \end{pmatrix}$ has column sums 3 and 9 and row sums 7 and 5, so $\|\mathbf{A}^\top\|_1 = 9 = \|\mathbf{A}\|_\infty$ and $\|\mathbf{A}^\top\|_\infty = 7 = \|\mathbf{A}\|_1$, as (b) predicts.

17.4.7 Exercises

17.4.7.1 A. Check your understanding

Exercise A1.

- Define the gauge p_K , the support function h_K and the dual norm $\|\cdot\|_*$, stating what each needs of the set or norm involved.
- Determine whether the following statement is correct, and justify your answer: the support function of every non-empty bounded subset of $\mathbb{R}^2$ is a norm.
- What is the dual of $\|\cdot\|_\infty$ on $\mathbb{R}^n$? What is the dual of that?
- Which result of §03 does the proof of Theorem 17.4.8 use, and at which point?
- Determine whether the following statement is correct, and justify your answer: $K^{\circ\circ} = K$ for every non-empty closed convex $K \subseteq \mathbb{R}^n$.

Solution.

- For K convex with $\mathbf{0}$ an interior point, $p_K(\mathbf{x}) = \inf\{t > 0 : \mathbf{x} \in tK\}$ (Definition 17.4.1). For K non-empty, $h_K(\mathbf{y}) = \sup_{\mathbf{x} \in K} \langle \mathbf{x}, \mathbf{y} \rangle$, possibly $+\infty$ (Definition 17.4.2). For a norm $\|\cdot\|$ with ball B , $\|\mathbf{y}\|_* = \max_{\mathbf{x} \in B} \langle \mathbf{x}, \mathbf{y} \rangle = h_B(\mathbf{y})$ (Definition 17.4.4).
- Incorrect. For $K = \{\mathbf{0}\}$, $h_K = 0$, which violates (N1). For the segment $[-\mathbf{e}_1, \mathbf{e}_1]$, $h_K(\mathbf{y}) = |y_1|$ vanishes at $\mathbf{e}_2$. Support functions are sublinear (Proposition 17.4.3 (a)) but need not be norms.
- $\|\cdot\|_1$, by Proposition 17.4.7 (b); and the dual of $\|\cdot\|_1$ is $\|\cdot\|_\infty$ again, by (a) of the same proposition, as Theorem 17.4.8 requires.
- Theorem 17.3.6, applied to the unit ball B at a point $\mathbf{x}$ with $\|\mathbf{x}\| = 1$, to produce a normal $\mathbf{a}$ at which the maximum defining $\|\mathbf{a}\|_*$ is attained at $\mathbf{x}$.
- Incorrect. For $K = \{\mathbf{e}_1\}$ in $\mathbb{R}^2$, $K^{\circ\circ}$ is the segment $[\mathbf{0}, \mathbf{e}_1]$. Theorem 17.4.5 needs $\mathbf{0} \in K$.

17.4.7.2 B. Practice

Exercise B1: A weighted norm and its dual.

On $\mathbb{R}^2$ let $N(\mathbf{x}) = \max(|x_1|, 2|x_2|)$.

- Prove that $N_*(\mathbf{y}) = |y_1| + \frac{1}{2}|y_2|$.
- Verify Theorem 17.4.8 for N by computing N_{**} directly.

Solution.

- (a) The ball of N is the rectangle $|x_1| \leq 1, |x_2| \leq \frac{1}{2}$. On it, $\langle \mathbf{x}, \mathbf{y} \rangle \leq |x_1||y_1| + |x_2||y_2| \leq |y_1| + \frac{1}{2}|y_2|$. The point $\mathbf{x} = (s_1, \frac{1}{2}s_2)$, where $s_i = 1$ if $y_i \geq 0$ and $s_i = -1$ otherwise, lies in the ball and gives equality. So $N_*(\mathbf{y}) = |y_1| + \frac{1}{2}|y_2|$.
- (b) The ball of N_* is $\{|y_1| + \frac{1}{2}|y_2| \leq 1\}$. On it,

$$\langle \mathbf{x}, \mathbf{y} \rangle \leq |x_1||y_1| + 2|x_2| \cdot \frac{1}{2}|y_2| \leq N(\mathbf{x})(|y_1| + \frac{1}{2}|y_2|) \leq N(\mathbf{x}).$$

If $N(\mathbf{x}) = |x_1|$, the point $\mathbf{y} = \pm \mathbf{e}_1$ (sign of x_1) lies in the ball and gives $\langle \mathbf{x}, \mathbf{y} \rangle = |x_1|$.
 If $N(\mathbf{x}) = 2|x_2|$, the point $\mathbf{y} = \pm 2\mathbf{e}_2$ (sign of x_2) lies in the ball, since $\frac{1}{2} \cdot 2 = 1$, and gives $\langle \mathbf{x}, \mathbf{y} \rangle = 2|x_2|$. So $N_{**} = N$.

Exercise B2: A hexagonal norm.

Let $K = \{\mathbf{x} \in \mathbb{R}^2 : |x_1| \leq 1, |x_2| \leq 1, |x_1 - x_2| \leq 1\}$.

- (a) Check that K satisfies the hypotheses of [Theorem 17.4.2](#) (a).
 (b) Show that $p_K(\mathbf{x}) = \max(|x_1|, |x_2|, |x_1 - x_2|)$, and compute $p_K((2, -1))$.
 (c) Determine whether p_K is induced by an inner product. Justify your answer.

Solution.

- (a) K is the intersection of six closed half-planes, such as $\{x_1 \leq 1\}$ and $\{x_1 - x_2 \geq -1\}$, each closed and convex (§03, before Definition 17.3.1); so K is closed and convex. It is bounded, since $\|\mathbf{x}\|_2 \leq \sqrt{2}$ when $|x_1|, |x_2| \leq 1$. It is symmetric, since each condition is unchanged by $\mathbf{x} \mapsto -\mathbf{x}$. And if $\|\mathbf{x}\|_2 < \frac{1}{2}$, then $|x_1|, |x_2| < \frac{1}{2}$ and $|x_1 - x_2| < 1$, so $\mathbf{x} \in K$ and $\mathbf{0}$ is an interior point.
- (b) For $t > 0$, $\mathbf{x} \in tK$ exactly when $\mathbf{x}/t \in K$, that is, when $|x_1| \leq t$, $|x_2| \leq t$ and $|x_1 - x_2| \leq t$. The infimum of such t is $M = \max(|x_1|, |x_2|, |x_1 - x_2|)$ when $\mathbf{x} \neq \mathbf{0}$, and $0 = M$ when $\mathbf{x} = \mathbf{0}$. So $p_K((2, -1)) = \max(2, 1, 3) = 3$.
- (c) No. With $\mathbf{u} = \mathbf{e}_1$ and $\mathbf{v} = \mathbf{e}_2$, $p_K(\mathbf{u} + \mathbf{v}) = \max(1, 1, 0) = 1$ and $p_K(\mathbf{u} - \mathbf{v}) = \max(1, 1, 2) = 2$, so the left side of the parallelogram law is $1 + 4 = 5$, while the right side is $2 \cdot 1 + 2 \cdot 1 = 4$. By Theorem 15.1.3, no inner product induces p_K .

Exercise B3: The polar of a triangle.

Let $K = \text{conv}\{(-1, -1), (2, -1), (-1, 2)\} \subseteq \mathbb{R}^2$.

- (a) Compute $h_K(\mathbf{e}_1)$, $h_K(-\mathbf{e}_1)$ and $h_K((1, 1))$.
 (b) Describe K° by three linear inequalities.
 (c) Show that $(-1, 0)$, $(0, -1)$ and $(1, 1)$ lie in K° , and that $(1, 0)$ does not.

Solution.

- (a) By [Proposition 17.4.3](#) (c), $h_K(\mathbf{y})$ is the largest of $\langle \mathbf{v}, \mathbf{y} \rangle$ over the three vertices $\mathbf{v}$, that is, of $-y_1 - y_2$, $2y_1 - y_2$, $-y_1 + 2y_2$. So $h_K(\mathbf{e}_1) = \max(-1, 2, -1) = 2$, $h_K(-\mathbf{e}_1) = \max(1, -2, 1) = 1$, and $h_K((1, 1)) = \max(-2, 1, 1) = 1$.

(b) $\mathbf{y} \in K^\circ$ exactly when $h_K(\mathbf{y}) \leq 1$, that is,

$$-y_1 - y_2 \leq 1, \quad 2y_1 - y_2 \leq 1, \quad -y_1 + 2y_2 \leq 1.$$

(c) At $(-1, 0)$ the three left sides are $1, -2, 1$; at $(0, -1)$ they are $1, 1, -2$; at $(1, 1)$ they are $-2, 1, 1$. All are ≤ 1 . At $(1, 0)$ the second is $2 > 1$, so $(1, 0) \notin K^\circ$, in agreement with $h_K(\mathbf{e}_1) = 2$.

17.4.7.3 C. Going deeper

Exercise C1: The norm from 1 to infinity.

Let $\mathbf{A} \in M_{m \times n}(\mathbb{R})$, and let $\|\mathbf{A}\|_{1 \rightarrow \infty}$ be the operator norm with $\|\cdot\|_1$ on $\mathbb{R}^n$ and $\|\cdot\|_\infty$ on $\mathbb{R}^m$.

- (a) Using Proposition 17.4.10 (a), prove that $\|\mathbf{A}\|_{1 \rightarrow \infty} = \max_{i,j} |a_{ij}|$.
- (b) Deduce from Proposition 17.4.10 (b) that the operator norm of $\mathbf{A}^\top$ with $\|\cdot\|_1$ on $\mathbb{R}^m$ and $\|\cdot\|_\infty$ on $\mathbb{R}^n$ is the same number.
- (c) Explain why this does not contradict the warning of Chapter 15 §03 that the entrywise maximum is not a matrix norm.

Solution.

- (a) By Proposition 17.4.7 (b), the dual of $\|\cdot\|_\infty$ is $\|\cdot\|_1$, so Proposition 17.4.10 (a) gives $\|\mathbf{A}\|_{1 \rightarrow \infty} = \max\{\mathbf{y}^\top \mathbf{A} \mathbf{x} : \|\mathbf{x}\|_1 \leq 1, \|\mathbf{y}\|_1 \leq 1\}$. Write $M = \max_{i,j} |a_{ij}|$. For such $\mathbf{x}, \mathbf{y}$,

$$\mathbf{y}^\top \mathbf{A} \mathbf{x} = \sum_{i,j} y_i a_{ij} x_j \leq M \sum_i |y_i| \sum_j |x_j| \leq M.$$

If $|a_{kl}| = M$, then $\mathbf{x} = \mathbf{e}_l$ and $\mathbf{y} = \pm \mathbf{e}_k$, with the sign of a_{kl} , give $\mathbf{y}^\top \mathbf{A} \mathbf{x} = M$. So $\|\mathbf{A}\|_{1 \rightarrow \infty} = M$.

- (b) By Proposition 17.4.10 (b) with $\alpha = 1$ and $\beta = \infty$, $\|\mathbf{A}^\top\|_{\beta* \rightarrow \alpha*} = \|\mathbf{A}^\top\|_{1 \rightarrow \infty}$ equals $\|\mathbf{A}\|_{1 \rightarrow \infty}$, using $\beta* = 1$ and $\alpha* = \infty$ from Proposition 17.4.7. This is consistent with (a), since $\mathbf{A}$ and $\mathbf{A}^\top$ have the same entries.
- (c) $\|\cdot\|_{1 \rightarrow \infty}$ uses **different** norms on the source and the target, so the product $\mathbf{A}\mathbf{B}$ of two square matrices is measured by a norm that does not compose: submultiplicativity in Theorem 15.3.2 (d) needs the target norm of $\mathbf{B}$ to be the source norm of $\mathbf{A}$. With one norm on $M_n(\mathbb{R})$ the entrywise maximum is not submultiplicative (Example (The entrywise maximum is not submultiplicative)), and nothing here says otherwise.

Exercise C2: Equivalence constants dualize.

Let N_1, N_2 be norms on $\mathbb{R}^n$ and $0 < m \leq M$ constants with $mN_1(\mathbf{x}) \leq N_2(\mathbf{x}) \leq MN_1(\mathbf{x})$ for every $\mathbf{x}$.

- (a) Prove that $\frac{1}{M}N_{1*}(\mathbf{y}) \leq N_{2*}(\mathbf{y}) \leq \frac{1}{m}N_{1*}(\mathbf{y})$ for every $\mathbf{y}$.
- (b) Apply (a) to $\|\mathbf{x}\|_2 \leq \|\mathbf{x}\|_1 \leq \sqrt{n}\|\mathbf{x}\|_2$ from Proposition 15.1.2, and compare the result with that proposition.

Solution.

- (a) If $N_2(\mathbf{x}) \leq 1$, then $N_1(\mathbf{x}) \leq 1/m$, so $\mathbf{x} = \frac{1}{m}\mathbf{x}'$ with $N_1(\mathbf{x}') \leq 1$, and $\langle \mathbf{x}, \mathbf{y} \rangle = \frac{1}{m}\langle \mathbf{x}', \mathbf{y} \rangle \leq \frac{1}{m}N_{1*}(\mathbf{y})$. Taking the maximum over the ball of N_2 gives $N_{2*}(\mathbf{y}) \leq \frac{1}{m}N_{1*}(\mathbf{y})$. If $N_1(\mathbf{x}) \leq 1$, then $N_2(\mathbf{x}) \leq M$, and [Proposition 17.4.6](#) (b) gives $\langle \mathbf{x}, \mathbf{y} \rangle \leq N_2(\mathbf{x})N_{2*}(\mathbf{y}) \leq MN_{2*}(\mathbf{y})$; so $N_{1*}(\mathbf{y}) \leq MN_{2*}(\mathbf{y})$.
- (b) Take $N_1 = \|\cdot\|_2$, $N_2 = \|\cdot\|_1$, $m = 1$, $M = \sqrt{n}$. The duals are $\|\cdot\|_2$ and $\|\cdot\|_\infty$ ([Proposition 17.4.7](#)), so (a) gives $\frac{1}{\sqrt{n}}\|\mathbf{y}\|_2 \leq \|\mathbf{y}\|_\infty \leq \|\mathbf{y}\|_2$. These are exactly the other two inequalities $\|\mathbf{y}\|_\infty \leq \|\mathbf{y}\|_2 \leq \sqrt{n}\|\mathbf{y}\|_\infty$ of [Proposition 15.1.2](#): that proposition is closed under duality, with the constants carried over unchanged.

Exercise C3: The gauge is the support function of the polar.

Let $K \subseteq \mathbb{R}^n$ be closed and convex, with $\mathbf{0}$ an interior point.

- (a) Prove that $p_K(\mathbf{x}) = h_{K^\circ}(\mathbf{x})$ for every $\mathbf{x}$. *Hint: for $t > 0$, rewrite “ $\mathbf{x} \in tK$ ” using [Theorem 17.4.5](#).*
- (b) Deduce [Theorem 17.4.8](#) again from (a), [Theorem 17.4.2](#) (b) and [Proposition 17.4.6](#) (d).

Solution.

- (a) The polar K° contains $\mathbf{0}$, so $h_{K^\circ}(\mathbf{x}) \geq \langle \mathbf{x}, \mathbf{0} \rangle = 0$. Let $t > 0$. By [Theorem 17.4.5](#), $K = K^{\circ\circ}$, so

$$\mathbf{x} \in tK \iff \mathbf{x}/t \in K^{\circ\circ} \iff \langle \mathbf{x}, \mathbf{y} \rangle \leq t \text{ for every } \mathbf{y} \in K^\circ \iff h_{K^\circ}(\mathbf{x}) \leq t.$$

So the admissible t in [Definition 17.4.1](#) are the $t > 0$ with $t \geq h_{K^\circ}(\mathbf{x})$, and their infimum is $h_{K^\circ}(\mathbf{x})$ (if $h_{K^\circ}(\mathbf{x}) = 0$, the admissible t are all $t > 0$, with infimum 0). In particular $h_{K^\circ}(\mathbf{x})$ is finite.

- (b) Let B be the ball of $\|\cdot\|$. By [Theorem 17.4.2](#) (b), B is closed and convex with $\mathbf{0}$ interior, and $\|\cdot\| = p_B$. By (a), $p_B = h_{B^\circ}$, and B° is the ball of $\|\cdot\|_*$ by [Proposition 17.4.6](#) (d). So $\|\mathbf{x}\| = \max\{\langle \mathbf{x}, \mathbf{y} \rangle : \|\mathbf{y}\|_* \leq 1\} = \|\mathbf{x}\|_{**}$, the maximum existing by [Definition 17.4.4](#) applied to $\|\cdot\|_*$.