

6.5 Row Operations and Multiplicativity

The previous section defined the determinant by the Leibniz formula and proved it is the only normalized alternating form. The formula settles every theoretical question in principle, but with $n!$ terms it is useless for computing, and it hides the two facts that make determinants worth having. The first is that $\det A \neq 0$ exactly when A is invertible. The second is that $\det(AB) = \det A \det B$. This section proves both. Along the way we get a fast algorithm, and we extend the determinant from matrices to operators.

6.5.1 Row operations

Row reduction is the workhorse of Chapter 2, and the three rules of a determinant say precisely how it reacts to each elementary row operation. Everything follows from Theorem 6.4.5, which lets us treat $\det$ as a function of the rows.

Theorem 6.5.1 — Row Operations and the Determinant

Let $A \in M_n(F)$ and let B be obtained from A by one operation.

- (a) **(Swap)** If B is obtained by $R_i \leftrightarrow R_j$ with $i \neq j$, then $\det B = -\det A$.
- (b) **(Scale)** If B is obtained by $R_i \rightarrow cR_i$ with $c \in F$, then $\det B = c \det A$.
- (c) **(Replace)** If B is obtained by $R_i \rightarrow R_i + cR_j$ with $i \neq j$ and $c \in F$, then $\det B = \det A$.
- (d) The same three statements hold for the corresponding operations on **columns**.
- (e) For the elementary matrices of Definition 2.4.1,

$$\det P_{ij} = -1, \quad \det D_i(c) = c, \quad \det(I_n + cE_{ij}) = 1 \quad (i \neq j),$$

and $\det(EA) = \det E \det A$ for every elementary matrix $E \in M_n(F)$.

Proof. By Theorem 6.4.5, $\det$ is an alternating n -linear function of the rows $\mathbf{r}_1, \dots, \mathbf{r}_n$ of A . We write $\det(\mathbf{r}_1, \dots, \mathbf{r}_n)$ for it in this proof.

- (b) This is linearity in row i .
- (c) By linearity in row i ,

$$\det B = \det(\dots, \mathbf{r}_i + c\mathbf{r}_j, \dots, \mathbf{r}_j, \dots) = \det(\dots, \mathbf{r}_i, \dots, \mathbf{r}_j, \dots) + c \det(\dots, \mathbf{r}_j, \dots, \mathbf{r}_j, \dots),$$

where the displayed arguments sit in positions i and j . The last determinant has two equal rows, so it is 0. Hence $\det B = \det A$.

- (d) This is the swap rule of Theorem 6.2.1, applied to the rows.
- (e) By Theorem 6.4.3, $\det$ is an alternating n -linear function of the columns, and the arguments for (a), (b), (c) apply word for word.
- (f) The matrices P_{ij} , $D_i(c)$ and $I_n + cE_{ij}$ are obtained from I_n by a swap, a scaling and a replacement respectively. By (a), (b), (c) and $\det I_n = 1$, their determinants are -1 , c and 1 . Now let $E = \rho(I_n)$ be elementary. By Theorem 2.4.1, $EA = \rho(A)$. By (a), (b) or (c), $\det \rho(A)$ is $\det A$ times -1 , c or 1 respectively, and this factor

is $\det E$. Hence $\det(EA) = \det E \det A$. This proves the theorem. ■

Read part (e) as a bookkeeping rule. A swap flips the sign, a scaling by c multiplies by c , and a replacement changes nothing. Replacement, the operation used most in elimination, is free.

Warning. A combined move is not a replacement. The move $R_2 \rightarrow 3R_2 - R_1$ looks harmless but multiplies the determinant by 3: it is the scaling $R_2 \rightarrow 3R_2$ followed by the replacement $R_2 \rightarrow R_2 - R_1$. For $A = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}$, with $\det A = 1$, it produces $\begin{pmatrix} 1 & 2 \\ 2 & 7 \end{pmatrix}$, whose determinant is 3. Keep replacements in the form “row i plus a multiple of another row”, with coefficient 1 on row i .

Since elimination brings any square matrix to an upper triangular one, and triangular determinants are products of diagonal entries (Theorem 6.4.6), we have an algorithm.

Example. (A 4×4 Determinant by Elimination)

Compute $\det A$ for

$$A = \begin{pmatrix} 0 & 2 & 4 & 2 \\ 1 & 1 & 0 & 3 \\ 2 & 3 & 1 & 5 \\ -1 & 2 & 3 & 1 \end{pmatrix} \in M_4(\mathbb{R}).$$

Solution. We record each operation and its effect, using [Theorem 6.5.1](#).

Step 1. The $(1, 1)$ -entry is 0, so swap: $R_1 \leftrightarrow R_2$. This multiplies the determinant by -1 :

$$\det A = -\det \begin{pmatrix} 1 & 1 & 0 & 3 \\ 0 & 2 & 4 & 2 \\ 2 & 3 & 1 & 5 \\ -1 & 2 & 3 & 1 \end{pmatrix}.$$

Step 2. Clear column 1 with the replacements $R_3 \rightarrow R_3 - 2R_1$ and $R_4 \rightarrow R_4 + R_1$, which change nothing. Row 2 has a common factor 2; pulling it out is linearity in row 2:

$$\det A = -\det \begin{pmatrix} 1 & 1 & 0 & 3 \\ 0 & 2 & 4 & 2 \\ 0 & 1 & 1 & -1 \\ 0 & 3 & 3 & 4 \end{pmatrix} = -2 \det \begin{pmatrix} 1 & 1 & 0 & 3 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 1 & -1 \\ 0 & 3 & 3 & 4 \end{pmatrix}.$$

Step 3. Clear column 2: $R_3 \rightarrow R_3 - R_2$ and $R_4 \rightarrow R_4 - 3R_2$. Then clear column 3: $R_4 \rightarrow R_4 - 3R_3$. None of these changes the determinant:

$$\det A = -2 \det \begin{pmatrix} 1 & 1 & 0 & 3 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -1 & -2 \\ 0 & 0 & -3 & 1 \end{pmatrix} = -2 \det \begin{pmatrix} 1 & 1 & 0 & 3 \\ 0 & 1 & 2 & 1 \\ 0 & 0 & -1 & -2 \\ 0 & 0 & 0 & 7 \end{pmatrix}.$$

Step 4. The last matrix is upper triangular, so by Theorem 6.4.6 its determinant is

$$1 \cdot 1 \cdot (-1) \cdot 7 = -7. \text{ Hence } \det A = -2 \cdot (-7) = 14.$$

Elimination costs roughly $n^3/3$ arithmetic operations, and reading off the triangular determinant takes only $n - 1$ more multiplications, against $n!(n - 1)$ multiplications for the Leibniz formula (Exercise (C2: Counting the cost)). For $n = 20$ that is a few thousand operations against about 4.6×10^{19} . Every practical determinant computation is some form of elimination.

6.5.2 Determinant and invertibility

The same bookkeeping answers the invertibility question. Row reduction multiplies the determinant by non-zero factors only, so it cannot turn a non-zero determinant into zero or back. And the end of the reduction is either I_n or a matrix with a zero row.

Theorem 6.5.2 — Invertible if and only if Non-Zero Determinant

A matrix $A \in M_n(F)$ is invertible if and only if $\det A \neq 0$.

Proof. Let R be the reduced row echelon form of A (Theorem 2.2.3, Theorem 2.3.4). By Definition 2.2.2 and Theorem 2.4.1, there are elementary matrices $E_1, \dots, E_k$ (possibly $k = 0$) with $R = E_k \cdots E_1 A$. Applying Theorem 6.5.1 (e) k times,

$$\det R = \det E_k \cdots \det E_1 \det A.$$

Each $\det E_m$ is -1 , 1 , or some $c \neq 0$, since a scaling operation requires $c \neq 0$. A product of non-zero elements of a field is non-zero (Theorem 0.7.1), so $\det R = 0$ if and only if $\det A = 0$.

By Lemma 2.4.3, either $R = I_n$, and then $\det R = 1 \neq 0$; or the last row of R is zero. In the second case, scaling that row by 0 leaves R unchanged, so $\det R = 0 \cdot \det R = 0$ by Theorem 6.5.1 (b). Hence $\det A \neq 0$ if and only if $R = I_n$, which by Theorem 2.4.4 holds if and only if A is invertible. ■

This is the promised addition to the list of equivalent conditions. We collect it with the conditions from Chapter 3 about the map $T_A: \mathbf{x} \mapsto A\mathbf{x}$, and with two conditions about rows that the determinant makes easy.

Theorem 6.5.3 — Invertible Matrix Theorem, with Determinants

Let $A \in M_n(F)$ and $T_A: F^n \rightarrow F^n$, $\mathbf{x} \mapsto A\mathbf{x}$. The following are equivalent, and each is equivalent to each of the eight conditions of Theorem 2.4.4.

- (a) A is invertible.
- (b) T_A is injective.
- (c) T_A is surjective.
- (d) $\text{rank } A = n$.

- (e) A^T is invertible.
- (f) The rows of A are linearly independent.
- (g) $\det A \neq 0$.

Proof. Item (a) is item (a) of Theorem 2.4.4, so it suffices to connect each new item with that theorem.

(a) $\Leftrightarrow$ (g). This is Theorem 6.5.2.

(a) $\Leftrightarrow$ (b). By Theorem 3.5.7 ((b) $\Leftrightarrow$ (c)) with $V = F^n$, T_A is injective if and only if $\ker T_A = \{\mathbf{x} : A\mathbf{x} = \mathbf{0}\}$ is $\{\mathbf{0}\}$. That is item (b) of Theorem 2.4.4.

(b) $\Leftrightarrow$ (c). This is Theorem 3.5.7 ((b) $\Leftrightarrow$ (d)), since F^n is finite-dimensional.

(c) $\Leftrightarrow$ (d). By Theorem 0.9.3, $\text{im } T_A$ is the set of combinations of the columns of A , that is, $\text{col}(A)$. So T_A is surjective if and only if $\text{col}(A) = F^n$. Since $\text{col}(A)$ is a subspace of F^n , this holds if and only if $\text{rank } A = \dim \text{col}(A) = n$ (Definition 2.5.4, Theorem 1.6.8).

(a) $\Leftrightarrow$ (e). By Theorem 6.4.5, $\det A^T = \det A$. Applying (a) $\Leftrightarrow$ (g) to A and to A^T , both “ A invertible” and “ A^T invertible” are equivalent to $\det A \neq 0$.

(e) $\Leftrightarrow$ (f). Let $\mathbf{r}_1, \dots, \mathbf{r}_n$ be the rows of A . The columns of A^T are $\mathbf{r}_1^T, \dots, \mathbf{r}_n^T$. By Theorem 0.9.5, $c_1\mathbf{r}_1 + \dots + c_n\mathbf{r}_n = \mathbf{0}$ if and only if $c_1\mathbf{r}_1^T + \dots + c_n\mathbf{r}_n^T = \mathbf{0}$, so the rows of A are independent exactly when the columns of A^T are. By Theorem 2.4.4 ((a) $\Leftrightarrow$ (f)) applied to A^T , this holds exactly when A^T is invertible. This proves the theorem. ■

The determinant is now one more item on the menu, and for the right problems it is the best one. For a concrete matrix with a parameter, one polynomial in the parameter decides invertibility at once. For an abstract matrix, such as $A + I$ when $A^k = 0$, the determinant is usually hopeless, and the kernel is the item to try first.

Quick check. For which $t \in \mathbb{R}$ is $\begin{pmatrix} 1 & t \\ t & 4 \end{pmatrix}$ invertible? Over $\mathbb{F}_3$, for which t ?

Solution. The determinant is $4 - t^2 = (2 - t)(2 + t)$. Over $\mathbb{R}$ the matrix is invertible exactly for $t \neq \pm 2$. Over $\mathbb{F}_3$, $4 = 1$, so the determinant is $1 - t^2$, which is 1 for $t = 0$ and 0 for $t = 1, 2$; the matrix is invertible only for $t = 0$.

6.5.3 Multiplicativity

Now the second fact. A natural first attempt is to multiply out the Leibniz formula for AB , whose entries are sums, and regroup. That is legitimate, but the bookkeeping is horrible. The uniqueness theorem does the work instead: we exhibit $B \mapsto \det(AB)$ as an alternating form in the columns of B , and then it has no choice.

Theorem 6.5.4 — Multiplicativity of the Determinant

For all $A, B \in M_n(F)$,

$$\det(AB) = \det A \det B.$$

Idea. Fix A . The j -th column of AB is $A\mathbf{b}_j$, so $g(\mathbf{b}_1, \dots, \mathbf{b}_n) = \det(A\mathbf{b}_1, \dots, A\mathbf{b}_n)$ is $\det(AB)$ viewed as a function of the columns of B . It inherits linearity in each $\mathbf{b}_j$ from $\det$ and from matrix multiplication, and equal columns $\mathbf{b}_p = \mathbf{b}_q$ give equal columns $A\mathbf{b}_p = A\mathbf{b}_q$. So g is an alternating form, hence a multiple of $\det$, and the multiple is $g(\mathbf{e}_1, \dots, \mathbf{e}_n) = \det A$.

Proof. Fix $A \in M_n(F)$, and define $g: F^n \times \dots \times F^n \rightarrow F$ by

$$g(\mathbf{b}_1, \dots, \mathbf{b}_n) = \det(A\mathbf{b}_1, \dots, A\mathbf{b}_n).$$

By Theorem 0.9.2 (columns), if B has columns $\mathbf{b}_1, \dots, \mathbf{b}_n$, then AB has columns $A\mathbf{b}_1, \dots, A\mathbf{b}_n$, so $g(\mathbf{b}_1, \dots, \mathbf{b}_n) = \det(AB)$.

g is n -linear. Fix k and all $\mathbf{b}_j$ with $j \neq k$. The map $\mathbf{b}_k \mapsto A\mathbf{b}_k$ is linear by Theorem 0.9.4, and $\det$ is linear in its k -th argument by Theorem 6.4.3. A composite of linear maps is linear, so g is linear in $\mathbf{b}_k$.

g is alternating. If $\mathbf{b}_p = \mathbf{b}_q$ with $p \neq q$, then $A\mathbf{b}_p = A\mathbf{b}_q$, so $g(\mathbf{b}_1, \dots, \mathbf{b}_n) = 0$ since $\det$ is alternating.

By Corollary 6.4.4, $g(\mathbf{b}_1, \dots, \mathbf{b}_n) = g(\mathbf{e}_1, \dots, \mathbf{e}_n) \det(\mathbf{b}_1, \dots, \mathbf{b}_n)$. Finally $A\mathbf{e}_j$ is the j -th column of A (Theorem 0.9.3), so $g(\mathbf{e}_1, \dots, \mathbf{e}_n) = \det A$. Hence $\det(AB) = \det A \det B$. ■

Remark. Over a commutative ring. This proof uses only Corollary 6.4.4 (in the explicit form of Theorem 6.4.2), linearity of matrix multiplication, and the Leibniz formula. None of these divides. So $\det(AB) = \det A \det B$ holds for $A, B \in M_n(R)$ over any commutative ring R , such as $F[x]$.

Alternatively, the building blocks of Chapter 2 give a proof. If A is invertible, then $A = E_1 \cdots E_k$ with elementary E_m (Theorem 2.4.4 (d)), and peeling off one factor at a time with Theorem 6.5.1 (e) gives $\det(AB) = \det E_1 \cdots \det E_k \det B$; the case $B = I_n$ of the same computation shows $\det A = \det E_1 \cdots \det E_k$. If A is not invertible, neither is AB : otherwise $(AB)C = I_n$ for some C , so $A(BC) = I_n$ and A would be invertible by Theorem 2.4.5. Then both sides are 0 by Theorem 6.5.2. This route leans on row reduction, which divides, so unlike the first proof it needs a field.

For reference, here are the two identities used most often, now both proved.

Theorem 6.5.5 — Properties of Determinants

Let $A, B \in M_n(F)$. Then:

1. $\det(AB) = \det A \det B$;
2. $\det A^T = \det A$.

Proof. Part 1 is [Theorem 6.5.4](#) and part 2 is [Theorem 6.4.5](#). ■

The first consequences are immediate.

Corollary 6.5.6 — Determinant of the Inverse

If $A \in M_n(F)$ is invertible, then $\det A \neq 0$ and $\det(A^{-1}) = (\det A)^{-1}$.

Proof. By [Theorem 6.5.4](#), $\det A \det(A^{-1}) = \det(AA^{-1}) = \det I_n = 1$. Hence $\det A \neq 0$, and $\det(A^{-1})$ is the inverse of $\det A$ in F . ■

Warning. **Multiplicativity does not make det additive, and it does not make products commute.** For $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 0 & 1 \\ 1 & 5 \end{pmatrix}$, $AB = \begin{pmatrix} 2 & 11 \\ 4 & 23 \end{pmatrix} \neq \begin{pmatrix} 3 & 4 \\ 16 & 22 \end{pmatrix} = BA$, yet $\det(AB) = \det(BA) = 2$, since both equal $\det A \det B = (-2)(-1)$. On the other hand, $\det(A+B)$ is not determined by $\det A$ and $\det B$ at all: I_2 and I_2 have determinants 1, 1 and $\det(I_2 + I_2) = 4$, while I_2 and $C = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ also have determinants 1, 1 but $\det(I_2 + C) = \det \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} = 2$.

Quick check. Let $A = \begin{pmatrix} 2 & 1 \\ 7 & 4 \end{pmatrix}$. Without multiplying any matrices, find $\det(A^5)$, $\det(A^{-1})$ and $\det(A^T A)$.

Solution. $\det A = 8 - 7 = 1$. By [Theorem 6.5.4](#), $\det(A^5) = (\det A)^5 = 1$. By [Corollary 6.5.6](#), $\det(A^{-1}) = 1^{-1} = 1$. By [Theorem 6.5.5](#), $\det(A^T A) = \det A^T \det A = (\det A)^2 = 1$.

6.5.4 The determinant of an operator

In Chapter 3 the trace passed from matrices to operators because similar matrices have the same trace ([Definition 3.10.2](#)). Multiplicativity gives the same for determinants.

Corollary 6.5.7 — Determinant Is a Similarity Invariant

If $A, B \in M_n(F)$ are similar, then $\det A = \det B$.

Proof. Let $B = P^{-1}AP$ with P invertible. By [Theorem 6.5.4](#) and [Corollary 6.5.6](#),

$$\det B = \det(P^{-1}) \det A \det P = (\det P)^{-1} \det P \det A = \det A,$$

where the middle step uses commutativity of multiplication in F .

The converse fails: I_2 and $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ both have determinant 1, but the only matrix similar to I_2 is $P^{-1}I_2P = I_2$ itself.

By Theorem 3.7.4, all matrices of one operator in the various bases are similar to each other. So the corollary lets us define:

Definition 6.5.1 — Determinant of an operator

Let V be a finite-dimensional vector space over F with $\dim V \geq 1$, and let $T \in \mathcal{L}(V)$. The **determinant** of T is

$$\det T := \det[T]_{\mathcal{B}},$$

where $\mathcal{B}$ is **any** basis of V . For $V = \{\mathbf{0}\}$ we set $\det T = 1$.

Well-definedness. If $\mathcal{B}$ and $\mathcal{C}$ are bases of V , then $[T]_{\mathcal{C}} = P^{-1}[T]_{\mathcal{B}}P$ with $P = [\text{id}]_{\mathcal{C}}^{\mathcal{B}}$ invertible, by Theorem 3.7.3 and Theorem 3.7.4 (a). By Corollary 6.5.7 the two determinants agree. The convention for $V = \{\mathbf{0}\}$ is the empty product, and it keeps the next theorem true in that case.

Examples.

- **Scalars.** For $c \in F$ and $\dim V = n \geq 1$, $[c \text{id}_V]_{\mathcal{B}} = cI_n$ in every basis, so $\det(c \text{id}_V) = c^n$ by Theorem 6.4.6.
- **A reflection.** $R(x, y) = (y, x)$ on $\mathbb{R}^2$ has standard matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, of determinant -1 . In the basis $((1, 1), (1, -1))$ its matrix is $\text{diag}(1, -1)$ (Example (The reflection in a mirror-adapted basis)), again of determinant -1 , as it must be.
- **Differentiation.** On $\mathbb{R}[x]_{\leq 3}$ with basis $(1, x, x^2, x^3)$, the matrix of D is upper triangular with zero diagonal (Example (Differentiation on Cubic Polynomials)), so $\det D = 0$.

The determinant of a map $T: V \rightarrow W$ between **different** spaces is not defined, even when $\dim V = \dim W$. A matrix $[T]_{\mathcal{B}}^{\mathcal{C}}$ exists, but changing $\mathcal{C}$ alone multiplies it on the left by an arbitrary invertible matrix, which can change its determinant to any non-zero value. For $\text{id}: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ with input basis $\mathcal{E}$ and output basis $((2, 0), (0, 1))$, the matrix is $\text{diag}(\frac{1}{2}, 1)$, of determinant $\frac{1}{2}$, not 1.

Theorem 6.5.8 — Properties of the Determinant of an Operator

Let V be a finite-dimensional vector space over F and $S, T \in \mathcal{L}(V)$. Then

$$\det(ST) = \det S \det T, \quad \det \text{id}_V = 1,$$

and T is invertible if and only if $\det T \neq 0$.

Proof. If $V = \{\mathbf{0}\}$, all determinants are 1 and the only operator is id_V , which is invertible. Otherwise fix a basis $\mathcal{B}$ of V . By Theorem 3.6.2, $[ST]_{\mathcal{B}} = [S]_{\mathcal{B}}[T]_{\mathcal{B}}$, so $\det(ST) = \det S \det T$ by Theorem 6.5.4. Next $[\text{id}_V]_{\mathcal{B}} = I_n$, of determinant 1. Finally, by Theorem 3.6.5 (b), T is invertible if and only if $[T]_{\mathcal{B}}$ is, which by Theorem 6.5.2 holds if and only if $\det T = \det[T]_{\mathcal{B}} \neq 0$.

So the determinant is a property of the operator, not of the coordinates. Differentiation on $\mathbb{R}[x]_{\leq 3}$ is not invertible because its determinant is 0, whichever basis we compute in. In the section on orientation and volume we will see what the number $\det T$ measures geometrically.

6.5.5 The determinant as a homomorphism

In Chapter 0 we met groups and homomorphisms, and promised that the determinant would be one. The general linear group $\mathrm{GL}_n(F)$ is the group of invertible matrices in $M_n(F)$ under multiplication, and $F \setminus \{0\}$ is a group under multiplication (Example (Groups)).

Theorem 6.5.9 — The Determinant Is a Group Homomorphism

Let $n \geq 1$. The determinant restricts to a **surjective** group homomorphism

$$\det: \mathrm{GL}_n(F) \rightarrow F \setminus \{0\}.$$

Its kernel is the **special linear group** $\mathrm{SL}_n(F) := \{A \in M_n(F) : \det A = 1\}$, which is therefore a subgroup of $\mathrm{GL}_n(F)$.

Proof. By Theorem 6.5.2, $\det A \neq 0$ for $A \in \mathrm{GL}_n(F)$, so $\det$ maps $\mathrm{GL}_n(F)$ into $F \setminus \{0\}$, and Theorem 6.5.4 says it is a homomorphism (Definition 0.10.3). For $c \neq 0$, the matrix $\mathrm{diag}(c, 1, \dots, 1)$ has determinant c by Theorem 6.4.6, which is non-zero, so it lies in $\mathrm{GL}_n(F)$; hence $\det$ is surjective. A matrix with determinant 1 is invertible by Theorem 6.5.2, so the kernel $\{A \in \mathrm{GL}_n(F) : \det A = 1\}$ is $\mathrm{SL}_n(F)$. It is a subgroup by Theorem 0.10.3.

For example, $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and the elementary matrices $I_n + cE_{ij}$ lie in $\mathrm{SL}_n(F)$, while P_{12} , of determinant -1 , does not. Over $\mathbb{R}$, the rotation matrices $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ have determinant $\cos^2 \theta + \sin^2 \theta = 1$ and lie in $\mathrm{SL}_2(\mathbb{R})$. The sign $\mathrm{sgn}: S_n \rightarrow \{1, -1\}$ of the third section is the other homomorphism promised in Chapter 0, and the two are linked: $\det P_\sigma = \mathrm{sgn}(\sigma)$ (Exercise (B2: The determinant of P_σ)), and $P_\sigma P_\tau = P_{\sigma\tau}$ (Lemma 2.6.6).

6.5.6 Which proofs survive over a ring

Remark. Commutative rings. Over a commutative ring R (as in the remark at the end of the previous section), the following proofs of this section use no division and remain valid for matrices in $M_n(R)$: Theorem 6.5.1, with $c \in R$ arbitrary in (b) and (c); Theorem 6.5.4, by its first proof; and Theorem 6.5.5. The proofs of Corollary 6.5.6 and Corollary 6.5.7 also survive for matrices that have an inverse in $M_n(R)$ (then $\det A \det(A^{-1}) = 1$, so $\det A$ has an inverse in R), for instance an invertible $P \in M_n(F)$ regarded inside $M_n(F[x])$; this is the case needed for characteristic polynomials. What fails over a ring is the converse direction: a matrix whose determinant is

non-zero need not be invertible, since $\det$ may be a non-zero element with no inverse, such as $2 \in \mathbb{Z}$ or $x \in F[x]$. By contrast, [Theorem 6.5.2](#), [Theorem 6.5.3](#) and the “Alternatively” proof of multiplicativity run through row reduction, which divides by pivots, and they are proved here only over a field.

6.5.7 Exercises

6.5.7.1 A. Check your understanding

Exercise A1.

- (a) State how each of the three elementary row operations changes the determinant.
- (b) True or false: the move $R_2 \rightarrow 3R_2 - R_1$ does not change the determinant. Justify your answer.
- (c) True or false: if $A, B \in M_n(F)$ have $\det A = \det B$, then A and B are similar. Justify your answer.
- (d) True or false: $\det(AB) = \det(BA)$ for all $A, B \in M_n(F)$. Justify your answer.
- (e) Explain why $\det T$ does not depend on the basis used to compute it.
- (f) Is $\det: M_n(F) \rightarrow F$ a group homomorphism? Explain.

Solution.

- (a) By [Theorem 6.5.1](#): a swap multiplies it by -1 , scaling a row by c multiplies it by c , and adding a multiple of one row to a different row leaves it unchanged.
- (b) False. The move is $R_2 \rightarrow 3R_2$ followed by $R_2 \rightarrow R_2 - R_1$, so it multiplies the determinant by 3. For I_2 it gives $\begin{pmatrix} 1 & 0 \\ -1 & 3 \end{pmatrix}$, of determinant $3 \neq 1$.
- (c) False. I_2 and $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ both have determinant 1, but $P^{-1}I_2P = I_2$ for every invertible P , so I_2 is similar only to itself.
- (d) True. Both equal $\det A \det B$ by [Theorem 6.5.4](#), since multiplication in F is commutative.
- (e) Matrices of T in two bases are similar ([Theorem 3.7.3](#)), and similar matrices have equal determinants ([Corollary 6.5.7](#)).
- (f) No: $M_n(F)$ is not a group under multiplication, since the zero matrix has no inverse. The restriction $\det: \mathrm{GL}_n(F) \rightarrow F \setminus \{0\}$ is a homomorphism ([Theorem 6.5.9](#)).

6.5.7.2 B. Practice

Exercise B1: Determinants by elimination.

Compute the determinants of the following real matrices by row reduction, recording

the effect of each operation.

$$A = \begin{pmatrix} 1 & 2 & 0 & 1 \\ 2 & 3 & 1 & 4 \\ 0 & 1 & 3 & 2 \\ -1 & 1 & 2 & 5 \end{pmatrix}, \quad B = \begin{pmatrix} 2 & -1 & 3 & 1 \\ 4 & -2 & 7 & 3 \\ -2 & 2 & -1 & 4 \\ 6 & -3 & 9 & 5 \end{pmatrix}.$$

Solution. For A : the replacements $R_2 \rightarrow R_2 - 2R_1$ and $R_4 \rightarrow R_4 + R_1$ give rows $(1, 2, 0, 1)$, $(0, -1, 1, 2)$, $(0, 1, 3, 2)$, $(0, 3, 2, 6)$. Then $R_3 \rightarrow R_3 + R_2$ and $R_4 \rightarrow R_4 + 3R_2$ give $(0, 0, 4, 4)$ and $(0, 0, 5, 12)$. Pulling the factor 4 out of row 3 leaves $(0, 0, 1, 1)$, and $R_4 \rightarrow R_4 - 5R_3$ gives $(0, 0, 0, 7)$. None of the replacements changes the determinant (Theorem 6.5.1), so

$$\det A = 4 \det \begin{pmatrix} 1 & 2 & 0 & 1 \\ 0 & -1 & 1 & 2 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 7 \end{pmatrix} = 4 \cdot (1 \cdot (-1) \cdot 1 \cdot 7) = -28,$$

by Theorem 6.4.6.

For B : $R_2 \rightarrow R_2 - 2R_1$, $R_3 \rightarrow R_3 + R_1$, $R_4 \rightarrow R_4 - 3R_1$ give rows $(2, -1, 3, 1)$, $(0, 0, 1, 1)$, $(0, 1, 2, 5)$, $(0, 0, 0, 2)$, without changing the determinant. The $(2, 2)$ -entry is 0, so swap $R_2 \leftrightarrow R_3$, which multiplies the determinant by -1 . The result is upper triangular with diagonal $2, 1, 1, 2$, so

$$\det B = -(2 \cdot 1 \cdot 1 \cdot 2) = -4.$$

Exercise B2: A parameter.

Let $t \in \mathbb{R}$ and $A_t = \begin{pmatrix} t & 1 & 1 \\ 1 & t & 1 \\ 1 & 1 & t \end{pmatrix}$. Find $\det A_t$ as a factored polynomial in t . Hence determine for which t the matrix A_t is invertible.

Solution. The replacements $R_1 \rightarrow R_1 + R_2$ and $R_1 \rightarrow R_1 + R_3$ do not change the determinant and make row 1 equal to $(t+2, t+2, t+2)$. By linearity in row 1 (this holds for every value of $t+2$, including 0),

$$\det A_t = (t+2) \det \begin{pmatrix} 1 & 1 & 1 \\ 1 & t & 1 \\ 1 & 1 & t \end{pmatrix} = (t+2) \det \begin{pmatrix} 1 & 1 & 1 \\ 0 & t-1 & 0 \\ 0 & 0 & t-1 \end{pmatrix} = (t+2)(t-1)^2,$$

where the second step uses $R_2 \rightarrow R_2 - R_1$, $R_3 \rightarrow R_3 - R_1$, and the last uses Theorem 6.4.6. By Theorem 6.5.2, A_t is invertible if and only if $t \neq -2$ and $t \neq 1$.

Exercise B3: The shift operator.

Let $T: \mathbb{R}[x]_{\leq 3} \rightarrow \mathbb{R}[x]_{\leq 3}$, $T(p) = p(x+1)$. Find $\det T$. Hence show that T is invertible.

Solution. In the basis $\mathcal{B} = (1, x, x^2, x^3)$, $T(1) = 1$, $T(x) = 1 + x$, $T(x^2) = 1 + 2x + x^2$

and $T(x^3) = 1 + 3x + 3x^2 + x^3$. So

$$[T]_{\mathcal{B}} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

which is upper triangular with ones on the diagonal. By [Definition 6.5.1](#) and [Theorem 6.4.6](#), $\det T = 1$. Since $\det T \neq 0$, T is invertible by [Theorem 6.5.8](#). (Its inverse is $p \mapsto p(x-1)$.)

6.5.7.3 C. Going deeper

Exercise C1: Orthogonal-type matrices.

- (a) Let $A \in M_n(\mathbb{R})$ satisfy $A^T A = I_n$. Prove that $\det A = 1$ or $\det A = -1$.
- (b) Show that both values occur, for every $n \geq 1$.
- (c) Over $\mathbb{F}_2$, what can $\det A$ be if $A^T A = I_n$?

Solution.

- (a) By [Theorem 6.5.5](#), $1 = \det I_n = \det(A^T A) = \det A^T \det A = (\det A)^2$. A real number with square 1 is 1 or -1 .
- (b) $I_n^T I_n = I_n$ and $\det I_n = 1$. For $D = \text{diag}(-1, 1, \dots, 1)$, $D^T D = D^2 = I_n$ and $\det D = -1$ by [Theorem 6.4.6](#).
- (c) The same computation gives $(\det A)^2 = 1$. In $\mathbb{F}_2$, $\det A \in \{0, 1\}$ and $0^2 = 0$, so $\det A = 1$. This agrees with (a), since $-1 = 1$ in $\mathbb{F}_2$.

Exercise C2: Left multiplication on matrices.

Let $A \in M_n(F)$, and let $L_A: M_n(F) \rightarrow M_n(F)$, $L_A(X) = AX$. This is a linear operator on a space of dimension n^2 . Prove that

$$\det L_A = (\det A)^n.$$

- (a) Order the matrix units column by column, $\mathcal{B} = (E_{11}, E_{21}, \dots, E_{n1}, E_{12}, \dots, E_{n2}, \dots, E_{1n}, \dots, E_{nn})$. Show that $[L_A]_{\mathcal{B}}$ consists of n copies of A placed along the diagonal, with zeros elsewhere.
- (b) Show that $L_{AB} = L_A L_B$, and prove the formula when A is an elementary matrix.
- (c) Prove the formula for invertible A , and then for non-invertible A .

Hint: for a swap matrix, identify $[L_{P_{ij}}]_{\mathcal{B}}$ as a permutation matrix and use Exercise (B2: The determinant of P_{σ}).

Solution.

- (a) By [Theorem 0.9.2](#), the l -th column of AE_{kl} is $A\mathbf{e}_k$, the k -th column of A , and the other columns are zero. So $AE_{kl} = \sum_i a_{ik} E_{il}$. In $\mathcal{B}$, the coordinates of this vector are zero outside the l -th block of n positions, and inside that block they are the

k -th column of A . Hence the columns of $[L_A]_{\mathcal{B}}$ belonging to the l -th block are those of A , placed in the rows of the l -th block: $[L_A]_{\mathcal{B}}$ has n copies of A along the diagonal.

- (b) $L_{AB}(X) = ABX = L_A(L_B(X))$ by associativity. Now let E be elementary. If $E = I_n + cE_{ij}$ with $i \neq j$, then by (a) $[L_E]_{\mathcal{B}}$ is triangular (upper if $i < j$, lower if $i > j$) with all diagonal entries 1, so $\det L_E = 1 = (\det E)^n$ by Theorem 6.4.6 and Theorem 6.5.1 (e). If $E = D_i(c)$, then $[L_E]_{\mathcal{B}}$ is diagonal with c appearing n times and 1 elsewhere, so $\det L_E = c^n = (\det E)^n$. If $E = P_{ij}$, then $[L_E]_{\mathcal{B}}$ has exactly one 1 in each row and column, so it is a permutation matrix P_{σ} (Definition 2.6.2), and σ exchanges the positions of E_{il} and E_{jl} for each l : σ is a product of n disjoint transpositions. By Exercise (B2: The determinant of P_{σ}), Theorem 6.3.1 and Corollary 6.3.2, $\det L_E = \operatorname{sgn}(\sigma) = (-1)^n = (\det E)^n$.
- (c) If A is invertible, then $A = E_1 \cdots E_k$ with elementary E_m (Theorem 2.4.4 (d)). By (b) and Theorem 6.5.8, $\det L_A = \det L_{E_1} \cdots \det L_{E_k} = (\det E_1)^n \cdots (\det E_k)^n = (\det E_1 \cdots \det E_k)^n = (\det A)^n$, where the last step is Theorem 6.5.4. If A is not invertible, there is $\mathbf{v} \neq \mathbf{0}$ with $A\mathbf{v} = \mathbf{0}$ (Theorem 2.4.4 (b)). The matrix $X = (\mathbf{v} \ \mathbf{0} \ \cdots \ \mathbf{0})$ is non-zero and $AX = \mathbf{0}$, so L_A is not injective, hence not invertible, and $\det L_A = 0$ by Theorem 6.5.8. Also $\det A = 0$ by Theorem 6.5.2, so $(\det A)^n = 0$ as $n \geq 1$.

Exercise C3: Non-square factors.

- (a) Give $A \in M_{2 \times 1}(\mathbb{R})$ and $B \in M_{1 \times 2}(\mathbb{R})$ with $\det(BA) \neq \det(AB)$.
- (b) Let $m > n$, $A \in M_{m \times n}(F)$ and $B \in M_{n \times m}(F)$. Prove that $\det(AB) = 0$.
- (c) Explain why (a) does not contradict Theorem 6.5.4.

Solution.

- (a) Let $A = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $B = \begin{pmatrix} 1 & 0 \end{pmatrix}$. Then $BA = (1)$, of determinant 1, and $AB = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, of determinant 0.
- (b) By Theorem 2.5.9, $\operatorname{rank}(AB) \leq \operatorname{rank} A \leq n$, since $\operatorname{col}(A)$ is spanned by n columns. So $\operatorname{rank}(AB) < m$, and the $m \times m$ matrix AB is not invertible by Theorem 6.5.3 (d). Hence $\det(AB) = 0$ by Theorem 6.5.2.
- (c) Theorem 6.5.4 concerns two **square** matrices of the same size, and then $\det(AB) = \det(BA)$. Here A and B are not square, $\det A$ and $\det B$ are not defined, and AB , BA have different sizes.