

15.6 Bounds for the Exponential, and Two Derivatives

Chapter 9 §09 built $e^{\mathbf{A}}$ out of a series and closed with a promise: “Chapter 15 will return to $e^{\mathbf{A}}$ with norms available, which give quantitative bounds instead of exact formulas.” That is what this section does. It does not rebuild the exponential; Definition 9.9.2 and Theorem 9.9.3 stand as they are, and the Jordan form is not touched. What is new is that the series now has a size, so its sum has a ceiling, its truncation has an error bound, and its behavior as $t \rightarrow \infty$ has a rate.

The second half takes the other step Chapter 9 could not: it differentiates. The derivative of the determinant and the derivative of the inverse are both needed later in the book, and both require one fact from calculus that linear algebra does not supply. That fact is stated before it is used.

Throughout, $\|\cdot\|$ is the operator norm on $M_n(\mathbb{C})$ induced by a norm on $\mathbb{C}^n$ (Definition 15.3.2), so $\|\mathbf{I}\| = 1$ and $\|\mathbf{XY}\| \leq \|\mathbf{X}\|\|\mathbf{Y}\|$ (Theorem 15.3.2 (c), (d)).

15.6.1 How big is the exponential

Chapter 9 §10’s criteria answered *whether*; the norm answers *how much*. The bound to aim at is the one the scalar case suggests: $|e^z| \leq e^{|z|}$, because the series has non-negative terms once every term is replaced by its modulus. The matrix proof is the same sentence with submultiplicativity in place of $|zw| = |z||w|$.

One scalar fact is needed and is not on the chapter’s list of imports. We name it here rather than let it in quietly.

Remark. A quoted fact. For every real $s \geq 0$ the series $\sum_{m \geq 0} s^m/m!$ converges, with sum e^s . This is fact (A1) of Chapter 9 §09, quoted there and quoted again here; this book does not prove it. Because its terms are non-negative, its partial sums increase to e^s , so every partial sum is at most e^s — that last step is **(A2), monotone convergence**, from the list in this chapter’s introduction.

Theorem 15.6.1. Bounds for the Exponential

Let $\mathbf{A} \in M_n(\mathbb{C})$ and write $a = \|\mathbf{A}\|$. Then

- (a) $\|e^{\mathbf{A}}\| \leq e^a$;
- (b) $\|e^{\mathbf{A}} - \mathbf{I}\| \leq e^a - 1$;
- (c) $\|e^{\mathbf{A}} - \mathbf{I} - \mathbf{A}\| \leq e^a - 1 - a \leq \frac{a^2 e^a}{2}$.

In particular $\|e^{t\mathbf{A}}\| \leq e^{|t|a}$ for every real t .

Idea. All three are the same computation stopped at a different place: bound the norm of a partial sum by the corresponding partial sum of $\sum s^m/m!$, then let the partial sums converge. The only thing that needs care is the passage to the limit, and Lemma 15.1.1 does it, exactly

as in Theorem 15.5.1. For the second inequality in (c), compare $m!$ with $2(m-2)!$.

Proof. Write $\mathbf{S}_M = \sum_{m=0}^M \mathbf{A}^m/m!$. By Theorem 9.9.3 the sequence $(\mathbf{S}_M)$ converges entrywise to $e^{\mathbf{A}}$, hence $\|\mathbf{S}_M - e^{\mathbf{A}}\| \rightarrow 0$ by Corollary 15.2.4, hence $\|\mathbf{S}_M\| \rightarrow \|e^{\mathbf{A}}\|$ by Lemma 15.1.1. The same three steps apply to $\mathbf{S}_M - \mathbf{I}$ and to $\mathbf{S}_M - \mathbf{I} - \mathbf{A}$. So in each part it suffices to bound the partial sums, uniformly in M .

- (a) Submultiplicativity gives $\|\mathbf{A}^m\| \leq a^m$ for $m \geq 1$, by induction, and $\|\mathbf{A}^0\| = \|\mathbf{I}\| = 1$. By subadditivity and the quoted fact above,

$$\|\mathbf{S}_M\| \leq \sum_{m=0}^M \frac{\|\mathbf{A}^m\|}{m!} \leq \sum_{m=0}^M \frac{a^m}{m!} \leq e^a.$$

Letting $M \rightarrow \infty$ gives $\|e^{\mathbf{A}}\| \leq e^a$.

- (b) The same estimate with the term $m = 0$ removed:

$$\|\mathbf{S}_M - \mathbf{I}\| \leq \sum_{m=1}^M \frac{a^m}{m!} \leq e^a - 1,$$

since the partial sums of $\sum_{m \geq 1} a^m/m!$ increase to $e^a - 1$.

- (c) Removing the terms $m = 0$ and $m = 1$ in the same way gives

$$\|e^{\mathbf{A}} - \mathbf{I} - \mathbf{A}\| \leq \sum_{m \geq 2} \frac{a^m}{m!} = e^a - 1 - a.$$

For the second inequality, note that $m! = m(m-1)(m-2)! \geq 2(m-2)!$ for every $m \geq 2$, so $a^m/m! \leq \frac{1}{2}a^2 \cdot a^{m-2}/(m-2)!$. Summing over $m \geq 2$ and substituting $l = m - 2$,

$$\sum_{m \geq 2} \frac{a^m}{m!} \leq \frac{a^2}{2} \sum_{l \geq 0} \frac{a^l}{l!} = \frac{a^2 e^a}{2}.$$

The final claim is (a) applied to $t\mathbf{A}$, whose norm is $|t|a$ by homogeneity. This proves the theorem. ■

Part (c) is the statement that $e^{\mathbf{A}}$ agrees with $\mathbf{I} + \mathbf{A}$ to second order in $\|\mathbf{A}\|$, with an explicit constant. It is the bound one uses when $\mathbf{A}$ is a small perturbation, and it is the counterpart, for the exponential, of Theorem 15.5.1 (c).

Example. Where the bound is exact, and where it is useless

Compare $\|e^{\mathbf{A}}\|_2$ with $e^{\|\mathbf{A}\|_2}$ for

- (a) $\mathbf{A} = \text{diag}(2, 1)$;
 (b) $\mathbf{A} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, and more generally $t\mathbf{A}$ for $t > 0$.

Solution.

- (a) $\mathbf{A}$ is Hermitian, hence normal, so its singular values are the moduli of its eigenvalues and $\|\mathbf{A}\|_2 = 2$ by Theorem 15.3.3 (c). By Definition 9.9.1 (or directly from the series, entry by entry) $e^{\mathbf{A}} = \text{diag}(e^2, e)$, again normal, so $\|e^{\mathbf{A}}\|_2 = e^2$. The bound $e^{\|\mathbf{A}\|_2} = e^2$ is attained exactly. The same happens for every $\mathbf{A} = c\mathbf{I}$ with $c > 0$, so no constant smaller than 1 can be inserted in Theorem 15.6.1 (a).
- (b) $\mathbf{A}^2 = -\mathbf{I}$, so $\mathbf{A}^3 = -\mathbf{A}$, $\mathbf{A}^4 = \mathbf{I}$, and grouping the series by parity gives, using fact (A1) of Chapter 9 §09 for the series of cos and sin and its (A3) to rearrange a convergent series into its even and odd parts,

$$e^{t\mathbf{A}} = (\cos t)\mathbf{I} + (\sin t)\mathbf{A} = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix},$$

a real orthogonal matrix. Hence $\|e^{t\mathbf{A}}\|_2 = 1$ for every t , while $\|t\mathbf{A}\|_2 = t$ and the bound reads e^t . At $t = 10$ the bound is about 22026 and the truth is 1.

Warning. The bound sees only the size of $\mathbf{A}$, never the direction of its spectrum.

Part (b) above is the extreme case: $\|e^{t\mathbf{A}}\|$ stays at 1 while $e^{t\|\mathbf{A}\|}$ runs away. The reason is that Theorem 15.6.1 throws away every cancellation in the series at the first step, when $\|\mathbf{A}^m\|$ is replaced by $\|\mathbf{A}\|^m$. Any statement about $e^{t\mathbf{A}}$ for large t must come from the eigenvalues instead, which is the subject of the next subsection.

Quick check. Why does Theorem 15.6.1 (a) fail for the Frobenius norm, and what is the correct statement for it?

Solution. Because $\|\mathbf{I}_n\|_F = \sqrt{n}$, so already at $\mathbf{A} = 0$ the claim reads $\sqrt{n} \leq 1$. Repeating the proof with the term $m = 0$ kept separate gives the correct version for any matrix norm: $\|e^{\mathbf{A}}\| \leq \|\mathbf{I}\| + e^{\|\mathbf{A}\|} - 1$. This is the same correction as in Exercise (C3: Without $\|\mathbf{I}\| = 1$), and it comes from the same place: only an induced norm has $\|\mathbf{I}\| = 1$.

15.6.2 Decay in the stable case

The warning above says the bound $e^{t\|\mathbf{A}\|}$ is the wrong instrument for large t . The right one reads the eigenvalues, and the bridge to the discrete theory of §04 is the observation that the exponential turns the *real part* of an eigenvalue into a *modulus*.

Lemma 15.6.2. The Spectrum of the Exponential

Let $\mathbf{A} \in M_n(\mathbb{C})$. Then $\text{spec}(e^{\mathbf{A}}) = \{e^\lambda : \lambda \in \text{spec}(\mathbf{A})\}$, and

$$\rho(e^{\mathbf{A}}) = e^\mu, \quad \mu = \max\{\text{Re } \lambda : \lambda \in \text{spec}(\mathbf{A})\}.$$

Idea. The eigenvalues of $e^{\mathbf{A}}$ are the numbers e^λ for $\lambda \in \text{spec}(\mathbf{A})$, which the Jordan form makes visible. Taking moduli turns e^λ into $e^{\text{Re}\lambda}$, and the largest of finitely many such numbers is the one with the largest exponent.

Proof. By Theorem 9.3.1 there are an invertible $\mathbf{P}$ and a Jordan matrix $\mathbf{J}$ with $\mathbf{A} = \mathbf{PJP}^{-1}$, and the diagonal of $\mathbf{J}$ lists the eigenvalues $\lambda_1, \dots, \lambda_n$ of $\mathbf{A}$ with multiplicity. By Theorem 9.9.4 (a), $e^{\mathbf{J}} = e^{\mathbf{P}^{-1}\mathbf{A}\mathbf{P}} = \mathbf{P}^{-1}e^{\mathbf{A}}\mathbf{P}$. By Theorem 9.9.3 each block $e^{\mathbf{J}_k(\lambda)}$ is upper triangular with e^λ on its diagonal, so $e^{\mathbf{J}}$ is upper triangular with diagonal $e^{\lambda_1}, \dots, e^{\lambda_n}$; hence $\text{spec}(e^{\mathbf{J}}) = \{e^{\lambda_i}\}$ by Theorem 8.7.4 (b). Similar matrices have the same characteristic polynomial (Theorem 6.10.3), so $\text{spec}(e^{\mathbf{A}}) = \text{spec}(e^{\mathbf{J}})$, which is the first claim.

For the second, $|e^\lambda| = e^{\text{Re}\lambda}$ by fact (A1) of Chapter 9 §09, and $s \mapsto e^s$ is non-decreasing on $\mathbb{R}$ — for $h \geq 0$, $e^{s+h} = e^s e^h \geq e^s$ by fact (A1) of Chapter 9 §09 — so the largest of the finitely many numbers $e^{\text{Re}\lambda_i}$ is e^μ . This proves the lemma. ■

Chapter 11 §10 called $\mathbf{A}$ **stable** when every $\lambda \in \text{spec}(\mathbf{A})$ has $\text{Re}\lambda < 0$ (Definition 11.10.2). In the language of the lemma, $\mathbf{A}$ is stable exactly when $\rho(e^{\mathbf{A}}) < 1$ — the continuous-time condition and Chapter 9 §10's discrete-time condition are the same condition, read through the exponential.

Theorem 15.6.3. Exponential Decay of a Stable System

Let $\mathbf{A} \in M_n(\mathbb{C})$, let $\mu = \max\{\text{Re}\lambda : \lambda \in \text{spec}(\mathbf{A})\}$, and let $\nu > \mu$ be real. Then there is a constant $C \geq 1$, depending on $\mathbf{A}$, on ν and on the norm, with

$$\|e^{t\mathbf{A}}\| \leq C e^{\nu t} \quad \text{for every } t \geq 0.$$

In particular, if $\mathbf{A}$ is **stable** then ν may be taken negative, and every solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$ satisfies

$$\|\mathbf{x}(t)\| \leq C e^{\nu t} \|\mathbf{x}(0)\| \xrightarrow[t \rightarrow \infty]{} 0.$$

Idea. Sample the flow at integer times and use §04. The matrix $\mathbf{M} = e^{-\nu}\mathbf{A}$ has $\rho(\mathbf{M}) = e^{\mu-\nu} < 1$ by Lemma 15.6.2, so Corollary 15.4.5 bounds all of its powers by one constant; unwinding the powers gives $\|e^{k\mathbf{A}}\| \leq C_0 e^{\nu k}$ at every integer k . Between the integers, $e^{t\mathbf{A}} = e^{(t-k)\mathbf{A}} e^{k\mathbf{A}}$ with $0 \leq t-k < 1$, and the crude bound Theorem 15.6.1 (a) is more than enough to cover a gap of length 1. The whole point is that the crude bound is applied only on a bounded interval, where crudeness costs a constant and not a rate.

Proof. Put $\mathbf{M} = e^{-\nu}\mathbf{A}$. Multiplying a matrix by a scalar c multiplies every eigenvalue by c , so by Lemma 15.6.2,

$$\rho(\mathbf{M}) = e^{-\nu}\rho(\mathbf{A}) = e^{-\nu}e^\mu = e^{\mu-\nu} < 1,$$

using $\mu - \nu < 0$. Fix any q with $\rho(\mathbf{M}) < q < 1$. By Corollary 15.4.5 (b) there is $C_0 \geq 1$ with $\|\mathbf{M}^k\| \leq C_0 q^k \leq C_0$ for every integer $k \geq 0$.

Since $\mathbf{A}$ commutes with itself, Theorem 9.9.4 (d) gives $e^{(s+u)\mathbf{A}} = e^{s\mathbf{A}}e^{u\mathbf{A}}$ for all real s, u ; in particular $(e^{\mathbf{A}})^k = e^{k\mathbf{A}}$ by induction on k . Hence $\mathbf{M}^k = e^{-\nu k}e^{k\mathbf{A}}$, and

$$\|e^{k\mathbf{A}}\| = e^{\nu k}\|\mathbf{M}^k\| \leq C_0 e^{\nu k} \quad (k = 0, 1, 2, \dots).$$

Now let $t \geq 0$ and let k be the integer part of t , so that $k \geq 0$ and $0 \leq t - k < 1$. Then $e^{t\mathbf{A}} = e^{(t-k)\mathbf{A}}e^{k\mathbf{A}}$, so by submultiplicativity and Theorem 15.6.1 (a),

$$\|e^{t\mathbf{A}}\| \leq \|e^{(t-k)\mathbf{A}}\| \|e^{k\mathbf{A}}\| \leq e^{(t-k)\|\mathbf{A}\|} C_0 e^{\nu k} \leq C_0 e^{\|\mathbf{A}\|} e^{\nu k},$$

using $0 \leq t - k < 1$ in the last step. Finally $e^{\nu k} = e^{\nu t} e^{-\nu(t-k)} \leq e^{\nu t} e^{|\nu|}$, again because $0 \leq t - k < 1$. So the constant

$$C = C_0 e^{\|\mathbf{A}\| + |\nu|} \geq 1$$

works for every $t \geq 0$.

For the last statement, suppose $\mathbf{A}$ is stable, so $\mu < 0$, and choose $\nu \in (\mu, 0)$. By Theorem 9.9.6 the unique solution with the given initial value is $\mathbf{x}(t) = e^{t\mathbf{A}}\mathbf{x}(0)$, so $\|\mathbf{x}(t)\| \leq \|e^{t\mathbf{A}}\| \|\mathbf{x}(0)\| \leq C e^{\nu t} \|\mathbf{x}(0)\|$ by Theorem 15.3.2 (a). Since $\nu < 0$, $e^{\nu t} \rightarrow 0$ as $t \rightarrow \infty$ — this is fact (A5) of Chapter 9 §09 with $j = 0$, quoted there and quoted again here — so $\|\mathbf{x}(t)\| \rightarrow 0$, and by Corollary 15.2.4 the vectors $\mathbf{x}(t)$ tend to $\mathbf{0}$. This proves the theorem. ■

Example. Decay after a detour

Let $\mathbf{A} = \begin{pmatrix} -1 & 5 \\ 0 & -1 \end{pmatrix}$. Find μ , compute $\|e^{t\mathbf{A}}\|_\infty$ exactly, and find the smallest constant C that works in Theorem 15.6.3 for $\nu = -\frac{1}{2}$.

Solution. The eigenvalues. $\mathbf{A}$ is upper triangular with both diagonal entries -1 , so $\text{spec}(\mathbf{A}) = \{-1\}$ (Theorem 8.7.4 (b)) and $\mu = -1 < 0$: the matrix is stable.

The exponential. Write $\mathbf{A} = -\mathbf{I} + 5\mathbf{N}$ with $\mathbf{N} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$. The two summands commute and $\mathbf{N}^2 = 0$, so Corollary 9.9.5 gives

$$e^{t\mathbf{A}} = e^{-t}(\mathbf{I} + 5t\mathbf{N}) = e^{-t} \begin{pmatrix} 1 & 5t \\ 0 & 1 \end{pmatrix}.$$

The absolute row sums are $e^{-t}(1 + 5t)$ and e^{-t} , so $\|e^{t\mathbf{A}}\|_\infty = e^{-t}(1 + 5t)$ for $t \geq 0$ by Theorem 15.3.3 (b).

The transient. The function $t \mapsto e^{-t}(1 + 5t)$ has derivative $e^{-t}(4 - 5t)$, so it **ris**es from 1 at $t = 0$ to $5e^{-4/5} \approx 2.247$ at $t = \frac{4}{5}$, and only then decays. A stable matrix does not decay from the start.

The constant. We need the smallest C with $e^{-t}(1 + 5t) \leq C e^{-t/2}$ for all $t \geq 0$, that is, $C = \max_{t \geq 0} (1 + 5t)e^{-t/2}$. The derivative of $(1 + 5t)e^{-t/2}$ is $e^{-t/2}(5 - \frac{1}{2}(1 + 5t))$, which is positive for $t < \frac{9}{5}$ and negative for $t > \frac{9}{5}$, so by fact (A6) of the chapter introduction the maximum is at $t = \frac{9}{5}$ and

$$C = 10 e^{-9/10} \approx 4.066.$$

The rate $e^{-t/2}$ is honest and the constant pays for the hump.

Remark. What this settles from Chapter 11 §10, and what it does not. That section introduced the Lyapunov equation $\mathbf{A}\mathbf{X} + \mathbf{X}\mathbf{A}^* = -\mathbf{Q}$ and the word *stable* (Definition 11.10.2). It then stated, without proof, that a stable $\mathbf{A}$ and a positive definite $\mathbf{Q}$ produce a positive definite solution $\mathbf{X}$, and conversely; it assigned the positivity half to Chapter 12, and assigned to this chapter “the consequence for differential equations — that a stable $\mathbf{A}$ makes every solution of $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$ decay, with $\mathbf{x}^*\mathbf{X}\mathbf{x}$ as the quantity that measures the decay”.

[Theorem 15.6.3](#) delivers the decay, with a rate, and does so without the Lyapunov equation at all. What it does not deliver is the existence of a positive definite $\mathbf{X}$ for a stable $\mathbf{A}$: that statement is not proved anywhere in this book, and nothing here depends on it. What can be said cheaply is the other half of the connection: **any** positive definite $\mathbf{X}$ satisfying $\mathbf{A}^*\mathbf{X} + \mathbf{X}\mathbf{A} = -\mathbf{Q}$ with $\mathbf{Q} \succ 0$ does measure the decay, in the sense that $\mathbf{x}^*\mathbf{X}\mathbf{x}$ is strictly decreasing along every non-zero solution of $\mathbf{x}' = \mathbf{A}\mathbf{x}$. That is [Exercise \(C3: A quantity that measures the decay\)](#), and it is three lines once the product rule below is available. Note the transpose: the equation the decay computation wants is Definition 11.10.2 for the pair $(\mathbf{A}^*, \mathbf{Q})$, and $\mathbf{A}$ is stable if and only if $\mathbf{A}^*$ is, since the eigenvalues of $\mathbf{A}^*$ are the conjugates of those of $\mathbf{A}$.

15.6.3 Differentiating a matrix-valued function

A matrix whose entries vary — a family $\mathbf{A}(t)$, a curve in $M_n(F)$ — is differentiated entrywise, exactly as Chapter 9 §09 differentiated $t \mapsto e^{t\mathbf{A}}$. The two derivatives this book needs later, of $\det$ and of inversion, are algebraic facts with one analytic ingredient, and the ingredient is named first.

Definition 15.6.1. Derivative of a Matrix-Valued Function

Let $I \subseteq \mathbb{R}$ be an open interval and $\mathbf{A}: I \rightarrow M_{m \times n}(\mathbb{C})$, written $\mathbf{A}(t) = (a_{ij}(t))$. Call $\mathbf{A}$ **differentiable** at $t_0 \in I$ if every entry function $a_{ij}: I \rightarrow \mathbb{C}$ is differentiable at t_0 , and then set

$$\mathbf{A}'(t_0) := (a'_{ij}(t_0)) \in M_{m \times n}(\mathbb{C}).$$

The prime here always means the derivative in t ; no dual map (Definition 4.4.1) appears anywhere in this section, so the notation cannot collide. By Corollary 15.2.4 the definition can be restated without coordinates: $\mathbf{A}$ is differentiable at t_0 with derivative $\mathbf{B}$ exactly when

$$\left\| \frac{\mathbf{A}(t) - \mathbf{A}(t_0)}{t - t_0} - \mathbf{B} \right\| \rightarrow 0 \quad (t \rightarrow t_0),$$

in any norm at all. Both readings are used below, the entrywise one to import calculus and the norm one to take limits.

Remark. The calculus this section quotes. Let $f, g: I \rightarrow \mathbb{C}$ be differentiable at t_0 .

(D1) (*Linearity and the product rule.*) $f + g$ and fg are differentiable at t_0 , with $(f + g)' = f' + g'$ and $(fg)' = f'g + fg'$; a constant function has derivative 0.

(D2) (*Differentiable implies continuous.*) f is continuous at t_0 .

Neither is proved here. (D1) is Chapter 9 §09's fact (A3), read for complex-valued functions of a real variable; (D2) is one more standard fact of the same kind, quoted here for the first time in this book. Both are of a piece with Chapter 13 §06's Taylor assumption: standard one-variable calculus, quoted so that the theorems below have hypotheses to stand on. Everything that follows them is linear algebra.

Lemma 15.6.4. Product Rule for Matrices

Let $\mathbf{A}: I \rightarrow M_{m \times n}(\mathbb{C})$ and $\mathbf{B}: I \rightarrow M_{n \times p}(\mathbb{C})$ be differentiable at t_0 . Then so is $t \mapsto \mathbf{A}(t)\mathbf{B}(t)$, with

$$(\mathbf{AB})'(t_0) = \mathbf{A}'(t_0)\mathbf{B}(t_0) + \mathbf{A}(t_0)\mathbf{B}'(t_0).$$

Proof. Fix a position (i, k) . The entry function is $t \mapsto \sum_{j=1}^n a_{ij}(t)b_{jk}(t)$, a finite sum of products of functions differentiable at t_0 . By (D1) it is differentiable at t_0 with derivative

$$\sum_{j=1}^n (a'_{ij}(t_0)b_{jk}(t_0) + a_{ij}(t_0)b'_{jk}(t_0)),$$

which is the (i, k) -entry of $\mathbf{A}'(t_0)\mathbf{B}(t_0) + \mathbf{A}(t_0)\mathbf{B}'(t_0)$. Since this holds at every position, [Definition 15.6.1](#) gives the claim. Note that the order of the factors is preserved throughout: matrices do not commute, and the product rule must not be written $2\mathbf{AA}'$ even for $\mathbf{B} = \mathbf{A}$. This proves the lemma. ■

15.6.4 The derivative of the inverse

Nothing so far says that $t \mapsto \mathbf{A}(t)^{-1}$ is even defined near t_0 , let alone differentiable. Both come from §05.

Theorem 15.6.5. Derivative of the Inverse

Let $\mathbf{A}: I \rightarrow M_n(\mathbb{C})$ be differentiable at $t_0 \in I$, with $\mathbf{A}(t_0)$ invertible. Then $\mathbf{A}(t)$ is invertible for all t in some open interval around t_0 , the function $t \mapsto \mathbf{A}(t)^{-1}$ is differentiable at t_0 , and

$$(\mathbf{A}^{-1})'(t_0) = -\mathbf{A}(t_0)^{-1} \mathbf{A}'(t_0) \mathbf{A}(t_0)^{-1}.$$

Idea. The formula is what one gets by differentiating $\mathbf{AA}^{-1} = \mathbf{I}$ with [Lemma 15.6.4](#) and solving — but that argument presupposes what has to be proved, namely that $\mathbf{A}^{-1}$ is differentiable. So run it in the honest order. The difference of two inverses factors as $\mathbf{X}^{-1} - \mathbf{Y}^{-1} = \mathbf{X}^{-1}(\mathbf{Y} - \mathbf{X})\mathbf{Y}^{-1}$; divide by $t - t_0$ and the middle factor becomes a difference quotient, which converges by hypothesis, while the outer factor converges because inversion is continuous (Corollary 15.5.2). The limit of the product is the product of the limits, and that is the formula.

Proof. Write $\mathbf{A}_0 = \mathbf{A}(t_0)$. By (D2) every entry of $\mathbf{A}$ is continuous at t_0 , so $\mathbf{A}(t) \rightarrow \mathbf{A}_0$ entrywise as $t \rightarrow t_0$, hence $\|\mathbf{A}(t) - \mathbf{A}_0\| \rightarrow 0$ by Corollary 15.2.4. By Corollary 15.5.2 there is therefore an open interval $J \ni t_0$ on which $\|\mathbf{A}(t) - \mathbf{A}_0\| < 1/\|\mathbf{A}_0^{-1}\|$, so that $\mathbf{A}(t)$ is invertible for $t \in J$, and on which $\|\mathbf{A}(t)^{-1} - \mathbf{A}_0^{-1}\| \rightarrow 0$ as $t \rightarrow t_0$.

Let $t \in J$ with $t \neq t_0$. Multiplying $\mathbf{A}_0 - \mathbf{A}(t)$ on the left by $\mathbf{A}(t)^{-1}$ and on the right by $\mathbf{A}_0^{-1}$,

$$\mathbf{A}(t)^{-1} - \mathbf{A}_0^{-1} = -\mathbf{A}(t)^{-1}(\mathbf{A}(t) - \mathbf{A}_0)\mathbf{A}_0^{-1}.$$

Divide by $t - t_0$:

$$\frac{\mathbf{A}(t)^{-1} - \mathbf{A}_0^{-1}}{t - t_0} = -\mathbf{A}(t)^{-1} \frac{\mathbf{A}(t) - \mathbf{A}_0}{t - t_0} \mathbf{A}_0^{-1}.$$

As $t \rightarrow t_0$ the middle factor tends to $\mathbf{A}'(t_0)$, by Definition 15.6.1. The left factor tends to $\mathbf{A}_0^{-1}$, by the first paragraph. Products of convergent matrix families converge to the product of the limits: if $\|\mathbf{X}(t) - \mathbf{X}\| \rightarrow 0$ and $\|\mathbf{Y}(t) - \mathbf{Y}\| \rightarrow 0$ then

$$\|\mathbf{X}(t)\mathbf{Y}(t) - \mathbf{X}\mathbf{Y}\| \leq \|\mathbf{X}(t) - \mathbf{X}\|\|\mathbf{Y}(t)\| + \|\mathbf{X}\|\|\mathbf{Y}(t) - \mathbf{Y}\|,$$

and $\|\mathbf{Y}(t)\|$ is bounded near t_0 because $\|\mathbf{Y}(t)\| \leq \|\mathbf{Y}\| + \|\mathbf{Y}(t) - \mathbf{Y}\|$. Applying this twice, the right-hand side converges to $-\mathbf{A}_0^{-1}\mathbf{A}'(t_0)\mathbf{A}_0^{-1}$. Hence the difference quotient on the left converges, which by Definition 15.6.1 is exactly the differentiability of $t \mapsto \mathbf{A}(t)^{-1}$ at t_0 , with the stated derivative. This proves the theorem. ■

Warning. The two copies of $\mathbf{A}^{-1}$ sit on either side and cannot be combined. The scalar formula $(1/a)' = -a'/a^2$ tempts one to write $-\mathbf{A}^{-2}\mathbf{A}'$, and that is wrong unless $\mathbf{A}'(t_0)$ commutes with $\mathbf{A}(t_0)$. For $\mathbf{A}(t) = \begin{pmatrix} 1 & t \\ 0 & 2 \end{pmatrix}$ at $t = 0$: we have $\mathbf{A}(0)^{-1} = \text{diag}(1, \frac{1}{2})$ and $\mathbf{A}'(0) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, so the correct answer is $-\text{diag}(1, \frac{1}{2})\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}\text{diag}(1, \frac{1}{2}) = \begin{pmatrix} 0 & -1/2 \\ 0 & 0 \end{pmatrix}$, whereas $-\mathbf{A}(0)^{-2}\mathbf{A}'(0) = -\begin{pmatrix} 1 & 0 \\ 0 & 1/4 \end{pmatrix}\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 0 & 0 \end{pmatrix}$.

15.6.5 Jacobi's formula

The determinant is a polynomial in the entries, so differentiating it needs no analysis beyond (D1); the work is in recognizing the answer.

Theorem 15.6.6. Jacobi's Formula

Let $\mathbf{A}: I \rightarrow M_n(\mathbb{C})$ be differentiable at $t_0 \in I$. Then $t \mapsto \det \mathbf{A}(t)$ is differentiable at t_0 , with

$$\left. \frac{d}{dt} \right|_{t_0} \det \mathbf{A}(t) = \text{tr}(\text{adj } \mathbf{A}(t_0) \mathbf{A}'(t_0)).$$

If moreover $\mathbf{A}(t_0)$ is invertible, this equals $\det \mathbf{A}(t_0) \text{tr}(\mathbf{A}(t_0)^{-1} \mathbf{A}'(t_0))$.

Idea. Use the Leibniz formula, which writes $\det$ as a sum of $n!$ products of n entries, one from each column. The product rule differentiates each such product into n terms, the

j -th of which has column j differentiated and the rest untouched. Collecting by j gives n determinants, the j -th being $\mathbf{A}$ with its j -th column replaced by $\mathbf{a}'_j$. Expanding that determinant along column j (Theorem 6.6.1) produces $\sum_i a'_{ij} C_{ij}$, and summing over j gives a double sum which is precisely a trace against the adjugate.

Proof. For $n = 1$ the statement reads $\frac{d}{dt}\big|_{t_0} a_{11}(t) = a'_{11}(t_0)$, since $\text{adj } \mathbf{A} = (1)$ by Definition 6.6.2, so assume $n \geq 2$; that is what lets us speak of cofactors. Write $\mathbf{A}(t) = (a_{ij}(t))$ with columns $\mathbf{a}_1(t), \dots, \mathbf{a}_n(t)$, and write C_{ij} for the cofactors of $\mathbf{A}(t_0)$ (Definition 6.6.1). By Theorem 6.4.3,

$$\det \mathbf{A}(t) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{\sigma(1)1}(t) \cdots a_{\sigma(n)n}(t),$$

a finite sum of products of n functions each differentiable at t_0 . By (D1), applied $n - 1$ times to each product and then to the sum, $\det \mathbf{A}$ is differentiable at t_0 with

$$\frac{d}{dt}\bigg|_{t_0} \det \mathbf{A} = \sum_{j=1}^n \sum_{\sigma \in S_n} \text{sgn}(\sigma) a'_{\sigma(j)j}(t_0) \prod_{l \neq j} a_{\sigma(l)l}(t_0),$$

where the outer sum collects, for each j , the terms in which the j -th factor was the one differentiated. By Theorem 6.4.3 read backwards, the inner sum is the determinant of the matrix $\mathbf{A}_j$ obtained from $\mathbf{A}(t_0)$ by replacing its j -th column with $\mathbf{a}'_j(t_0)$. Expanding $\det \mathbf{A}_j$ along that column with Theorem 6.6.1 (a), and using that the cofactors of column j do not involve column j and are therefore the cofactors C_{ij} of $\mathbf{A}(t_0)$ itself,

$$\det \mathbf{A}_j = \sum_{i=1}^n a'_{ij}(t_0) C_{ij}.$$

Summing over j ,

$$\frac{d}{dt}\bigg|_{t_0} \det \mathbf{A} = \sum_{j=1}^n \sum_{i=1}^n C_{ij} a'_{ij}(t_0).$$

It remains to recognize the right-hand side. By Definition 6.6.2, $(\text{adj } \mathbf{A}(t_0))_{ji} = C_{ij}$, so

$$\text{tr}(\text{adj } \mathbf{A}(t_0) \mathbf{A}'(t_0)) = \sum_{j=1}^n \sum_{i=1}^n (\text{adj } \mathbf{A}(t_0))_{ji} a'_{ij}(t_0),$$

which is the same double sum. For the last claim, if $\mathbf{A}(t_0)$ is invertible then $\text{adj } \mathbf{A}(t_0) = \det \mathbf{A}(t_0) \mathbf{A}(t_0)^{-1}$ by Theorem 6.6.2, and the trace is linear (Theorem 0.9.6), so the scalar $\det \mathbf{A}(t_0)$ comes out in front. This proves the theorem. ■

Corollary 15.6.7. The Determinant's Derivative at the Identity Is the Trace

For every $\mathbf{B} \in M_n(\mathbb{C})$,

$$\frac{d}{dt}\bigg|_{t=0} \det(\mathbf{I} + t\mathbf{B}) = \text{tr } \mathbf{B}.$$

Proof. Apply [Theorem 15.6.6](#) to $\mathbf{A}(t) = \mathbf{I} + t\mathbf{B}$, which is differentiable with $\mathbf{A}'(t) = \mathbf{B}$ and $\mathbf{A}(0) = \mathbf{I}$. Since $\mathbf{I}$ is invertible with $\mathbf{I}^{-1} = \mathbf{I}$ and $\det \mathbf{I} = 1$, the second form of the formula gives $\operatorname{tr}(\mathbf{I}\mathbf{B}) = \operatorname{tr} \mathbf{B}$. ■

So the determinant, which is a product of n numbers, has the trace, a sum of n numbers, as its linearization at $\mathbf{I}$. That is the honest source of the identity $\det e^{\mathbf{A}} = e^{\operatorname{tr} \mathbf{A}}$ of Theorem 9.9.4 (e), and [Exercise \(C1: The determinant of the exponential, again\)](#) reproves that identity along these lines.

Example. Both derivatives on one family

Let $\mathbf{A}(t) = \begin{pmatrix} 1+t & t^2 \\ t & 1 \end{pmatrix}$.

- (a) Compute $\frac{d}{dt} \det \mathbf{A}(t)$ directly and by [Theorem 15.6.6](#).
- (b) Compute $(\mathbf{A}^{-1})'(0)$ by [Theorem 15.6.5](#), and check it against a direct expansion.

Solution.

- (a) *Directly.* $\det \mathbf{A}(t) = (1+t) \cdot 1 - t^2 \cdot t = 1+t-t^3$, so the derivative is $1-3t^2$.

By the formula. $\operatorname{adj} \mathbf{A}(t) = \begin{pmatrix} 1 & -t^2 \\ -t & 1+t \end{pmatrix}$ (Definition 6.6.2) and $\mathbf{A}'(t) = \begin{pmatrix} 1 & 2t \\ t & 0 \end{pmatrix}$, so

$$\operatorname{adj} \mathbf{A}(t) \mathbf{A}'(t) = \begin{pmatrix} 1-t^2 & 2t \\ 1 & -2t^2 \end{pmatrix},$$

whose trace is $1-t^2-2t^2 = 1-3t^2$. The two agree.

- (b) $\mathbf{A}(0) = \mathbf{I}$ and $\mathbf{A}'(0) = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$, so [Theorem 15.6.5](#) gives

$$(\mathbf{A}^{-1})'(0) = -\mathbf{I} \begin{pmatrix} 1 & 0 \\ 1 & 0 \end{pmatrix} \mathbf{I} = \begin{pmatrix} -1 & 0 \\ -1 & 0 \end{pmatrix}.$$

Check. $\mathbf{A}(t)^{-1} = \frac{1}{1+t-t^3} \begin{pmatrix} 1 & -t^2 \\ -t & 1+t \end{pmatrix}$. Near $t=0$ the scalar factor is $1-t+(\text{terms in } t^2)$, so

$$\mathbf{A}(t)^{-1} = \begin{pmatrix} 1-t & 0 \\ -t & 1 \end{pmatrix} + (\text{terms in } t^2),$$

whose derivative at 0 is $\begin{pmatrix} -1 & 0 \\ -1 & 0 \end{pmatrix}$, as predicted. The formula did the work that the quotient rule would otherwise have had to do four times.

Quick check. [Theorem 15.6.6](#) holds with no invertibility hypothesis, yet the second form of the formula has one. Where exactly is invertibility used, and what does the formula say at a t_0 where $\det \mathbf{A}(t_0) = 0$?

Solution. Only in the last line, where $\text{adj } \mathbf{A}(t_0)$ is rewritten as $\det \mathbf{A}(t_0) \mathbf{A}(t_0)^{-1}$. The adjugate itself is defined for every square matrix (Definition 6.6.2), so the first form survives. At a t_0 with $\det \mathbf{A}(t_0) = 0$ it still gives a number: for $\mathbf{A}(t) = \text{diag}(t, t)$ at $t_0 = 0$ we get $\text{adj } \mathbf{A}(0) = 0$, so the derivative of $\det \mathbf{A}(t) = t^2$ is 0 there, which is correct. The second form would read “0 times something undefined”.

15.6.6 Exercises

15.6.6.1 A. Check your understanding

Exercise. A1

- State the three bounds of [Theorem 15.6.1](#), and say which property of the norm each one needs.
- Determine whether the following statement is correct, and justify your answer: if $\|\mathbf{A}\|_2$ is large then $\|e^{\mathbf{A}}\|_2$ is large.
- State [Theorem 15.6.3](#), and say what the constant C is paying for.
- State Jacobi’s formula ([Theorem 15.6.6](#)), and name the fact from calculus its proof quotes.
- Write down the derivative of $t \mapsto \mathbf{A}(t)^{-1}$ and explain in one sentence why it is not $-\mathbf{A}^{-2} \mathbf{A}'$.

Solution.

- $\|e^{\mathbf{A}}\| \leq e^{\|\mathbf{A}\|}$, $\|e^{\mathbf{A}} - \mathbf{I}\| \leq e^{\|\mathbf{A}\|} - 1$, and $\|e^{\mathbf{A}} - \mathbf{I} - \mathbf{A}\| \leq e^{\|\mathbf{A}\|} - 1 - \|\mathbf{A}\| \leq \|\mathbf{A}\|^2 e^{\|\mathbf{A}\|} / 2$. All three need subadditivity (to bound a sum term by term) and submultiplicativity (to get $\|\mathbf{A}^m\| \leq \|\mathbf{A}\|^m$); the first also needs $\|\mathbf{I}\| = 1$, which is what makes the $m = 0$ term equal 1.
- Incorrect. For $\mathbf{A} = t \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ we have $\|\mathbf{A}\|_2 = t$, arbitrarily large, while $e^{\mathbf{A}}$ is a rotation matrix and $\|e^{\mathbf{A}}\|_2 = 1$ ([Example \(Where the bound is exact, and where it is useless\)](#)).
- If μ is the largest real part of an eigenvalue of $\mathbf{A}$ and $\nu > \mu$, there is $C \geq 1$ with $\|e^{t\mathbf{A}}\| \leq C e^{\nu t}$ for all $t \geq 0$. The constant pays for the transient: $\|e^{t\mathbf{A}}\|$ may first grow, as in [Example \(Decay after a detour\)](#), and it also pays for the gap between ν and μ .
- $\frac{d}{dt} \det \mathbf{A}(t) = \text{tr}(\text{adj } \mathbf{A}(t) \mathbf{A}'(t))$, for a matrix-valued $\mathbf{A}$ differentiable at t . The quoted fact is (D1): the linearity of differentiation and the product rule for complex-valued functions of a real variable.
- $(\mathbf{A}^{-1})' = -\mathbf{A}^{-1} \mathbf{A}' \mathbf{A}^{-1}$. The two copies of $\mathbf{A}^{-1}$ sit on opposite sides of $\mathbf{A}'$, and matrices do not commute, so they may not be collected into $\mathbf{A}^{-2}$ unless $\mathbf{A}'$ commutes with $\mathbf{A}$.

15.6.6.2 B. Practice

Exercise. B1: Bound against truth

For each matrix, compute $\|e^{\mathbf{A}}\|_{\infty}$ exactly and compare with $e^{\|\mathbf{A}\|_{\infty}}$.

- (a) $\mathbf{A} = \begin{pmatrix} 0 & 3 \\ 0 & 0 \end{pmatrix}$.
 (b) $\mathbf{A} = \text{diag}(1, -4)$.

Solution.

- (a) $\mathbf{A}^2 = 0$, so the series stops: $e^{\mathbf{A}} = \mathbf{I} + \mathbf{A} = \begin{pmatrix} 1 & 3 \\ 0 & 1 \end{pmatrix}$, with $\|e^{\mathbf{A}}\|_{\infty} = 4$. Also $\|\mathbf{A}\|_{\infty} = 3$, so the bound is $e^3 \approx 20.09$. Correct, and loose by a factor of about 5.
 (b) $e^{\mathbf{A}} = \text{diag}(e, e^{-4})$, so $\|e^{\mathbf{A}}\|_{\infty} = e \approx 2.718$. Here $\|\mathbf{A}\|_{\infty} = 4$ and the bound is $e^4 \approx 54.6$. The bound is governed by the entry of largest modulus, -4 , while the exponential is governed by the largest **real part**, 1. This is the discrepancy [Theorem 15.6.3](#) exists to remove.

Exercise. B2: A rate for a stable matrix

Let $\mathbf{A} = \begin{pmatrix} -2 & 4 \\ 0 & -2 \end{pmatrix}$.

- (a) Compute μ and $e^{t\mathbf{A}}$, and give $\|e^{t\mathbf{A}}\|_{\infty}$ for $t \geq 0$.
 (b) Find the largest value of $\|e^{t\mathbf{A}}\|_{\infty}$ on $t \geq 0$.
 (c) Find the smallest C that works in [Theorem 15.6.3](#) for $\nu = -1$.

Solution.

- (a) $\mathbf{A}$ is upper triangular with diagonal $-2, -2$, so $\text{spec}(\mathbf{A}) = \{-2\}$ (Theorem 8.7.4 (b)) and $\mu = -2$. Writing $\mathbf{A} = -2\mathbf{I} + 4\mathbf{N}$ with $\mathbf{N} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$, the summands commute and $\mathbf{N}^2 = 0$, so by Corollary 9.9.5

$$e^{t\mathbf{A}} = e^{-2t} \begin{pmatrix} 1 & 4t \\ 0 & 1 \end{pmatrix}, \quad \|e^{t\mathbf{A}}\|_{\infty} = e^{-2t}(1 + 4t).$$

- (b) The derivative of $e^{-2t}(1 + 4t)$ is $e^{-2t}(4 - 2 - 8t) = e^{-2t}(2 - 8t)$, which is positive for $t < \frac{1}{4}$ and negative after, so by fact (A6) the maximum is at $t = \frac{1}{4}$. The value there is $2e^{-1/2} \approx 1.213$.
 (c) We need the smallest C with $e^{-2t}(1 + 4t) \leq Ce^{-t}$, that is, $C = \max_{t \geq 0} (1 + 4t)e^{-t}$. Its derivative is $e^{-t}(4 - 1 - 4t) = e^{-t}(3 - 4t)$, positive for $t < \frac{3}{4}$ and negative after, so by fact (A6) the maximum is at $t = \frac{3}{4}$ and $C = 4e^{-3/4} \approx 1.890$.

Exercise. B3: Two derivatives

Let $\mathbf{A}(t) = \begin{pmatrix} \cos t & -\sin t \\ \sin t & \cos t \end{pmatrix}$.

- (a) Compute $\frac{d}{dt} \det \mathbf{A}(t)$ by [Theorem 15.6.6](#) and check it against the exact value of $\det \mathbf{A}(t)$.
 (b) Compute $(\mathbf{A}^{-1})'(t)$ by [Theorem 15.6.5](#) and check it by differentiating $\mathbf{A}(t)^{-1}$ directly.

Solution.

- (a) $\det \mathbf{A}(t) = \cos^2 t + \sin^2 t = 1$, so the derivative is 0. By the formula: $\operatorname{adj} \mathbf{A}(t) = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix}$ and $\mathbf{A}'(t) = \begin{pmatrix} -\sin t & -\cos t \\ \cos t & -\sin t \end{pmatrix}$, so

$$\operatorname{adj} \mathbf{A}(t) \mathbf{A}'(t) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

whose trace is 0. The two agree. (The diagonal entries are $-\cos t \sin t + \sin t \cos t$ and $\sin t \cos t - \cos t \sin t$, both zero.)

- (b) Since $\det \mathbf{A}(t) = 1$, $\mathbf{A}(t)^{-1} = \operatorname{adj} \mathbf{A}(t) = \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix}$, whose entrywise derivative is $\begin{pmatrix} -\sin t & \cos t \\ -\cos t & -\sin t \end{pmatrix}$. By the formula, using the product $\operatorname{adj} \mathbf{A}(t) \mathbf{A}'(t)$ computed in (a),

$$-\mathbf{A}^{-1} \mathbf{A}' \mathbf{A}^{-1} = -\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \cos t & \sin t \\ -\sin t & \cos t \end{pmatrix} = \begin{pmatrix} -\sin t & \cos t \\ -\cos t & -\sin t \end{pmatrix}.$$

The two agree.

15.6.6.3 C. Going deeper

Exercise. C1: The determinant of the exponential, again

Let $\mathbf{A} \in M_n(\mathbb{C})$ and put $\varphi(t) = \det e^{t\mathbf{A}}$.

- (a) Prove that φ is differentiable with $\varphi'(t) = (\operatorname{tr} \mathbf{A})\varphi(t)$.
 (b) Hence deduce $\det e^{\mathbf{A}} = e^{\operatorname{tr} \mathbf{A}}$, recovering Theorem 9.9.4 (e).

Hint for (b): consider $\psi(t) = e^{-t \operatorname{tr} \mathbf{A}} \varphi(t)$.

Solution.

- (a) The function $t \mapsto e^{t\mathbf{A}}$ is differentiable with derivative $\mathbf{A}e^{t\mathbf{A}}$ (Theorem 9.9.4 (b)), and $e^{t\mathbf{A}}$ is invertible with inverse $e^{-t\mathbf{A}}$ (Theorem 9.9.4 (c)). So Theorem 15.6.6 applies in its second form:

$$\varphi'(t) = \det(e^{t\mathbf{A}}) \operatorname{tr}(e^{-t\mathbf{A}} \mathbf{A} e^{t\mathbf{A}}).$$

By Theorem 0.9.6 (3) with the two factors $e^{-t\mathbf{A}}$ and $\mathbf{A}e^{t\mathbf{A}}$, we get $\operatorname{tr}(e^{-t\mathbf{A}} \mathbf{A} e^{t\mathbf{A}}) = \operatorname{tr}(\mathbf{A} e^{t\mathbf{A}} e^{-t\mathbf{A}}) = \operatorname{tr} \mathbf{A}$, giving $\varphi'(t) = (\operatorname{tr} \mathbf{A})\varphi(t)$.

- (b) Put $c = \operatorname{tr} \mathbf{A}$ and $\psi(t) = e^{-ct} \varphi(t)$. By the product rule (D1) and (a),

$$\psi'(t) = -c e^{-ct} \varphi(t) + e^{-ct} c \varphi(t) = 0$$

for every t , where $\frac{d}{dt} e^{-ct} = -c e^{-ct}$ is Theorem 9.9.4 (b) in the case $n = 1$. A differentiable function with vanishing derivative is constant — fact (A4) of Chapter 9 §09 — so $\psi(t) = \psi(0) = \det(e^0) = \det \mathbf{I} = 1$. Hence $\varphi(t) = e^{ct}$, and $t = 1$ gives $\det e^{\mathbf{A}} = e^{\operatorname{tr} \mathbf{A}}$.

Exercise. C2: Comparing two exponentials

Let $\mathbf{A}, \mathbf{B} \in M_n(\mathbb{C})$ with $\mathbf{AB} = \mathbf{BA}$.

- (a) Prove that $\|e^{\mathbf{A}} - e^{\mathbf{B}}\| \leq e^{\|\mathbf{B}\|}(e^{\|\mathbf{A}-\mathbf{B}\|} - 1)$.
- (b) Deduce that $\mathbf{M} \mapsto e^{\mathbf{M}}$ is continuous at every $\mathbf{M}$ that commutes with a neighborhood of itself, and identify those $\mathbf{M}$: they are exactly the scalar matrices $c\mathbf{I}$.
- (c) Say exactly where the commuting hypothesis is used, and why the argument cannot simply be dropped.

Solution.

- (a) Since $\mathbf{A}$ and $\mathbf{B}$ commute, so do $\mathbf{B}$ and $\mathbf{A}-\mathbf{B}$. By Theorem 9.9.4 (d), $e^{\mathbf{A}} = e^{\mathbf{B}+(\mathbf{A}-\mathbf{B})} = e^{\mathbf{B}}e^{\mathbf{A}-\mathbf{B}}$. Hence

$$e^{\mathbf{A}} - e^{\mathbf{B}} = e^{\mathbf{B}}(e^{\mathbf{A}-\mathbf{B}} - \mathbf{I}),$$

and submultiplicativity with Theorem 15.6.1 (a) and (b) gives

$$\|e^{\mathbf{A}} - e^{\mathbf{B}}\| \leq \|e^{\mathbf{B}}\| \|e^{\mathbf{A}-\mathbf{B}} - \mathbf{I}\| \leq e^{\|\mathbf{B}\|}(e^{\|\mathbf{A}-\mathbf{B}\|} - 1).$$

- (b) Take $\mathbf{B} = c\mathbf{I}$, which commutes with every matrix. For $\mathbf{A}$ with $\|\mathbf{A} - c\mathbf{I}\| = \delta$, part (a) gives $\|e^{\mathbf{A}} - e^{c\mathbf{I}}\| \leq e^{|c|}(e^{\delta} - 1)$, and $e^{\delta} - 1 \rightarrow 0$ as $\delta \rightarrow 0$, because $t \mapsto e^t$ is differentiable by fact (A2) of Chapter 9 §09 and so continuous at 0. So $e^{\mathbf{A}} \rightarrow e^{c\mathbf{I}}$ as $\mathbf{A} \rightarrow c\mathbf{I}$.
- (c) Only in the first line, where $e^{\mathbf{B}+(\mathbf{A}-\mathbf{B})}$ is split as $e^{\mathbf{B}}e^{\mathbf{A}-\mathbf{B}}$; Theorem 9.9.4 (d) has commutation as a hypothesis, and Chapter 9 §09 exhibits matrices with $e^{\mathbf{A}+\mathbf{B}} \neq e^{\mathbf{A}}e^{\mathbf{B}}$. Without it there is no factorization to take norms of, and a different argument is needed. (The inequality itself is in fact true in general, but proving it needs an integral representation of $e^{\mathbf{A}} - e^{\mathbf{B}}$, which this book does not have.)

Exercise. C3: A quantity that measures the decay

Let $\mathbf{A}, \mathbf{X}, \mathbf{Q} \in M_n(\mathbb{C})$ with $\mathbf{X}$ and $\mathbf{Q}$ positive definite (Definition 12.1.1) and

$$\mathbf{A}^*\mathbf{X} + \mathbf{X}\mathbf{A} = -\mathbf{Q}.$$

Let $\mathbf{x}: \mathbb{R} \rightarrow \mathbb{C}^n$ be differentiable with $\mathbf{x}' = \mathbf{A}\mathbf{x}$, and put $V(t) = \mathbf{x}(t)^*\mathbf{X}\mathbf{x}(t)$.

- (a) Prove that V is real-valued and differentiable, with $V'(t) = \mathbf{x}(t)^*(\mathbf{A}^*\mathbf{X} + \mathbf{X}\mathbf{A})\mathbf{x}(t)$.
- (b) Deduce that if $\mathbf{x}(0) \neq \mathbf{0}$, then V is strictly decreasing.
- (c) Explain how this relates to the Lyapunov equation of Definition 11.10.2, whose unknown sits between $\mathbf{A}$ and $\mathbf{A}^*$ the other way round.

Hint for (a): $\mathbf{x}^$ is a $1 \times n$ matrix-valued function, and $\overline{f'} = \overline{f'}$ for a differentiable $f: \mathbb{R} \rightarrow \mathbb{C}$.*

Solution.

- (a) $V(t)$ is a 1×1 matrix, and $V(t)^* = \mathbf{x}^*\mathbf{X}^*\mathbf{x} = \mathbf{x}^*\mathbf{X}\mathbf{x} = V(t)$ since $\mathbf{X}$ is Hermitian, so $V(t)$ is real. Each entry of $t \mapsto \mathbf{x}(t)^*$ is the conjugate of an entry of $\mathbf{x}$, hence differentiable with $(\mathbf{x}^*)' = (\mathbf{x}')^* = (\mathbf{A}\mathbf{x})^* = \mathbf{x}^*\mathbf{A}^*$. By Lemma 15.6.4, applied twice to the product $\mathbf{x}^* \cdot \mathbf{X} \cdot \mathbf{x}$ and using $\mathbf{X}' = 0$,

$$V'(t) = \mathbf{x}^*\mathbf{A}^*\mathbf{X}\mathbf{x} + \mathbf{x}^*\mathbf{X}\mathbf{A}\mathbf{x} = \mathbf{x}^*(\mathbf{A}^*\mathbf{X} + \mathbf{X}\mathbf{A})\mathbf{x}.$$

- (b) By hypothesis $V'(t) = -\mathbf{x}(t)^*\mathbf{Q}\mathbf{x}(t)$. If $\mathbf{x}(t_1) = \mathbf{0}$ for some t_1 , then by Theorem 9.9.6

$\mathbf{x}(t) = e^{t\mathbf{A}}\mathbf{x}(0)$ for every t , and $e^{t_1\mathbf{A}}$ is invertible by Theorem 9.9.4 (c), so $\mathbf{x}(0) = \mathbf{0}$, contradicting the hypothesis. So $\mathbf{x}(t) \neq \mathbf{0}$ for every t , and $\mathbf{Q} \succ 0$ gives $\mathbf{x}(t)^*\mathbf{Q}\mathbf{x}(t) > 0$, hence $V'(t) < 0$ for every t . A differentiable real function with everywhere negative derivative is strictly decreasing; that last step is the mean value theorem, one more fact of one-variable calculus quoted here and not proved.

- (c) Definition 11.10.2 writes the equation as $\mathbf{A}\mathbf{X} + \mathbf{X}\mathbf{A}^* = -\mathbf{Q}$; the equation used here is that one for the pair $(\mathbf{A}^*, \mathbf{Q})$. The two are interchanged by replacing $\mathbf{A}$ with $\mathbf{A}^*$, which changes nothing about stability, since the eigenvalues of $\mathbf{A}^*$ are the complex conjugates of those of $\mathbf{A}$ and conjugation preserves real parts. So the decay of V is exactly the statement Chapter 11 §10 pointed forward to, for whichever of the two conventions one adopts — granted a positive definite solution $\mathbf{X}$, whose existence is not proved in this book.