

22.5 Burnside and the Double Commutant

Twice already the book has computed a commutant: Chapter 9 §04 found the matrices commuting with a basic Weyr matrix, and Chapter 11 §10 found the matrices commuting with a block diagonal matrix whose blocks have separated spectra. Both times the question was *given one operator, who commutes with it?* This section turns the question round. Given a whole algebra of operators, how much of the algebra can be read off from its commutant, and when is the algebra as large as it could possibly be? Two theorems answer this. Burnside's theorem says that over an algebraically closed field an algebra with no invariant subspace to hide in must be everything; the double commutant theorem says that a semisimple algebra is exactly the set of operators commuting with everything that commutes with it.

Throughout, V is a finite-dimensional vector space over a field F , and $\text{End}(V)$ is the space $\mathcal{L}(V)$ of operators regarded as an algebra under composition. A **subalgebra** $A \subseteq \text{End}(V)$ contains id_V by (SA3) of Definition 22.1.1; this is not decoration, and the warning after [Theorem 22.5.5](#) shows what goes wrong without it.

22.5.1 Commutants, and when an operator is a scalar

For a subalgebra $A \subseteq \text{End}(V)$, Definition 22.1.6 sets

$$A' = \{t \in \text{End}(V) : ta = at \text{ for all } a \in A\},$$

and Proposition 22.1.4 says A' is again a subalgebra containing id_V .

Applying the prime twice gives the **double commutant** $A'' := (A')'$. One inclusion is free and worth recording at once: by Proposition 22.1.4 (c),

$$A \subseteq A''.$$

The content of this section is that the reverse inclusion holds under a hypothesis we can name exactly.

Both theorems below end by proving that some operator is a scalar, and §01 has already supplied the fact that lets us conclude this. By Proposition 22.1.5 the center of $M_n(F)$ consists of the scalar matrices or, in the operator form recorded there, for a finite-dimensional $V \neq \{0\}$ the operators commuting with every operator on V are exactly the scalar multiples of id_V :

$$\text{End}(V)' = \{\lambda \text{id}_V : \lambda \in F\}.$$

Nothing new is proved here; it is the operator form that the proofs below call on.

22.5.2 Algebras that act irreducibly

Recall from §03 that V is **irreducible** as an A -space (§03 calls such a space simple, Definition 22.3.3) when $V \neq \{0\}$ and the only A -invariant subspaces of V are $\{0\}$ and V . The following one-line consequence is used in every proof of this section, so we isolate it.

Lemma 22.5.1. An Irreducible Algebra Moves One Vector Everywhere

Let $A \subseteq \text{End}(V)$ be a subalgebra with $\text{id}_V \in A$, acting irreducibly on V . Then $Av := \{av :$

$a \in A\} = V$ for every $\mathbf{v} \neq \mathbf{0}$.

Proof. The set $A\mathbf{v}$ is the image of the linear map $A \rightarrow V$, $a \mapsto a\mathbf{v}$, hence a subspace. It is A -invariant: $b(a\mathbf{v}) = (ba)\mathbf{v} \in A\mathbf{v}$ for $b \in A$, since $ba \in A$. It contains $\text{id}_V \mathbf{v} = \mathbf{v} \neq \mathbf{0}$, so it is not $\{\mathbf{0}\}$. Irreducibility leaves only $A\mathbf{v} = V$. ■

Warning. The assumption $\text{id}_V \in A$ is doing real work here. Take $V = F$, a line, and $A = \{0\}$, a subspace closed under composition but not a subalgebra, since it does not contain id_V . The only subspaces of V at all are $\{0\}$ and V , so A satisfies the letter of irreducibility, yet $A\mathbf{v} = \{0\} \neq V$ and $A \neq \text{End}(V) = F$. Every statement below about an algebra acting irreducibly assumes the identity belongs to the algebra.

22.5.3 Burnside's theorem

Here is the question. An algebra $A \subseteq \text{End}(V)$ that acts irreducibly has nowhere to hide: no proper non-zero subspace survives it. Does that force A to be all of $\text{End}(V)$? Over $\mathbb{C}$ the answer is yes. The proof is worth the trouble: the corollary below, which says that an irreducibly acting set of operators already spans $\text{End}(V)$, is what the representation theory later in this chapter uses to turn statements about a group into statements about all matrices.

Theorem 22.5.2. Burnside's Theorem

Let F be an **algebraically closed** field, that is, one in which every non-constant polynomial in $F[x]$ has a root. Let V be a vector space over F with $1 \leq \dim V < \infty$, and let $A \subseteq \text{End}(V)$ be a subalgebra with $\text{id}_V \in A$. If V is irreducible as an A -space, then

$$A = \text{End}(V).$$

Idea. Step roadmap. ① Among the non-zero members of A , let r be the smallest rank, attained at a . ② Show $r = 1$. If $r \geq 2$, the image $\text{im } a$ has two independent vectors $a\mathbf{u}_1, a\mathbf{u}_2$, and irreducibility supplies $b \in A$ carrying $a\mathbf{u}_1$ to $\mathbf{u}_2$. The operator ab maps $\text{im } a$ into itself; an eigenvalue λ of that restriction — this is the one place algebraic closure is spent — makes $aba - \lambda a$ a member of A of strictly smaller rank, hence zero, and then $a\mathbf{u}_2 = \lambda a\mathbf{u}_1$ contradicts independence. ③ One rank-one operator $\mathbf{v}_0\varphi_0$ in A gives all of them: multiplying on the left moves $\mathbf{v}_0$ anywhere, and multiplying on the right moves φ_0 anywhere. ④ Rank-one operators span $\text{End}(V)$.

Proof. Write $\mathbf{v}\varphi$ for the operator $\mathbf{x} \mapsto \varphi(\mathbf{x})\mathbf{v}$, where $\mathbf{v} \in V$ and $\varphi \in V^*$; it has rank 1 exactly when $\mathbf{v} \neq \mathbf{0}$ and $\varphi \neq 0$, and every operator of rank at most 1 has this form, since an operator with image $\text{span}(\mathbf{v})$, $\mathbf{v} \neq \mathbf{0}$, sends each $\mathbf{x}$ to a unique multiple $\varphi(\mathbf{x})\mathbf{v}$, and φ so defined is linear.

Step 1: a member of least rank. Since $\text{id}_V \in A$ and $\dim V \geq 1$, the set $\{a \in A : a \neq 0\}$

is non-empty. Let

$$r = \min\{\text{rank } a : a \in A, a \neq 0\} \geq 1,$$

and fix $a \in A$ with $a \neq 0$ and $\text{rank } a = r$.

Step 2: $r = 1$. Suppose $r \geq 2$. Then $\dim \text{im } a \geq 2$, so there are $\mathbf{u}_1, \mathbf{u}_2 \in V$ with $a\mathbf{u}_1$ and $a\mathbf{u}_2$ independent; in particular $a\mathbf{u}_1 \neq \mathbf{0}$. By Lemma 22.5.1 there is $b \in A$ with $b(a\mathbf{u}_1) = \mathbf{u}_2$.

Put $W = \text{im } a$, a subspace of dimension r . Then $ab(W) \subseteq \text{im } a = W$, so $s := (ab)|_W$ is an operator on W . Its characteristic polynomial p_s is monic of degree $r \geq 1$, hence non-constant, hence has a root $\lambda \in F$ because F is algebraically closed; by Theorem 8.2.1, λ is an eigenvalue of s , so $\ker(s - \lambda \text{id}_W) \neq \{\mathbf{0}\}$.

Consider $c := aba - \lambda a$, which lies in A because A is closed under products and linear combinations. For every $\mathbf{x} \in V$,

$$c\mathbf{x} = (ab - \lambda \text{id}_V)(a\mathbf{x}) = (s - \lambda \text{id}_W)(a\mathbf{x}),$$

the second equality because $a\mathbf{x} \in W$. Hence $\text{im } c = (s - \lambda \text{id}_W)(W)$, and Theorem 3.3.1 applied to $s - \lambda \text{id}_W$ on W gives

$$\text{rank } c = r - \dim \ker(s - \lambda \text{id}_W) \leq r - 1.$$

By the minimality of r , an element of A of rank less than r must be 0, so $c = 0$, that is, $aba = \lambda a$. Applying both sides to $\mathbf{u}_1$ and using $ba\mathbf{u}_1 = \mathbf{u}_2$,

$$a\mathbf{u}_2 = (aba)\mathbf{u}_1 = \lambda a\mathbf{u}_1,$$

which contradicts the independence of $a\mathbf{u}_1$ and $a\mathbf{u}_2$. Therefore $r = 1$.

Step 3: A contains every operator of rank at most one. By Step 2 there is $a_0 \in A$ with $\text{rank } a_0 = 1$; write $a_0 = \mathbf{v}_0\varphi_0$ with $\mathbf{v}_0 \neq \mathbf{0}$ and $\varphi_0 \neq 0$.

Left multiplication. For $b \in A$ and $\mathbf{x} \in V$ we get $ba_0\mathbf{x} = \varphi_0(\mathbf{x})b\mathbf{v}_0$, that is, $ba_0 = (b\mathbf{v}_0)\varphi_0$. By Lemma 22.5.1, $A\mathbf{v}_0 = V$, so

$$\mathbf{v}\varphi_0 \in A \quad \text{for every } \mathbf{v} \in V.$$

Right multiplication. For $c \in A$ we get $a_0c = \mathbf{v}_0(\varphi_0 \circ c)$. Put

$$S = \{\varphi_0 \circ c : c \in A\} \subseteq V^*,$$

the image of the linear map $A \rightarrow V^*$, $c \mapsto \varphi_0 \circ c$, and hence a subspace of V^* .

Claim. $S = V^*$.

Proof. Let $U = \{\mathbf{x} \in V : \varphi(\mathbf{x}) = 0 \text{ for every } \varphi \in S\}$, a subspace of V . It is A -invariant: if $\mathbf{x} \in U$ and $d \in A$, then for every $c \in A$ we have $cd \in A$ and so $\varphi_0(c(d\mathbf{x})) = \varphi_0((cd)\mathbf{x}) = 0$, whence $d\mathbf{x} \in U$. It is not all of V : taking $c = \text{id}_V$ shows $\varphi_0 \in S$, and $\varphi_0 \neq 0$ gives some $\mathbf{x}$ with $\varphi_0(\mathbf{x}) \neq 0$. Irreducibility therefore forces $U = \{\mathbf{0}\}$.

Let $d = \dim S$ and pick a basis $\varphi_1, \dots, \varphi_d$ of S . The linear map $\Theta: V \rightarrow F^d$, $\Theta(\mathbf{x}) = (\varphi_1(\mathbf{x}), \dots, \varphi_d(\mathbf{x}))$, has kernel exactly $U = \{\mathbf{0}\}$, since a vector killed by a basis of S is killed by all of S . So Θ is injective and Theorem 3.3.1 gives $\dim V \leq d$. On the other hand S is a subspace of V^* , and $\dim V^* = \dim V$ by Corollary 4.1.4, so $d \leq \dim V$. Hence $d = \dim V^*$ and $S = V^*$ by Theorem 1.6.8.

Now let $\mathbf{v} \in V$ and $\varphi \in V^*$ be arbitrary. Choose $c \in A$ with $\varphi = \varphi_0 \circ c$, which the Claim allows, and $b \in A$ with $b\mathbf{v}_0 = \mathbf{v}$, which Lemma 22.5.1 allows when $\mathbf{v} \neq \mathbf{0}$ (and for $\mathbf{v} = \mathbf{0}$ take $b = 0$). Then for every $\mathbf{x}$,

$$(b a_0 c)\mathbf{x} = b(\varphi_0(c\mathbf{x}) \mathbf{v}_0) = \varphi(\mathbf{x}) \mathbf{v},$$

so $\mathbf{v}\varphi = ba_0c \in A$.

Step 4: rank-one operators span. Let $t \in \text{End}(V)$, let $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of V and $(\varphi_1, \dots, \varphi_n)$ its dual basis (Definition 4.1.3). The operator $\sum_{i=1}^n (t\mathbf{v}_i)\varphi_i$ sends $\mathbf{v}_j$ to $t\mathbf{v}_j$ for each j , because $\varphi_i(\mathbf{v}_j) = \delta_{ij}$; two operators agreeing on a basis are equal, so $t = \sum_i (t\mathbf{v}_i)\varphi_i$. Each summand lies in A by Step 3, and A is a subspace, so $t \in A$. This proves $A = \text{End}(V)$.

The theorem is stronger than it may look. It says that irreducibility, a statement about *subspaces*, forces a statement about *dimension*: over an algebraically closed field, an irreducibly acting subalgebra of $\text{End}(V)$ has dimension exactly $(\dim V)^2$, the largest possible. There is no room between “acts irreducibly” and “is everything”.

The version about sets rather than algebras is worth recording alongside it, since a group of invertible operators is a set closed under composition and not a subspace.

Corollary 22.5.3. An Irreducible Multiplicative Set Spans the Whole Algebra

Let F be algebraically closed, V a vector space over F with $1 \leq \dim V < \infty$, and let $\mathcal{S} \subseteq \text{End}(V)$ be a set with $\text{id}_V \in \mathcal{S}$ that is closed under composition. If the only subspaces $U \subseteq V$ with $s(U) \subseteq U$ for all $s \in \mathcal{S}$ are $\{\mathbf{0}\}$ and V , then $\text{span}(\mathcal{S}) = \text{End}(V)$.

Proof. Put $A = \text{span}(\mathcal{S})$. It is a subspace containing id_V , and it is closed under multiplication: a product of two linear combinations of members of $\mathcal{S}$ expands, by bilinearity of composition, into a linear combination of products of members of $\mathcal{S}$, which lie in $\mathcal{S}$. So A is a subalgebra. A subspace U is A -invariant if and only if it is invariant under every member of $\mathcal{S}$, since invariance is preserved by linear combinations. So V is irreducible as an A -space, and Theorem 22.5.2 gives $A = \text{End}(V)$.

Warning. Burnside is false over a field that is not algebraically closed. Let $F = \mathbb{R}$, $V = \mathbb{R}^2$ and

$$A = \left\{ \begin{pmatrix} x & -y \\ y & x \end{pmatrix} : x, y \in \mathbb{R} \right\},$$

the span of $\mathbf{I}_2$ and $\mathbf{Q} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Since $\mathbf{Q}^2 = -\mathbf{I}_2$, this is a subalgebra, and it acts irreducibly: an invariant line would be spanned by a real eigenvector of $\mathbf{Q}$, and $p_{\mathbf{Q}} = x^2 + 1$ has no real root. Yet $\dim A = 2$, not 4, so $A \neq M_2(\mathbb{R})$. This is the same witness §04 used against the second half of Schur’s lemma (Theorem 22.4.1), and the reason is the same: an

operator on a real space need have no eigenvalue, and Step 2 of the proof needs one.

Quick check. Let $A \subseteq M_2(\mathbb{C})$ be the subalgebra generated by $\mathbf{X} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $\mathbf{Z} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Without computing products, what must $\dim A$ be, and why?

Solution. It must be 4, so $A = M_2(\mathbb{C})$. A one-dimensional invariant subspace would be a common eigenvector of $\mathbf{X}$ and $\mathbf{Z}$. The eigenvectors of $\mathbf{Z}$ are the multiples of $\mathbf{e}_1$ and of $\mathbf{e}_2$, and $\mathbf{X}\mathbf{e}_1 = \mathbf{e}_2$ is not a multiple of $\mathbf{e}_1$, nor is $\mathbf{X}\mathbf{e}_2 = \mathbf{e}_1$ a multiple of $\mathbf{e}_2$. So A acts irreducibly on $\mathbb{C}^2$, and Theorem 22.5.2 gives $A = M_2(\mathbb{C})$. Directly: $\mathbf{I}_2, \mathbf{X}, \mathbf{Z}$ and $\mathbf{XZ} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ are independent, which confirms it.

22.5.4 The double commutant theorem

Burnside handles the irreducible case. The general case is not “ A is everything” — a diagonal algebra is not everything — but a statement about how faithfully A is recorded by its commutant. The right hypothesis is semisimplicity of the action, and it cannot be weakened to nothing, as the warning below shows.

The engine is a single step: an operator in A'' can be matched by a member of A on **one** given vector.

Lemma 22.5.4. Matching on One Vector

Let W be a finite-dimensional vector space over F and let $B \subseteq \text{End}(W)$ be a subalgebra with $\text{id}_W \in B$ which acts **semisimply** on W : every B -invariant subspace of W has a B -invariant complement (Definition 22.3.4). Then for every $t \in B''$ and every $\mathbf{w} \in W$ there is $b \in B$ with $b\mathbf{w} = t\mathbf{w}$.

Idea. The subspace $B\mathbf{w}$ is invariant, so it has an invariant complement, and the projection p onto $B\mathbf{w}$ along that complement commutes with B . Being in B'' , the operator t commutes with p , hence cannot move $\mathbf{w}$ out of the image of p , which is $B\mathbf{w}$.

Proof. Put $U = B\mathbf{w} = \{b\mathbf{w} : b \in B\}$, a subspace of W (the image of $b \mapsto b\mathbf{w}$) which is B -invariant, since $b_1(b\mathbf{w}) = (b_1b)\mathbf{w}$ for $b_1 \in B$. By hypothesis there is a B -invariant subspace U_1 with $W = U \oplus U_1$. Let $p \in \text{End}(W)$ be the projection onto U along U_1 (Theorem 3.10.1 (b)).

$p \in B'$. Indeed, let $b \in B$ and write $\mathbf{x} = \mathbf{u} + \mathbf{u}_1$ with $\mathbf{u} \in U$, $\mathbf{u}_1 \in U_1$. Then $b\mathbf{u} \in U$ and $b\mathbf{u}_1 \in U_1$ by invariance, so $b\mathbf{x} = b\mathbf{u} + b\mathbf{u}_1$ is the decomposition of $b\mathbf{x}$, giving $p(b\mathbf{x}) = b\mathbf{u} = b(p\mathbf{x})$.

Since $t \in B''$ and $p \in B'$, we have $tp = pt$. Also $\mathbf{w} = \text{id}_W \mathbf{w} \in B\mathbf{w} = U$, so $p\mathbf{w} = \mathbf{w}$. Therefore

$$t\mathbf{w} = t(p\mathbf{w}) = p(t\mathbf{w}) \in \text{im } p = U = B\mathbf{w},$$

so $t\mathbf{w} = b\mathbf{w}$ for some $b \in B$. ■

Matching on one vector is not yet matching everywhere. The trick that upgrades it is to run the lemma not on V but on n copies of V at once, where one cleverly chosen vector carries a whole basis.

Theorem 22.5.5. Double Commutant Theorem

Let F be **any** field, let V be a finite-dimensional vector space over F , and let $A \subseteq \text{End}(V)$ be a subalgebra with $\text{id}_V \in A$. Assume that A acts **semisimply** on V : every A -invariant subspace of V has an A -invariant complement (Definition 22.3.4). Then

$$A'' = A.$$

Idea. Step roadmap. ① [Lemma 22.5.4](#) matches $t \in A''$ with some $a \in A$ on one vector. ② To match on a whole basis at once, let A act diagonally on $V^n = V \oplus \cdots \oplus V$ ($n = \dim V$) and apply the lemma there to the single vector whose n entries are the basis vectors. ③ For that we must check two things about the copied algebra $\tilde{A}$: that it still acts semisimply, which Theorem 22.3.3 supplies, and that $\tilde{t}$ lies in $\tilde{A}''$, which is a computation with the $n \times n$ array of components of an operator on V^n .

Proof. The inclusion $A \subseteq A''$ was noted above, so only $A'' \subseteq A$ needs proof. Let $t \in A''$, let $n = \dim V$; if $n = 0$ there is nothing to prove, so let $n \geq 1$.

Step 1: the copied algebra. Let $V^n = V \oplus \cdots \oplus V$ be the product of n copies of V (Definition 3.9.1), and for $s \in \text{End}(V)$ let $\tilde{s} \in \text{End}(V^n)$ be given by $\tilde{s}(\mathbf{x}_1, \dots, \mathbf{x}_n) = (s\mathbf{x}_1, \dots, s\mathbf{x}_n)$. The map $s \mapsto \tilde{s}$ is linear and satisfies $\widetilde{s_1 s_2} = \tilde{s}_1 \tilde{s}_2$ and $\text{id}_V = \text{id}_{V^n}$, so $\tilde{A} = \{\tilde{a} : a \in A\}$ is a subalgebra of $\text{End}(V^n)$ containing the identity.

Step 2: $\tilde{A}$ acts semisimply on V^n . Since A acts semisimply on V , Theorem 22.3.3 writes $V = W_1 + \cdots + W_q$ with each W_j an irreducible A -invariant subspace. For $1 \leq k \leq n$ let $\iota_k : V \rightarrow V^n$ place a vector in the k -th slot and $\mathbf{0}$ elsewhere; it is injective and satisfies $\iota_k(s\mathbf{x}) = \tilde{s} \iota_k(\mathbf{x})$. Hence $\iota_k(W_j)$ is $\tilde{A}$ -invariant, and it is irreducible, because ι_k carries the A -invariant subspaces of W_j bijectively onto the $\tilde{A}$ -invariant subspaces of $\iota_k(W_j)$. Every element of V^n is the sum of its n slots, so

$$V^n = \sum_{k=1}^n \sum_{j=1}^q \iota_k(W_j),$$

a sum of irreducible $\tilde{A}$ -invariant subspaces. By the converse half of Theorem 22.3.3, $\tilde{A}$ acts semisimply on V^n .

Step 3: $\tilde{t} \in \tilde{A}''$. Let $\pi_k : V^n \rightarrow V$ be the k -th coordinate map. Every $s \in \text{End}(V^n)$ is determined by its components $s_{kl} = \pi_k \circ s \circ \iota_l \in \text{End}(V)$, through $\pi_k(s\mathbf{x}) = \sum_l s_{kl} \mathbf{x}_l$.

Comparing components in $s\tilde{a} = \tilde{a}s$ gives

$$(s\tilde{a})_{kl} = s_{kl} a, \quad (\tilde{a}s)_{kl} = a s_{kl},$$

so $s \in \tilde{A}'$ if and only if every s_{kl} lies in A' . Now let $s \in \tilde{A}'$. Then $(\tilde{t}s)_{kl} = t s_{kl}$ and $(s\tilde{t})_{kl} = s_{kl} t$, and these agree because $s_{kl} \in A'$ and $t \in A''$. Hence $\tilde{t}$ commutes with every $s \in \tilde{A}'$, that is, $\tilde{t} \in \tilde{A}''$.

Step 4: conclude. Let $(\mathbf{v}_1, \dots, \mathbf{v}_n)$ be a basis of V and put $\tilde{\mathbf{v}} = (\mathbf{v}_1, \dots, \mathbf{v}_n) \in V^n$. By Steps 1–3 the hypotheses of [Lemma 22.5.4](#) hold for $B = \tilde{A}$ on $W = V^n$, with $\tilde{t} \in \tilde{A}''$. So there is $a \in A$ with $\tilde{a}\tilde{\mathbf{v}} = \tilde{t}\tilde{\mathbf{v}}$, that is, $a\mathbf{v}_i = t\mathbf{v}_i$ for every i . Two operators agreeing on a basis are equal, so $t = a \in A$. This proves $A'' \subseteq A$, and with the reverse inclusion, $A'' = A$. ■

Notice which hypotheses were used where. Algebraic closure was **not** used: the double commutant theorem holds over every field, unlike Burnside. Finite dimension was used for the basis in Step 4 and inside Theorem 22.3.3. Semisimplicity was used exactly once, to produce the invariant complement in [Lemma 22.5.4](#).

Warning. Neither hypothesis can simply be dropped.

Without the identity. Take $A = \{0\} \subseteq \text{End}(F^2)$, a subspace closed under composition but with no identity element of $\text{End}(F^2)$ in it. Every subspace of F^2 is A -invariant and every subspace has a complement, so the condition on invariant complements holds; but $A' = M_2(F)$ and $A'' = \{\lambda \mathbf{I}_2\}$ by Proposition 22.1.5, so $A'' \neq A$.

Without semisimplicity. Let $A \subseteq M_2(F)$ be the upper triangular matrices. The line $\text{span}(\mathbf{e}_1)$ is A -invariant, and it is the **only** invariant line: if $\mathbf{v} = (v_1, v_2)$ has $v_2 \neq 0$, then $\mathbf{E}_{12}\mathbf{v} = v_2\mathbf{e}_1$ is a non-zero multiple of $\mathbf{e}_1$ and does not lie in $\text{span}(\mathbf{v})$. A complement of $\text{span}(\mathbf{e}_1)$ would be an invariant line other than $\text{span}(\mathbf{e}_1)$, so there is none and A does not act semisimply. Here $A' = \{\lambda \mathbf{I}_2\}$: a matrix commuting with $\mathbf{E}_{11}$ and $\mathbf{E}_{22}$ is diagonal, and commuting with $\mathbf{E}_{12}$ then forces the two diagonal entries to agree. Hence $A'' = \{\lambda \mathbf{I}_2\}' = M_2(F)$, which is strictly larger than A .

Remark. Semisimplicity is sufficient but not necessary. Let $\mathbf{N} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and $A = \{x\mathbf{I}_2 + y\mathbf{N}\}$. Solving $\mathbf{XN} = \mathbf{NX}$ gives exactly the matrices $x\mathbf{I}_2 + y\mathbf{N}$, so $A' = A$ and therefore $A'' = A$ — even though A does not act semisimply, the invariant line $\text{span}(\mathbf{e}_1) = \ker \mathbf{N}$ again being the only one.

Remark. The name has two sources. In finite dimensions the result above is due to Wedderburn, and the argument through n copies of V is the finite-dimensional shadow of Jacobson's density theorem, which we do not prove. For algebras of operators on a Hilbert space there is an analytic statement of the same shape, von Neumann's double commutant theorem, which is outside this book.

22.5.5 The commutant of one operator, revisited

The two earlier computations of the book now read as computations of A' for the smallest interesting algebra: the one generated by a single operator.

Corollary 22.5.6. The Commutant of a Single Operator

Let V be a finite-dimensional vector space over F , let $T \in \mathcal{L}(V)$, and let $A = F[T] \subseteq \text{End}(V)$ be the algebra of all polynomials in T . Then

$$A' = \{S \in \mathcal{L}(V) : ST = TS\}.$$

Consequently the following two earlier results are computations of A' .

- (a) If $F = \mathbb{C}$ and the matrix of T in some basis is $\mathbf{A}_1 \oplus \cdots \oplus \mathbf{A}_r$ with the $\mathbf{A}_p$ of pairwise disjoint spectra, then in that same basis every member of A' has a block **diagonal** matrix for the partition into those block sizes (Corollary 11.10.4).
- (b) If the matrix of T in some basis is a basic Weyr matrix of Weyr structure $(w_1, \dots, w_s)$, then in that same basis every member of A' has a block **upper triangular** matrix for the partition $w_1 + \cdots + w_s$ (Proposition 9.4.6).

If moreover T is diagonalizable with distinct eigenvalues $\lambda_1, \dots, \lambda_k$ and $d_i = \dim E_{\lambda_i}(T)$, then A acts semisimply, $\dim A = k$, $\dim A' = \sum_{i=1}^k d_i^2$, and $A'' = A$.

Idea. Nothing in (a) and (b) is new: whatever commutes with T commutes with every polynomial in it, so the two earlier computations were computations of A' all along. The diagonalizable case is where the second commutant enters. The Lagrange polynomials evaluated at T are the projections onto the eigenspaces, and they turn out to be a basis of A ; a commuting operator is exactly one that preserves each eigenspace and is otherwise unconstrained, which counts A' ; and the same projections assemble an invariant complement for any invariant subspace, which is the hypothesis [Theorem 22.5.5](#) needs.

Proof. An operator commutes with every polynomial in T as soon as it commutes with T , and $T \in A$; this gives the displayed description of A' . Parts (a) and (b) are then the cited results verbatim, read for the matrix of T in the given basis.

For the last sentence, let T be diagonalizable, $E_i = E_{\lambda_i}(T)$, so $V = E_1 \oplus \cdots \oplus E_k$ by Theorem 8.4.1 (c). Let $\ell_i \in F[x]$ be the Lagrange polynomial with $\ell_i(\lambda_j) = \delta_{ij}$ (Theorem 5.6.1) and $P_i = \ell_i(T)$. Evaluating on each E_j , where T acts as λ_j , shows P_i is the projection onto E_i along the other summands. Since $T^m = \sum_i \lambda_i^m P_i$ for every $m \geq 0$, the algebra A is spanned by $P_1, \dots, P_k$, which are independent because $P_i P_j = \delta_{ij} P_i$ and each $P_i \neq 0$: applying $\sum_i c_i P_i = 0$ to a non-zero vector of E_j gives $c_j = 0$. So $\dim A = k$.

By Theorem 8.9.1, every $S \in A'$ maps each E_i into itself; conversely such an S commutes with T , because both ST and TS act as $\lambda_i S$ on E_i . So A' is the set of operators preserving all the E_i , and restricting gives a linear bijection $A' \rightarrow \text{End}(E_1) \times \cdots \times \text{End}(E_k)$; hence $\dim A' = \sum_i d_i^2$. Each P_i preserves every E_j , so $P_i \in A'$.

Finally, A acts semisimply. Let U be an A -invariant subspace. Then U is T -invariant, and $T|_U$ is diagonalizable (Corollary 8.8.3), so U has a basis of eigenvectors of T (Theorem 8.4.1,

(a) $\Leftrightarrow$ (b)) and therefore $U = (U \cap E_1) \oplus \cdots \oplus (U \cap E_k)$. Choose a complement C_i of $U \cap E_i$ inside E_i for each i (Theorem 1.7.6) and put $C = C_1 \oplus \cdots \oplus C_k$. Then $V = U \oplus C$, and C is A -invariant because $P_j(C) = C_j \subseteq C$ and the P_j span A . So Theorem 22.5.5 applies and gives $A'' = A$. ■

So the diagonalizable case is an instance of Theorem 22.5.5, reached here by hand; the Weyr case in (b) is **not**, since $F[\mathbf{W}]$ does not act semisimply once $s \geq 2$. That is the honest shape of the relationship: the two earlier sections computed first commutants in cases where the second commutant theorem has nothing to say, and this section explains what extra hypothesis would make the computation reversible.

Example. The Two Extremes

Inside $M_n(F)$ with $n \geq 2$, compute A' and A'' for (i) A the diagonal matrices, and (ii) A the scalar matrices.

Solution.

- (i) Let $A = \{\text{diag}(c_1, \dots, c_n)\}$. Then $\mathbf{X}$ commutes with $\mathbf{E}_{ii}$ for every i exactly when $\mathbf{X}$ is diagonal, since $(\mathbf{X}\mathbf{E}_{ii})_{kl} = x_{ki}\delta_{il}$ and $(\mathbf{E}_{ii}\mathbf{X})_{kl} = \delta_{ik}x_{il}$, and comparing at (k, i) with $k \neq i$ gives $x_{ki} = 0$. So $A' = A$, and applying the prime again, $A'' = A' = A$. This agrees with Theorem 22.5.5: the diagonal algebra acts semisimply, F^n being the direct sum of the n coordinate lines, each of which is invariant and one-dimensional, hence irreducible.
- (ii) Let $A = \{\lambda\mathbf{I}_n\}$. Every matrix commutes with a scalar matrix, so $A' = M_n(F)$, and $A'' = M_n(F)' = \{\lambda\mathbf{I}_n\} = A$ by Proposition 22.1.5. Again $A'' = A$, as it must be: the scalars act semisimply because every subspace is invariant and every subspace has a complement.

The two examples sit at opposite ends. In (i) the commutant is as small as the algebra; in (ii) the commutant is everything. In both, applying the prime a second time returns to the start, which is what the theorem promises whenever the action is semisimple.

22.5.6 Exercises

22.5.6.1 A. Check your understanding

Exercise. A1

- (a) State Burnside's theorem, with every hypothesis.
- (b) Give the field hypothesis of Theorem 22.5.5, and say why it differs from the one in (a).
- (c) True or false, with a reason: if $A \subseteq \text{End}(V)$ is a subalgebra containing id_V , then $A''' = A'$.
- (d) Explain in one sentence why $\{\mathbf{0}\}$ and V being the only A -invariant subspaces does **not** by itself imply $A\mathbf{v} = V$ for $\mathbf{v} \neq \mathbf{0}$.

Solution.

- (a) F algebraically closed, V finite-dimensional over F with $\dim V \geq 1$, $A \subseteq \text{End}(V)$ a subalgebra containing id_V , and V irreducible as an A -space; the conclusion is $A = \text{End}(V)$.
- (b) [Theorem 22.5.5](#) has no field hypothesis at all: it holds over every field. Algebraic closure entered Burnside only to produce an eigenvalue of $(ab)|_{\text{im } a}$, and the double commutant proof never needs an eigenvalue — it needs an invariant complement, which semisimplicity hands over directly.
- (c) True, and it holds for any subset: it is Proposition 22.1.4 (d). The two ingredients are (c) of that proposition, $\mathcal{S} \subseteq \mathcal{S}''$, and (b), that $\mathcal{S} \subseteq \mathcal{T}$ implies $\mathcal{T}' \subseteq \mathcal{S}'$. Applying the prime to $A \subseteq A''$ gives $A''' \subseteq A'$, and applying the inclusion $B \subseteq B''$ to $B = A'$ gives $A' \subseteq A'''$.
- (d) Because $A\mathbf{v}$ could be $\{\mathbf{0}\}$ if A contained no operator sending $\mathbf{v}$ anywhere non-zero; the identity in A is what rules this out, as in the warning after [Lemma 22.5.1](#).

22.5.6.2 B. Practice

Exercise. B1: Determine which are irreducible

For each subalgebra of $M_2(\mathbb{C})$, determine whether it acts irreducibly on $\mathbb{C}^2$, and hence whether it equals $M_2(\mathbb{C})$. Justify your answer.

- (a) The algebra generated by $\begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$.
- (b) The algebra generated by $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and $\begin{pmatrix} 1 & 0 \\ 0 & i \end{pmatrix}$.
- (c) The algebra of all matrices $\begin{pmatrix} a & b \\ 0 & a \end{pmatrix}$.

Solution.

- (a) Not irreducible. With $\mathbf{N} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ we have $\mathbf{N}^2 = \mathbf{0}$, so the algebra is $\{x\mathbf{I}_2 + y\mathbf{N}\}$, and $\text{span}(\mathbf{e}_1) = \text{im } \mathbf{N}$ is invariant, proper and non-zero. It is not $M_2(\mathbb{C})$; its dimension is 2.
- (b) Irreducible, so it equals $M_2(\mathbb{C})$ by [Theorem 22.5.2](#). The eigenvectors of $\text{diag}(1, i)$ are the multiples of $\mathbf{e}_1$ and of $\mathbf{e}_2$ (the diagonal entries are distinct), and the swap matrix maps each of those lines to the other, so no line is invariant under both.
- (c) Not irreducible: this is the algebra of (a), written out. The invariant line is again $\text{span}(\mathbf{e}_1)$.

Exercise. B2: A commutant and a double commutant

Let $\mathbf{A} = \text{diag}(1, 1, 2) \in M_3(\mathbb{R})$ and let $A = \mathbb{R}[\mathbf{A}] \subseteq M_3(\mathbb{R})$.

- (a) Compute $\dim A$.

- (b) Describe A' explicitly and compute $\dim A'$.
- (c) Compute A'' , and check the answer against [Theorem 22.5.5](#).

Solution.

- (a) The eigenvalues are 1 and 2, each with a non-zero eigenspace, so $m_{\mathbf{A}} = (x - 1)(x - 2)$ and, by the count in [Corollary 22.5.6](#) with $k = 2$, $\dim A = 2$: a basis is $\mathbf{I}_3, \mathbf{A}$, equivalently $\mathbf{P}_1 = \text{diag}(1, 1, 0)$, $\mathbf{P}_2 = \text{diag}(0, 0, 1)$.
- (b) $E_1 = \text{span}(\mathbf{e}_1, \mathbf{e}_2)$ and $E_2 = \text{span}(\mathbf{e}_3)$. Every matrix in A' preserves both ([Theorem 8.9.1](#)), and conversely a matrix preserving both commutes with $\mathbf{A}$, since $\mathbf{A}$ acts as a scalar on each; so a matrix lies in A' exactly when it has the shape

$$\begin{pmatrix} p & q & 0 \\ u & v & 0 \\ 0 & 0 & w \end{pmatrix}.$$

Its dimension is $2^2 + 1^2 = 5$.

- (c) A matrix commuting with all of these must commute in particular with $\text{diag}(1, 1, 0)$ and $\text{diag}(0, 0, 1)$, hence be of the same block shape, and its 2×2 block must commute with all of $M_2(\mathbb{R})$, hence be scalar ([Proposition 22.1.5](#)). So $A'' = \{\alpha \mathbf{P}_1 + \beta \mathbf{P}_2\} = A$, of dimension 2. Since $\mathbf{A}$ is diagonalizable, A acts semisimply, and [Theorem 22.5.5](#) predicts exactly $A'' = A$.

Exercise. B3: One theorem, once

Let F be algebraically closed, let V be finite-dimensional over F with $\dim V \geq 2$, and let $A \subseteq \text{End}(V)$ be a **commutative** subalgebra containing id_V . Prove that A does not act irreducibly on V .

Solution. Suppose it did. By [Theorem 22.5.2](#), $A = \text{End}(V)$. But $\text{End}(V)$ is not commutative when $\dim V \geq 2$: choosing a basis and the operators with matrices $\mathbf{E}_{12}$ and $\mathbf{E}_{21}$ gives $\mathbf{E}_{12}\mathbf{E}_{21} = \mathbf{E}_{11} \neq \mathbf{E}_{22} = \mathbf{E}_{21}\mathbf{E}_{12}$. This contradicts commutativity of A , so A is not irreducible.

22.5.6.3 C. Going deeper

Exercise. C1: The rank-one step needs its hypothesis

In Step 2 of the proof of [Theorem 22.5.2](#), the operator ab was restricted to $W = \text{im } a$ and an eigenvalue was extracted.

- (a) Prove that $ab(W) \subseteq W$, and give an example of $a, b \in M_2(F)$ with $ba(\text{im } a) \not\subseteq \text{im } a$, so that the order of the product matters.
- (b) Take $F = \mathbb{R}$, $A = \{x\mathbf{I}_2 + y\mathbf{Q}\}$ with $\mathbf{Q} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Compute the minimal rank r of a non-zero member of A and confirm that the conclusion $r = 1$ fails.

- (c) Which sentence of Step 2 fails for this A ? Say exactly where the hypothesis on F was used.

Solution.

- (a) $ab(W) = a(b(W)) \subseteq \text{im } a = W$, since a is applied last. For the other order take $a = \mathbf{E}_{11}$ and $b = \mathbf{E}_{21}$ in $M_2(F)$. Then $\text{im } a = \text{span}(\mathbf{e}_1)$ and $ba\mathbf{e}_1 = \mathbf{E}_{21}\mathbf{e}_1 = \mathbf{e}_2 \notin \text{span}(\mathbf{e}_1)$.
- (b) A non-zero $x\mathbf{I}_2 + y\mathbf{Q}$ has determinant $x^2 + y^2 > 0$ for real x, y not both zero, so it is invertible and has rank 2. Hence $r = 2$, and the conclusion $r = 1$ fails; consistently, A contains no rank-one matrix and is not all of $M_2(\mathbb{R})$.
- (c) The sentence “its characteristic polynomial has a root $\lambda \in F$ because F is algebraically closed”. With $a = \mathbf{I}_2$ and $b = \mathbf{Q}$ we get $W = \mathbb{R}^2$ and $s = \mathbf{Q}$, whose characteristic polynomial $x^2 + 1$ has no real root. Everything before that sentence is valid over any field.

Exercise. C2: Commutants under a change of basis

Let V be finite-dimensional over F , let $A \subseteq \text{End}(V)$ be a subalgebra containing id_V , and let $g \in \text{End}(V)$ be invertible. Put $gAg^{-1} = \{gag^{-1} : a \in A\}$.

- (a) Prove that gAg^{-1} is a subalgebra containing id_V and that $(gAg^{-1})' = gA'g^{-1}$.
- (b) Deduce that if $A'' = A$ then $(gAg^{-1})'' = gAg^{-1}$.
- (c) Hence show that the conclusion of [Theorem 22.5.5](#) is a property of the algebra up to similarity, and use this to give, for each $n \geq 2$, a subalgebra of $M_n(F)$ with $A'' \neq A$.

Solution.

- (a) The map $a \mapsto gag^{-1}$ is linear, sends id_V to id_V , and satisfies $(ga_1g^{-1})(ga_2g^{-1}) = g(a_1a_2)g^{-1}$, so the image is a subalgebra containing the identity. For the commutant: t commutes with every gag^{-1} if and only if $g^{-1}tg$ commutes with every a , that is, if and only if $g^{-1}tg \in A'$, that is, $t \in gA'g^{-1}$.
- (b) Apply (a) twice: $(gAg^{-1})'' = (gA'g^{-1})' = gA''g^{-1} = gAg^{-1}$.
- (c) By (b), whether $A'' = A$ holds is unchanged by replacing A with a similar copy, so it is a property of the similarity class. For a failure in $M_n(F)$ take $A_n =$ all upper triangular matrices. The same computation as in the warning after [Theorem 22.5.5](#), done with the matrix units $\mathbf{E}_{ii}$ and $\mathbf{E}_{i,i+1}$, gives $A'_n = \{\lambda\mathbf{I}_n\}$, hence $A''_n = M_n(F) \neq A_n$; and by (b) every similar copy gA_ng^{-1} fails too.

Exercise. C3: Is this look-alike a theorem?

Let V be finite-dimensional over an algebraically closed F and let $A \subseteq \text{End}(V)$ be a subalgebra with $\text{id}_V \in A$.

- (a) Prove: if V is irreducible as an A -space then $A' = \{\lambda \text{id}_V\}$.
- (b) Is the converse true? That is, does $A' = \{\lambda \text{id}_V\}$ force V to be irreducible? Justify your answer.

Hint for (b): consider the upper triangular matrices in $M_2(F)$.

Solution.

- (a) By [Theorem 22.5.2](#), $A = \text{End}(V)$, so $A' = \text{End}(V)' = \{\lambda \text{id}_V\}$ by Proposition 22.1.5. (This recovers the second half of Schur's lemma, Theorem 22.4.1, in this special case.)
- (b) No. Let A be the upper triangular matrices in $M_2(F)$. The warning after [Theorem 22.5.5](#) computes $A' = \{\lambda \mathbf{I}_2\}$, yet $\text{span}(\mathbf{e}_1)$ is a proper non-zero invariant subspace, so F^2 is not irreducible as an A -space — and indeed $A \neq M_2(F)$. A trivial commutant is therefore strictly weaker than irreducibility; it is semisimplicity, not a small commutant, that makes the commutant remember the algebra.